



medicina *intensiva*

<http://www.medintensiva.org/en/>



SCIENTIFIC LETTER

Serum phosphate abnormalities in critically ill children

Alteraciones en el fósforo sérico en niños en estado crítico

Dear Editor:

Disturbances in phosphate levels are common in critically ill children,¹ and have been associated with malnutrition, refeeding syndrome,^{2–4} parenteral nutrition (PN),⁵ and the use of diuretics and vasoactive drugs.^{1,4,6} Some studies have reported an association between hypophosphatemia and hyperphosphatemia and mortality, severity status^{1,2} and the duration of admission to the Pediatric Intensive Care Unit (PICU).^{1,3}

Following approval by the Research Ethics Committee, a retrospective, single-center, observational study was carried out. The study aimed to analyze the related factors and the impact of phosphate disturbances in children admitted to the PICU.

We included patients for whom phosphate levels were measured during admission to the PICU over a period of 12 months. Patient age, sex, date of admission and discharge, anthropometric data, personal history, clinical severity scales and blood phosphate, calcium, creatinine, albumin, parathormone (PTH), magnesium and lactate values were recorded at different timepoints: T₀: day 1 and day 2; T₁: days 3–7; T₂: days 8–14; and T_{discharge}. Treatment with phosphate, drugs, blood products, renal replacement therapy (RRT), enteral nutrition (EN) or PN, invasive mechanical ventilation (IMV) or noninvasive mechanical ventilation (NIMV), extracorporeal membrane oxygenation (ECMO) and mortality were also recorded. The normal range of phosphatemia was between 4–7 mg/dl. Continuous variables were analyzed using the Mann–Whitney *U* test and the median test, categorical variables using the Chi-square test and Fisher's exact test, and correlation analysis using Pearson's correlation coefficient (*r*). Statistical significance was considered for *p* < 0.05. First, a univariate analysis was performed, followed by a multivariate logistic regression analysis of the factors related to phosphate disturbances.

Of the 370 PICU admissions, 232 patients were included in the analysis (41.8% female; see Table 1 E of the Supplementary Material). The median length of stay in the PICU was 3 days (IQR: 2–9). Ten patients died (4.3%). As in healthy children, age and serum phosphate levels showed an inverse proportional relationship. Surgical patients had higher phosphate levels upon admission than the rest of the patients (5.6 [4.7–6.8] vs. 4.4 mg/dl [3.7–5.1]; *p* < 0.001).

Fig. 1E of the Supplementary material shows the evolution of the phosphate, calcium and creatinine values, and the percentages of hypophosphatemia and hyperphosphatemia. In surgical patients, the phosphate levels were higher on admission and subsequently decreased. This pattern was not observed in medical patients (see Fig. 2E of the Supplementary material).

Tables 1 and 2 show the variables associated with hyperphosphatemia and hypophosphatemia in the univariate and multivariate analyses, respectively.

Hyperphosphatemia

As shown in Table 1, children with hyperphosphatemia at and during admission had lower ages and weights, higher severity scores and longer lengths of stay in the PICU. Fifty percent of the deceased patients developed hyperphosphatemia during their admission compared to 20.8% of the survivors (*p* = 0.041).

Hyperphosphatemia occurred more frequently in patients treated with corticosteroids (33.3% vs. 19.8%; *p* = 0.030), vasoactive drugs (36.5% vs. 10.5%; *p* < 0.001), furosemide (39.2% vs. 6.4%; *p* < 0.001), acetazolamide (50% vs. 22%; *p* = 0.017), vitamin D (47.6% vs. 21.2%; *p* = 0.006) and iron (35% vs. 20.2%; *p* = 0.043).

Multivariate analysis revealed that hyperphosphatemia at admission was associated with younger age, surgical admission and a higher Pediatric Multiple Organ Dysfunction Score (P-MODS). Hyperphosphatemia during admission was associated with younger age, surgical admission, non-invasive mechanical ventilation (NIMV), artificial nutrition, renal replacement therapy (RRT), and a longer duration of admission, as has been described in adults⁷ (Table 2).

In patients undergoing RRT, phosphorus levels decrease if dialysis fluids are not supplemented with phosphate,⁸ but our patients received phosphate supplements in their RRT fluids. Perhaps because of this, they more frequently presented hyperphosphatemia.⁶

DOI of original article:
<https://doi.org/10.1016/j.medine.2025.502217>

<https://doi.org/10.1016/j.medine.2025.502217>

2173-5727/© 2025 Elsevier España, S.L.U. and SEMICYUC. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

Please cite this article as: N. Del Amo Carramiñana, P. Lasarte Merino, C. Pascual Alonso et al., Serum phosphate abnormalities in critically ill children, *Medicina Intensiva*, <https://doi.org/10.1016/j.medine.2025.502217>

Table 1 Factors associated with phosphatemia disturbances on admission and during admission to the PICU.

	Hyperphosphatemia on admission		Hyperphosphatemia during admission		Hypophosphatemia on admission		Hypophosphatemia during admission	
	n/N (%)	p-Value	n/N (%)	p-Value	n/N (%)	p-Value	n/N (%)	p-Value
Sex								
Female	14/88 (15.9%)	0.4	22/97 (22.7%)	0.856	19/88 (21.6%)	0.866	41/97 (42.3%)	0.375
Male	15/126 (11.9%)		32/135 (23.7%)		26/126 (20.6%)		65/135 (48.1%)	
Post-surgical admission								
Yes	27/114 (23.7%)	<0.001	33/117 (28.2%)	0.073	13/114 (11.4%)	<0.001	53/117 (45.3%)	0.904
No	2/100 (2%)		21/115 (18.3%)		32/100 (32%)		53/115 (46.1%)	
IMV								
Yes	18/75 (24%)	0.001	41/86 (47.7%)	<0.01	13/75 (17.3%)	0.33	39/86 (45.3%)	0.936
No	11/139 (7.9%)		13/146 (8.9%)		32/139 (23%)		67/146 (45.9%)	
NIMV								
Yes	21/82 (25.6%)	<0.001	41/94 (43.6%)	<0.001	10/82 (12.2%)	0.012	41/94 (43.6%)	0.601
No	8/132 (6.1%)		13/138 (9.4%)		35/132 (26.5%)		65/138 (47.1%)	
Artificial nutrition								
Yes	24/87 (27.6%)	<0.001	47/102 (46.1%)	<0.001	14/87 (16.1%)	0.142	43/102 (42.2%)	0.339
No	5/127 (3.9%)		7/130 (5.4%)		31/127 (24.4%)		63/130 (48.5%)	
Parenteral nutrition								
Yes	1/21 (4.8%)	0.213	7/24 (29.2%)	0.444	8/21 (38.1%)	0.045	12/24 (50%)	0.637
No	28/192 (14.6%)		46/207 (22.2%)		37/192 (19.3%)		93/207 (44.9%)	
RRT								
Yes	6/17 (35.3%)	0.006	16/19 (84.2%)	<0.001	3/17 (17.6%)	0.721	11/19 (57.9%)	0.265
No	23/197 (11.7%)		38/213 (17.8%)		42/197 (21.3%)		95/213 (44.6%)	
ECMO								
Yes	1/9 (11.1%)	0.827	8/11 (72.7%)	<0.001	1/9 (11.1%)	0.456	5/11 (45.5%)	0.987
No	28/205 (13.7%)		46/221 (20.8%)		44/205 (21.5%)		101/221 (45.7%)	
Cardiac arrest								
Yes	2/7 (28.6%)	0.236	4/8 (50%)	0.05	2/7 (28.6%)	0.605	3/8 (37.5%)	0.66
No	26/200 (13%)		45/216 (20.8%)		41/200 (20.5%)		98/216 (45.4%)	
Mortality								
Yes	1/8 (12.5%)	0.929	5/10 (50%)	0.041	5/8 (72.5%)	0.003	3/10 (30%)	0.309
No	28/206 (13.6%)		49/222 (22.1%)		40/206 (19.4%)		103/222 (46.4%)	

ECMO: extracorporeal membrane oxygenation; CA: cardiac arrest; RRT: renal replacement therapy; PICU: Pediatric Intensive Care Unit; IMV: invasive mechanical ventilation; NIMV: noninvasive mechanical ventilation.

Table 2 Multivariate analysis of factors related to phosphate disturbances on admission and during admission to the PICU.

Variable	On admission			
	Hyperphosphatemia		Hypophosphatemia	
	OR (95%CI)	p-Value	OR (95%CI)	p-Value
Age (months)	0.956 (0.934–0.978)	<0.001	1.007 (1.001–1.012)	0.026
Surgical intervention	25.144 (3.462–182.617)	0.001	0.293 (0.129–0.666)	0.030
PRISM score III	1.061 (0.907–1.242)	0.459	1.082 (0.978–1.197)	0.127
PELOD-2 score	1.042 (0.811–1.338)	0.748	1.143 (0.939–1.392)	0.182
P-MODS	1.568 (1.019–2.411)	0.041	0.661 (0.486–0.897)	0.008
FSS score	0.662 (0.370–1.186)	0.166	0.914 (0.790–1.057)	0.223
Variable	During admission			
	Hyperphosphatemia		Hypophosphatemia	
	OR (95%CI)	p-Value	OR (95%CI)	p-Value
Age (months)	0.984 (0.974–0.993)	<0.001	1.008 (1.003–1.012)	0.001
Surgery	7.585 (2.466–23.334)	<0.001	0.981 (0.556–1.733)	0.948
Length of stay	1.026 (1.000–1.052)	0.048	1.001 (0.988–1.015)	0.840
NIMV	4.552 (1.572–13.178)	0.005	0.873 (0.450–1.694)	0.688
Artificial nutrition	4.381 (1.393–13.784)	0.012	1.100 (0.531–2.276)	0.798
Parenteral nutrition	0.109 (0.010–1.227)	0.073	0.977 (0.364–2.623)	0.963
RRT	38.973 (2.694–563.897)	0.007	1.101 (0.292–4.153)	0.887

95%CI: 95% confidence interval; Artificial nutrition: nasogastric tube/transpyloric tube/gastrostomy; OR: odds ratio; PELOD: Pediatric Logistic Organ Dysfunction; FSS: Functional Status Scale; P-MODS: Pediatric Multiple Organ Dysfunction Score; PRISM-III: Pediatric Risk of Mortality; IQR: interquartile range; RRT: renal replacement therapy; PICU: Pediatric Intensive Care Unit; NIMV: noninvasive mechanical ventilation.

Hyperphosphatemia was more frequent in patients who required mechanical ventilation, which coincides with the observations of other studies. No pathophysiological reason has been found to explain this association. Patients with hyperphosphatemia had higher severity scores, suggesting that hyperphosphatemia could be a marker of severity in critically ill children.^{6,7,9}

Hypophosphatemia

Hypophosphatemia occurred more frequently than hyperphosphatemia, which aligns with data found in the literature.⁶ The main causes of hypophosphatemia are intracellular redistribution due to respiratory alkalosis, decreased intake, or hemodilution when phosphate-free fluids are used.³

Hypophosphatemia was more prevalent among medical patients, those subjected to NIMV and PN, and those who died upon admission to the PICU. Patients presenting hypophosphatemia at and during admission were older and had a greater body weight (Table 1).

Some studies have found an association between hypophosphatemia and the duration of mechanical ventilation.^{1,6} The univariate study showed an association between hypophosphatemia upon admission and NIMV; however, this association was not confirmed in the multivariate study. We found no association between hypophosphatemia and PN, corticosteroids, furosemide or vasoactive drugs, as previously reported by other authors.^{1,10}

The univariate analysis revealed an association between hypophosphatemia at admission and mortality. A total of 72.5% of the deceased patients had hypophosphatemia at admission compared to 19.4% of the survivors; $p=0.003$. However, this association was not maintained in the multivariate analysis.

In the multivariate analysis, hypophosphatemia at admission was associated with older age, medical causes of admission and lower P-MODS scores. Hypophosphatemia during admission was only associated with older age (Table 2).

The main limitations of our study are its retrospective design, which precludes the establishment of causal relationships, and the fact that phosphate levels were not determined for some patients, which may cause the incidence of the alterations to be overestimated.

In conclusion, phosphate disturbances are frequent in children admitted to the PICU. Hyperphosphatemia was more prevalent in surgical patients, smaller patients, children subjected to NIMV, patients receiving artificial nutrition and patients with RRT. It was also associated with a longer duration of admission. Hypophosphatemia upon admission and hyperphosphatemia during admission were more prevalent in deceased patients, though a causal relationship could not be established.

CRedit authorship contribution statement

All authors have made substantial contributions in each of the following areas:

Nuria del Amo-Carramiñana, Paula Lasarte-Merino, Celia Pascual-Alonso and Rafael González-Cortés: acquisition, analysis and interpretation of the data, drafting of the article, and final approval of the manuscript.

Jesús López-Herce: conception and design of the study, acquisition, analysis and interpretation of the data, drafting of the article, critical review of the intellectual content, and the final approval of the manuscript.

Declaration of generative AI and AI-assisted technologies in the writing process

The authors declare that they have not used artificial intelligence for any activities related to the study or in the writing of the manuscript.

Financing

The authors have not received any funding to conduct this study.

Declaration of competing interest

The authors declare that they have no conflicts of interest.

Acknowledgments

We would like to acknowledge the physicians and nurses of the Department of Pediatric Intensive Care Medicine at Hospital General Universitario Gregorio Marañón in Madrid, Spain, for their help in data collection and patient management.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.medine.2025.502217>.

References

1. El Shazly AN, Soliman DR, Assar EH, Behiry EG, Gad Ahmed IAEN. Phosphate disturbance in critically ill children: Incidence, associated risk factors and clinical outcomes. *Ann Med Surg.* 2017;21:118–23.

2. De Menezes FS, Leite HP, Fernandez J, Benzecry SG, de Carvalho WB. Hypophosphatemia in children hospitalized within an intensive care unit. *J Intensive Care Med.* 2006;21:235–9.
3. Shah SK, Irshad M, Gupta N, Kabra SK, Lodha R. Hypophosphatemia in critically ill children: risk factors, outcome and mechanism. *Indian J Pediatr.* 2016;83:1379–85.
4. Santana e Meneses JF, Leite HP, de Carvalho WB, Lopes E. Hypophosphatemia in critically ill children: prevalence and associated risk factors. *Pediatr Crit Care Med.* 2009;10:234–8.
5. De Mattos Springer AM, Russo Hortencio TD, Melro EC, de Souza TH, Negrão Nogueira RJ. Hypophosphatemia in critically ill pediatric patients receiving enteral and oral nutrition. *J Parenter Enter Nutr.* 2022;46:842–9.
6. Akbaş Y, Koker A, Erkek N. Are we aware that hyperphosphatemia affects mortality and morbidity as much as hypophosphatemia in pediatric intensive care patients? *Pediatr Endocrinol Rev.* 2019;17:35–40.
7. Sin JCK, Laupland KB, Ramanan M, Tabah A. Phosphate abnormalities and outcomes among admissions to the intensive care unit: a retrospective multicentre cohort study. *J Crit Care.* 2021;64:154–9.
8. Santiago MJ, López-Herce J, Urbano J, Bellón JM, del Castillo J, Carrillo A. Hypophosphatemia and phosphate supplementation during continuous renal replacement therapy in children. *Kidney Int.* 2009;75:312–6.
9. Al Harbi SA, Al-Dorzi HM, Al Meshari AM, Tamim H, Abdukahil SAI, Sadat M, et al. Association between phosphate disturbances and mortality among critically ill patients with sepsis or septic shock. *BMC Pharmacol Toxicol.* 2021;22:30.
10. Reintam Blaser A, Gunst J, Ichai C, Casaer MP, Benstoem C, Besch G, et al. Hypophosphatemia in critically ill adults and children — a systematic review. *Clin Nutr.* 2021;40:1744–54.

Nuria Del Amo Carramiñana^a, Paula Lasarte Merino^a,
Celia Pascual Alonso^a, Jesús López-Herce^{a,b,c,*},
Rafael González Cortés^{a,b,c}

^a *Departamento de Salud Pública y Maternoinfantil, Universidad Complutense de Madrid, Madrid, Spain*

^b *Servicio de Cuidados Intensivos Pediátricos, Hospital General Universitario Gregorio Marañón, Madrid, Spain*

^c *Primary Care Interventions to Prevent Maternal and Child Chronic Diseases of Perinatal and Development Origin Network (RICORS) RD21/0012/0011, Instituto de Salud Carlos III, Madrid, Spain*

* Corresponding author.

E-mail address: pielvi@hotmail.com (J. López-Herce).