



IMAGES IN INTENSIVE MEDICINE

Pneumomediastinum radiologic signs: Naclerio's V and continuous diaphragm sign



Signos radiológicos de neumomediastino: V de Naclerio y signo de diafragma continuo

Christopher Barrera-Hoffmann^a, Yadira Mariaca-Ortiz^b, Enrique Monares-Zepeda^{c,*}

^a Critical Care Medicine, Hospital General de Zona No. 71 IMSS, Veracruz, Mexico

^b Nephrology, Instituto Nacional de Cardiología "Ignacio Chávez", Mexico City, Mexico

^c Coordination of Intensive Therapy and Hemodynamics in Secondary-Level Hospitals, IMSS Bienestar, Mexico City, Mexico




Figure 1

* Corresponding author.

E-mail address: enrique.monares@hotmail.com (E. Monares-Zepeda).




Figure 2

A previously healthy 21-year-old woman primigravida at 36 weeks of pregnancy underwent vaginal delivery of a male infant. In the immediate postpartum period, she developed subcutaneous emphysema and chest discomfort. A chest X-ray (Fig. 1) revealed subcutaneous emphysema and signs of pneumomediastinum, including Naclerio's V (blue area) and the continuous diaphragm sign (yellow lines). She was treated with conventional oxygen therapy and analgesics. A non-contrast chest CT (Fig. 2) confirmed the presence of pneumomediastinum. During her hospital stay, she was managed conservatively, with gradual improvement of symptoms and resolution of the subcutaneous emphysema. As there were no signs of recurrence, she was discharged after five days.

Funding

The authors received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflicts of interest

The authors report no conflict of interest.