



## SCIENTIFIC LETTER

### Myocardial injury as a prognostic factor in critically ill patients with severe SARS-CoV-2 pneumonia



### Daño miocárdico como factor pronóstico en neumonía grave por SARS-CoV-2

Dear Editor,

The highly contagious severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) resulted in a global outbreak of coronavirus disease 2019 (COVID-19). The manifestations of SARS-CoV-2 infection range from a total absence of symptoms to rapidly progressing life-threatening disease.<sup>1</sup> Cardiovascular complications reported in patients with COVID-19 admitted to intensive care units (ICU) include arrhythmias (16%–44%), shock (30.6%), acute heart failure (10%), myocarditis (1%), and myocardial injury (22%–33%); these complications have been associated with mortality as high as 67%.<sup>2,3</sup> Patients with COVID-19-related cardiovascular complications have elevated inflammatory markers (e.g., interleukin-6, C-reactive protein (CRP), and cardiac troponin (cTn), among others).<sup>4</sup> Notably, hs-cTn, which is indicative of myocardial injury, is associated with severe COVID-19 and worse outcomes in mixed populations of patients with severe and non-severe disease.<sup>5,6</sup> Nevertheless, the relationship between hs-cTn and mortality in ICU patients has not been firmly established. Based on current evidence, we hypothesized that myocardial injury defined as hs-cTn > 47 ng/L.<sup>7</sup> could worsen outcomes in patients with severe SARS-CoV-2 pneumonia.

This study was approved by local ethical committee at Joan XXIII University Hospital (IRB#CEIM/066/2020). Written informed consent was waived owing the nature of the study.

From a total of 181 patients with confirmed severe SARS-CoV-2 pneumonia, we analyzed 140 patients in which hs-cTn levels were available within 72 h after admission to the ICU of a university hospital in Spain between March 2020 and February 2021 (Figure 1, Supplementary material).

Table 1 reports patient's baseline characteristics.

Median hs-cTn was 11 ng/L [IQR 4.3–30.7]; hs-cTn was >47 ng/L in 30 (21.4%) patients. Compared to patients without myocardial injury, those with hs-cTn > 47 ng/L

had higher median concentrations of procalcitonin (0.25 [IQR 0.14–2.2] mg/dL vs. 0.15 [IQR 0.07–0.44] mg/dL,  $p=0.016$ ), CRP (21.8 [IQR 13.1–26.7] mg/dL vs. 12.5 [IQR 7.3–21.6] mg/dL,  $p=0.012$ ), and D-dimer (2600 [IQR 1040–15,120] ng/mL vs. 1100 [IQR 600–1800] ng/mL,  $p=0.001$ ), and a greater percentage developed shock (43.3% vs. 16.4%,  $p=0.002$ ), left ventricular systolic dysfunction (31% vs. 3.1%,  $p=0.000$ ), and acute kidney injury (43.3% vs. 24.5%,  $p<0.05$ ) (Table 1).

Of the 140 patients admitted to the ICU with severe SARS-CoV-2 pneumonia, 23 (16.4%) died within 28 days. In the bivariate analyses mortality was significantly associated with age, APACHE II, creatinine, procalcitonin, and myocardial injury (Table 1, Supplementary material). In the multivariate analysis myocardial injury was independently associated with 28-day mortality (OR: 3.60, 95%CI 1.07–12.16,  $p<0.01$ ) (Fig. 1). Cox hazard regression analysis found the risk of death was higher in patients with myocardial injury (HR = 4.56, 95% CI 1.64–12.69,  $p=0.004$ ) (Figure 2, Supplementary material).

Our results suggest that myocardial injury is common in patients with severe SARS-CoV-2 pneumonia and that early development of myocardial injury could increase the risk of death during the ICU stay.

In this sense, in a report of 138 hospitalized patients with COVID-19, Wang et al.,<sup>8</sup> found that 46% had at least one coexisting medical conditions and those admitted to the ICU were more likely to have cardiovascular complications, including myocardial injury in 22.2%. Moreover, Zhou et al.<sup>9</sup> found that patients with pre-existing cardiovascular disease had higher cardiac markers and higher rates of ICU admission and mortality.

We found that 21.4% of patients with severe COVID-19 developed myocardial injury. Myocardial injury was significantly associated with the development of cardiovascular complications like shock, left ventricular systolic dysfunction, and acute kidney injury. Like in other studies reporting strong correlations between hs-cTn and other inflammatory factors in acute infections (viral or bacterial)<sup>10</sup>, we found higher values of procalcitonin and CRP in patients with myocardial injury. These findings could be related with pro-inflammatory or pro-coagulable states and could help explain the organic damage and unfavorable prognosis.

Several studies have evaluated the relationship between myocardial injury and mortality in mixed patient populations. In this respect, a recent meta-analysis found that cardiac injury was associated with a greater than sevenfold increase in the risk of mortality, ICU admission, and severe COVID-19.<sup>6</sup> Nevertheless, few studies have evaluated the prognostic value of high-sensitivity troponin in critically ill

**Abbreviations:** SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; hs-cTn, high sensitive cardiac troponin; ICU, intensive care unit; APACHE II, Acute Physiology and Chronic Health Evaluation II score; CRP, C-reactive protein.

**Table 1** Baseline characteristics of patients with and without myocardial injury.

	Cohort population [n 140]	With myocardial injury [n 30]	Without myocardial injury [n 110]	p value
<b>Demographics</b>				
Female, n (%)	42 (30%)	10 (33.3%)	32 (29.1%)	0.653
Age, median (IQR)	66 (54–72)	65 (55.8–73)	66 (54–71.2)	0.415
BMI, median (IQR)	27.9 (26.1–32.4)	27.7 (25.9–32.2)	28.2 (26.1–32.7)	0.603
Overweight, n (%)	64 (45.7%)	15 (50%)	49 (44.5%)	0.595
Obesity, n (%)	39 (28%)	9 (30%)	30 (27.3%)	0.768
Morbid obesity, n (%)	12 (8.6%)	1 (3.3%)	11 (10%)	0.248
Hypertension, n (%)	67 (48%)	15 (50%)	52 (47.3%)	0.791
Diabetes mellitus, n (%)	33 (23.6%)	8 (23.6%)	25 (22.7%)	0.652
Cardiovascular disease	17 (12%)	6 (20%)	11 (10%)	0.137
Chronic atrial fibrillation, n (%)	6 (4.3%)	1 (3.3%)	5 (4.5%)	1
Chronic cardiac dysfunction, n (%)	2 (1.4%)	2 (6.7%)	0 (0%)	0.006
Chronic cardiac ischemia, n (%)	12 (8.6%)	5 (16.7%)	7 (6.4%)	0.074
Chronic renal disease, n (%)	7 (5%)	2 (6.7%)	5 (4.5%)	0.637
<b>Severity scores</b>				
CHARLSON, median (IQR)	3 (1–4)	3 (1–5.25)	2 (1–3)	0.1
APACHE II, median (IQR)	16 (12–20)	17.5 (14.75–21.5)	15 (12–19.75)	0.068
SAPS III, median (IQR)	52 (48–56)	52.5 (49.5–56)	52 (48–56)	0.299
SOFA 24h, median (IQR)	4 (3–6)	4 (3.75–7)	4 (3–6)	0.134
<b>Analytics</b>				
Lactate mmol/L, median (IQR)	1.8 (1.4–2.2)	1.93 (1.5–2.5)	1.78 (1.46–2.2)	0.191
Creatinine mg/dL, median (IQR)	0.8 (0.63–1)	1 (0.7–1.5)	0.77 (0.63–0.98)	0.012
PCT ng/mL, median (IQR); [n]	0.19 (0.07–0.5) [124]	0.25 (0.14–2.2) [29]	0.15 (0.07–0.44) [95]	0.012
CRP mg/dL, median (IQR)	15.5 (7.8–23)	21.8 (13.1–26.7)	12.5 (7.38–21.6)	0.016
Leukocytes × 10 <sup>3</sup> , median (IQR)	8.5 (6.5–11.4)	9 (7.2–12.4)	8.4 (6.4–11.4)	0.186
IL-6 pg/mL, median (IQR); [n]	32 (8.2–111) [82]	38.7 (4.7–135) [15]	31.5 (8.3–107) [67]	0.764
Ferritin ng/mL, median (IQR); [n]	945 (537–1650) [100]	945 (577–1792) [22]	956 (526–1552) [78]	0.934
D-dimer ng/mL, median (IQR); [n]	1.25 (0.7–2.4) [125]	2.6 (1.04–15.12) [27]	1.1 (0.6–1.8) [98]	0.001
ProBNP ng/mL, median (IQR); [n]	312 (149–32515) [40]	2287 (270–17776) [12]	235 (135–1182) [28]	0.011
hs-Tn ng/L, median (IQR)	11 (4.3–30.7) [140]	162.5 (92–673)	8.5 (4–14)	0.000
<b>Use of drugs at 3rd day</b>				
Corticoids, n (%)	95 (67.9%)	19 (63.3%)	76 (69.1%)	0.549
Noepinephrine, n (%)	100 (71.4%)	24 (80%)	76 (69.1%)	0.241
Furosemide, n (%)	58 (41.4%)	15 (50%)	43 (39.1%)	0.282
<b>Renal</b>				
Hydric balance 3d ml, mean (±SD)	–788.8 (±1925)	–413.2 (±1636)	–889.7 (±1990)	0.238
Diuresis 3d ml, mean (±SD)	4162.9 (±1338.7)	4231 (±1855)	4144 (±1172)	0.813
AKI, n (%)	39 (27.9%)	13 (43.3%)	27 (24.5%)	0.043
CRRT, n (%)	7 (5%)	3 (10%)	4 (3.6%)	0.156
<b>Hemodynamic</b>				
HR on admission, median (IQR)	79 (70–92)	84.5 (66–102.8)	110 (70.8–89)	0.239
Median BP on admission, mean (±SD)	91.3 (±17.6)	88.5 (±21.1)	92 (±16.5)	0.408
Cardiovascular complications, n (%)	122 (87.1%)			
Shock, n (%)	30 (21.4%)	13 (43.3%)	18 (16.4%)	0.002
LVD, n (%); [n]	11 (11.8%) [93]	9 (31%)	2 (3.1%)	0.000
RVD, n (%); [n]	3 (2.6%) [116]	2 (8%)	1 (1.1%)	0.117
Arrhythmias, n (%)	77 (55%)	16 (53.3%)	61 (55.5%)	0.836
Sinus bradycardia, n (%)	62 (44.3%)	10 (33.3%)	52 (47.3%)	0.173
Atrial-ventricular block, n (%)	3 (2.1%)	0 (0%)	3 (2.7%)	0.226
Atrial fibrillation, n (%)	6 (4.3%)	4 (13.3%)	2 (1.8%)	0.019
Long QT, n (%)	13 (9.3%)	4 (13.3%)	9 (8.2%)	0.389

Table 1 (Continued)

	Cohort population [n 140]	With myocardial injury [n 30]	Without myocardial injury [n 110]	p value
MI, n (%)	30 (21.4%)			
<b>Pulmonary</b>				
PaO <sub>2</sub> /FiO <sub>2</sub> at intubation, median (IQR); [n]	91 (75–115) [120]	82.5 (70–100) [28]	95 (75–120) [92]	0.207
MV, n (%)	124 (88.6%)	30 (100%)	94 (85.5%)	0.026
MV days, median (IQR); [n]	19 (9–32.3) [130]	18.5 (8.25–22)	26 (10–33) [90]	0.195
<b>Outcomes</b>				
LOS, median (IQR)	21 (11–33)	24 (15.5–32.3) [24]	14 (9–35) [97]	0.161
28-day mortality, n (%)	23 (16.4%)	9 (30%)	14 (12.7%)	0.024
ICU mortality, n (%)	36 (25.7%)	10 (38.5%)	26 (24.8%)	0.161

Data with [n] is the number of participants in the corresponding cell. BMI = body mass index, APACHE II = acute physiology and chronic evaluation score, SOFA = sequential organ failure assessment, SAPS III = simplified acute physiology score III, PCT = procalcitonin, CRP = C reactive protein, IL-6 = interleukin-6, ProBNP = n-terminal pro-B-type natriuretic peptide, hs-Tn = high sensitive cardiac troponin, AKI = acute kidney injury, CRRT = continuous renal replacement therapy, HR = heart rate, BP = blood pressure, LVD = left ventricular systolic dysfunction, RVD = right ventricular systolic dysfunction, MV = mechanical ventilation, LOS = length of stay, and ICU = intensive care unit.




Figure 1 Variables associated with 28-day mortality (logistic regression).

patients. Analyzing 111 ICU patients with confirmed COVID-19, Larcher et al.<sup>5</sup> found that hs-cTn > 22 ng/L at admission was independently associated with in-hospital mortality. Similarly, after adjusting for age, APACHE II score, creatinine, shock, and procalcitonin, we found that myocardial injury was associated with 28-day mortality.

Our study has several limitations. First, it was conducted at a single center and included only 140 patients; data from larger populations at multiple centers are needed to confirm the relation between myocardial injury and high risk of mortality. Second, logistic limitations arising from the urgency of containing the COVID-19 pandemic resulted in incomplete data about some aspects of cardiovascular complications and inflammation (e.g., echocardiography findings, troponin or interleukin-6 levels, or glucocorticoid treatment) in some cases, thus limiting our ability to identify

potential mechanisms and patterns associated with myocardial injury. Third, because we evaluated only the first three days after ICU admission, the incidence of myocardial injury and other cardiovascular complications during the course of the disease may have been underestimated; however, by excluding the possible influence of later complications, this approach might also better gauge the direct relationship between SARS-CoV-2 infection and cardiovascular complications. Finally, to attribute the death of an individual patient directly to myocardial injury would require longer follow-up.

We conclude that myocardial injury is common among patients with severe SARS-CoV-2 pneumonia admitted to ICUs and that it seems to be associated with an intense

inflammatory response as well as a higher risk of organ failure and ICU mortality.

## Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

## Conflict of interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.medin.2022.05.003](https://doi.org/10.1016/j.medin.2022.05.003).

## References

- Clerkin KJ, Fried JA, Raikhelkar J, Sayer G, Griffin JM, Masoumi A, et al. COVID-19 and cardiovascular disease. *Circulation*. 2020;141:1648–55, <http://dx.doi.org/10.1161/circulationaha.120.046941>.
- Arentz M, Yim E, Klaff L, Lokhandwala S, Riedo FX, Chong M, et al. Characteristics and outcomes of 21 critically ill patients with COVID-19 in Washington State. *JAMA*. 2020;323:1612–4, <http://dx.doi.org/10.1001/jama.2020.4326>.
- Estella Á, Garcia Garmendia JL, de la Fuente C, Machado Casas JF, Yuste ME, Amaya Villar R, et al. A multicenter prospective study. *Med Intensiva*. 2021;8:S0210–5691, <http://dx.doi.org/10.1016/j.medin.2021.02.013>.
- Ruan Q, Yang K, Wang W, Jiang L, Song J. Clinical predictors of mortality due to 16 COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med*. 2020;46:846–8, <http://dx.doi.org/10.1007/s00134-020-05991-x>.
- Larcher R, Besnard N, Akouz A, Rabier E, Teule L, Vandercamere T, et al. Admission high-sensitive cardiac troponin T level increase is independently associated with higher mortality in critically ill patients with COVID-19: a multicenter study. *J Clin Med*. 2021;10:1656, <http://dx.doi.org/10.3390/jcm10081656>.
- Santoso A, Pranata R, Wibowo A, Al-Farabi MJ, Huang I, Antariksa B. Cardiac injury is associated with mortality and critically ill pneumonia in COVID-19: a meta-analysis. *Am J Emerg Med*. 2021;44:352–7, <http://dx.doi.org/10.1016/j.ajem.2020.04.052>.
- Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, et al., Executive Group on behalf of the Joint European Society of Cardiology (ESC)/American College of Cardiology (ACC)/American Heart Association (AHA)/World Heart Federation (WHF) Task Force for the Universal Definition of Myocardial Infarction. Fourth Universal Definition of Myocardial Infarction (2018). *J Am Coll Cardiol*. 2018;72:2231–64, <http://dx.doi.org/10.1016/j.jheart.2018.08.004>.
- Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. 2020;323:1061–9, <http://dx.doi.org/10.1001/jama.2020.1585>.
- Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395:1054–62, [http://dx.doi.org/10.1016/s0140-6736\(20\)30566-3](http://dx.doi.org/10.1016/s0140-6736(20)30566-3).
- Barman HA, Atici A, Sahin I, Alici G, Aktas Tekin E, Baycan ÖF, et al. Prognostic significance of cardiac injury in COVID-19 patients with and without coronary artery disease. *Coron Artery Dis*. 2021;32:359–66, <http://dx.doi.org/10.1097/mca.0000000000000914>.

C. Villavicencio\*, X. Daniel, C. Ferré, M. Cartanyá, Á. Pobo, I. Oliva, M. Roure, J. Leache, M. Bodí

*Critical Care Department, Joan XXIII – University Hospital, Tarragona, Spain*

\* Corresponding author.

E-mail address: [Christiananda@hotmail.com](mailto:Christiananda@hotmail.com)

(C. Villavicencio).

<https://doi.org/10.1016/j.medin.2022.05.003>  
0210-5691/ © 2022 Elsevier España, S.L.U. y SEMICYUC. All rights reserved.

## Assessment of plasma endocan for the prediction of mortality in COVID-19 patients undergoing veno-venous ECMO: A pilot study



## Evaluación del endocan para la predicción de la mortalidad en pacientes COVID-19 sometidos a ECMO veno-venosa: un estudio piloto

Dear Editor:

Patients with severe COVID-19 related acute respiratory distress syndrome (ARDS) may benefit from V-V ECMO support. However, this technique remains associated with frequent

complications and high mortality.<sup>1</sup> The early identification of patients with V-V ECMO who are likely to survive on ICU discharge therefore seems of major interest.

Endocan is a circulating proteoglycan secreted by the pulmonary vasculature under inflammatory conditions such as ARDS.<sup>2</sup> Recent data suggest that monitoring of blood endocan during 1st week following intensive care unit (ICU) admission may correlate with the severity of ARDS in Covid-19.<sup>3</sup> Thus, we aimed in this study to assess whether plasma endocan measurements performed on the day of V-V ECMO implantation (D0) and repeated seven days later (D7) may be effective in predicting mortality on ICU discharge.

This study was conducted in a 50-bed mixed ICU, from October 2020 to June 2021. We included all consecutive COVID-19 patients undergoing V-V ECMO implantation and with available results of plasma endocan measured on day of ECMO implantation.