



ORIGINAL ARTICLE

Can end-tidal CO₂ measurement replace arterial partial CO₂ in emergency department respiratory distress management?



Kudret Selki, Mehmet Cihat Demir, Erdinç Şengüldür, Emre Erdem, Hatice Güldal, Murat Taşdemir, Alp Kaan Furkan Kıcıroğlu, Mustafa Boğan*

Düzce University, School of Medicine, Department of Emergency Medicine, Düzce, Turkey

Received 6 November 2023; accepted 31 March 2024

KEYWORDS

End-tidal carbon dioxide;
Respiratory distress;
Arterial carbon dioxide;
Venous carbon dioxide;
Emergency department

Abstract

Objective: To assess the feasibility of using end-tidal carbon dioxide (EtCO₂) as a non-invasive substitute for partial pressure of arterial carbon dioxide (PaCO₂) in emergency department (ED) triage and follow-up, and to explore the potential of partial pressure of venous carbon dioxide (PvCO₂) as an alternative to PaCO₂.

Design: Prospective cross-sectional study.

Setting: Tertiary university hospital.

Patients or participants: 97 patients presenting with acute respiratory distress to the ED.

Interventions: EtCO₂, arterial blood gases, and venous blood gases measured at admission (0 min), 60 min, and 120 min.

Main variables of interest: CO₂ levels.

Results: Among 97 patients (mean age: 70.93 ± 9.6 years; 60.8% male), EtCO₂ > 45 mmHg at admission showed strong positive correlations with PaCO₂ and PvCO₂ ($r = 0.844$, $r = 0.803$; $p < 0.001$, respectively). Significant positive correlation was observed between 60-min EtCO₂ and PaCO₂ ($r = 0.729$; $p < 0.001$). Strong correlation between PaCO₂ and PvCO₂ at 120 min when EtCO₂ > 45 mmHg ($r = 0.870$; $p < 0.001$). EtCO₂ was higher in hospitalized patients compared to discharged ones.

Conclusions: EtCO₂ appears promising as a substitute for PaCO₂ in ED patients with acute respiratory distress within the initial two hours of treatment. Venous blood gas sampling offers a less invasive alternative to arterial sampling, facilitating simultaneous blood tests.

© 2024 Elsevier España, S.L.U. and SEMICYUC. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

* Corresponding author.

E-mail address: mustafabogan@duzce.edu.tr (M. Boğan).

PALABRAS CLAVE

Dióxido de carbono al final de la espiración;
 Insuficiencia respiratoria;
 Dióxido de carbono arterial;
 Dióxido de carbono venoso;
 Departamento de emergencias

¿Puede la medición del CO₂ al final de la espiración reemplazar el CO₂ parcial arterial en el tratamiento de la dificultad respiratoria en el departamento de urgencias?

Resumen

Objetivo: Evaluar la viabilidad de utilizar el dióxido de carbono al final de la espiración (EtCO₂) como un sustituto no invasivo de la presión parcial de dióxido de carbono arterial (PaCO₂) en el triaje y seguimiento en el departamento de emergencias (ED), y explorar el potencial de la presión parcial de dióxido de carbono venoso (PvCO₂) como alternativa a PaCO₂.

Diseño: Estudio prospectivo transversal.

Ámbito: Hospital universitario terciario.

Pacientes o participantes: 97 pacientes que se presentaron con dificultad respiratoria en el ED.

Intervenciones: Se midieron EtCO₂, gases en sangre arterial y gases en sangre venosa al ingreso (0 min), 60 min y 120 min.

Principales variables de interés: Niveles de CO₂.

Resultados: Entre 97 pacientes (edad media: 70,93 ± 9,6 años; 60,8% hombres), EtCO₂ > 45 al ingreso mostró correlaciones positivas fuertes con PaCO₂ y PvCO₂ (r = 0,844, r = 0,803; p < 0,001, respectivamente). Se observó una correlación positiva significativa entre EtCO₂ a los 60 min y PaCO₂ (r = 0,729; p < 0,001). Fuerte correlación entre PaCO₂ y PvCO₂ a los 120 min cuando EtCO₂ > 45 (r = 0,870; p < 0,001). EtCO₂ fue mayor en los pacientes hospitalizados en comparación con los dados de alta.

Conclusiones: EtCO₂ parece prometedor como sustituto de PaCO₂ en pacientes del ED con dificultad respiratoria dentro de las dos primeras horas de tratamiento. La toma de muestras de gases en sangre venosa ofrece una alternativa menos invasiva a la toma de muestras arterial, facilitando pruebas de sangre simultáneas.

© 2024 Elsevier España, S.L.U. y SEMICYUC. Se reservan todos los derechos, incluidos los de minería de texto y datos, entrenamiento de IA y tecnologías similares.

Introduction

The first point of admission for patients experiencing acute respiratory distress is usually the emergency department (ED). Establishing a safe and protected setting for individuals experiencing respiratory distress is paramount. Furthermore, it is imperative to efficiently detect and diagnose any underlying factors contributing to their shortness of breath. For this reason, measuring devices that evaluate respiratory status with noninvasive, easily applicable, and reliable methods in busy ED environments are of interest. End-tidal carbon dioxide (EtCO₂) measurement strengthens its place in clinical practice day by day as a handy and valuable method for evaluating patient ventilation in emergency settings.¹⁻⁶

The use of EtCO₂ monitoring is becoming increasingly common in a variety of clinical settings, both prehospital and hospital.⁷ In addition to being considered the most reliable confirmation method of the correct placement of the endotracheal tube, it also helps the physician in life support efforts, such as determining the quality of cardiopulmonary resuscitation and detecting the return of spontaneous circulation.^{8,9} Capnography is used in various situations, including procedural sedation and analgesia, patient transfer, pulmonary embolism, and acute follow-up of chronic obstructive pulmonary disease (COPD).¹⁰⁻¹⁴

EtCO₂ is the level of carbon dioxide released at the end of expiration. For noninvasive EtCO₂ measurement, capnometry, which provides only numerical values, or capnography, which provides both graphical and numerical results, is used.

Measurement is made as a mainstream or a sidestream. In sidestream, the CO₂ sensor is located inside the monitor, and a small portion of the exhaled air reaches the breathing circuit through a cable with a delay. In the mainstream type, the CO₂ sensor is directly connected to the breathing circuit and displays the result without delay. While sidestream can be used in intubated and non-intubated patients, mainstream is primarily used in intubated patients.^{2,6,12,15} While increased dead space and clogging with secretions are substantial disadvantages in the sidestream type, these are no longer a problem in the mainstream type, and the results are more reliable.^{2,6}

Partial pressure of arterial carbon dioxide (PaCO₂) is the gold standard for diagnosing and treating patients presenting to the ED with respiratory distress. However, in busy EDs, there is a need for fast, practical, cost-effective, and noninvasive methods that will help us comprehend the severity of patients while they are still in the triage phase. In many of the studies in the literature on non-intubated patients, correlation studies were conducted between EtCO₂ measurements obtained through sidestream detectors and PaCO₂.^{1,16-18}

The main aim of this study is to determine whether the EtCO₂ value obtained using the mainstream detector at the first presentation to the ED in non-intubated patients can be safely used instead of the patient's PaCO₂ value. The reason for using the mainstream detector, which is used in intubated patients, instead of the sidestream detector in non-intubated patients is that the previously described dis-

advantages that may alter the measurement results, such as dead space and obstruction by secretions, are less likely to be encountered. The secondary aim of the study was to determine whether partial pressure of venous carbon dioxide (PvCO₂) measurement would be an alternative to partial pressure of arterial carbon dioxide (PaCO₂) measurement, especially in patients who are not suitable for EtCO₂ (confused or intubated). The tertiary aim of the study was to determine whether EtCO₂ measurement was associated with hospitalizations.

Methods

Study design and setting

It was planned as a prospective cross-sectional study with patients who presented to the ED with respiratory distress between April and May 2023 (61 days). The study was conducted in the ED of a 317-bed tertiary university hospital with approximately 90,000 patient admissions per year. Local Ethics Committee approval was received for the study (Approval ID: 2023/42, dated March 20, 2023). Patients were included in the study by obtaining informed consent. Patient data were recorded on the study form simultaneously during the patients' application. Descriptive data were obtained from the hospital's electronic database and ED records.

Participants and measurements

Patients with a primary complaint of dyspnea regardless of etiology, older than 18 years of age, able to blow into a capnograph device and followed up in the emergency department for at least 2 h were included in the study. Of the 623 patients admitted to the emergency department with respiratory distress. We excluded 111 patients who refused to participate in the study, 162 patients with confused, uncooperative or ineffective blowing, 87 patients with inconclusive blood gas results due to coagulation, 25 patients undergoing tracheal intubation, 141 patients who have not been in the ED for at least 2 h (Fig. 1). Participation in the study was purely voluntary and not reliant on coercion. Demographic information of the patients, EtCO₂ level, PaCO₂ and PvCO₂ levels, pulse oximetry saturation level (SpO₂), and treatments given were recorded at the moment of first admission (0th minute), 60th minute, and 120th minute. The patients' hospitalization-discharge status, vital signs, and initial blood gas pH values were also noted. Arterial blood gas samples were obtained simultaneously from the radial artery, brachial artery, or femoral artery, and venous blood gas samples were obtained from the brachial vein simultaneously with heparinized syringes and quickly delivered to the laboratory.

Measuring EtCO₂

EtCO₂ measurements were performed by attaching a 3D-printed disposable apparatus (colorFabb 1.75 mm filament was used with a Creality Ender 3 Pro 3D printer) to the airway adapter of a mainstream EMMA® Capnograph device (PHASEIN AB Svärdvägen, Danderyd, Sweden). It is produced




Figure 1 Flowchart of patients excluded from the study.

for intubated patients. It was obtained with spontaneous single pulses from patients presenting with respiratory distress. Carbon dioxide level was classified as follows: <35 mmHg (hypocarbica), 35–45 mmHg (normocarbica), and >45 mmHg (hypercarbica).¹⁹ End tidal carbon dioxide measurement takes 3 s and arterial and venous carbon dioxide measurement takes approximately 10 min.

Statistical analysis

Statistical analyses were performed using SPSS software for Windows, version 23 (IBM, Chicago, IL, United States). Descriptive statistics are presented as numbers and percentages. Demographic data were presented as mean \pm standard deviation (SD), or median (interquartile range). Spearman correlation analysis was used for correlation analyses between EtCO₂, PaCO₂, PvCO₂, SpO₂, and cigarette pack/year. If the Spearman correlation coefficient (*r*) is below 0.20, it is considered an insignificant correlation. A weak correlation is marked between 0.20–0.50, a moderate correlation is observed between 0.50–0.70, and a strong correlation is observed when it is greater than 0.70.²⁰ Coefficients were given as significance values *p*. Correlation was considered significant at *p* < 0.05. The Pearson chi-square test and Fisher's exact test (when the expected number was less than five) were used for independent categorical variables. Bonferroni correction was made for subgroup analysis, and *p* < 0.016 was considered significant. The McNemar test was used for dependent categorical variables. For the relationship between the treatment given with EtCO₂ and hospitalization discharge, the Mann-Whitney *U* test was used for independent two-group comparisons that did not show normal distribution, and the Kruskal Wallis test was used for multi-group comparisons. Student-t test was used for numerical two-group variables with normal distribution. *p* < 0.05 was considered significant.

Results

Of the 623 patients admitted to the ED with respiratory distress, 512 agreed to participate in the study (82%). Of these, 415 were excluded from the study for reasons explained in the method. The study was planned with 97 patients (Fig. 1). The average age of the patients was 70.9 \pm 9.6 years, and 60.8% (*n* = 59) were male. 43.3% (*n* = 42) of the patients who admitted were ex-smokers. The most common comorbidities were hypertension (HT) with 76% (*n* = 74) and COPD with 58% (*n* = 57). 34% of the patients were hypercarbic at first presentation (0th minute, PaCO₂ at admission >45 mmHg). Descriptive data of the patients are presented in Table 1

The results of EtCO₂ values between hospitalized and discharged patients, including measurements at admission, 60th and 120th minutes, are shown in Table 2. EtCO₂ value was statistically significantly higher in all measurements in hospitalized patients than in discharged patients.

Table 3 shows the results of the correlations between EtCO₂ and PaCO₂ and PvCO₂ measurements at the time of admission, the sixtieth and the 120th minute, in detail. When the EtCO₂ value was above 45 mmHg, a strong positive (*r* = 0.844 and *r* = 0.803) and significant relationship was

Table 1 Descriptive data of patients with respiratory distress (*n* = 97).

Features	<i>n</i> (%)
Gender, Male	59 (60.8%)
Age, years (mean + SD)	70.9 \pm 9.6
Initial arterial blood gas pH value [mean + SD (min–max)]	7.37 \pm 0.08 (7.081–7.57)
Treatment at admission	
Inhaler	45 (46.4%)
bronchodilator	12 (12.4%)
Diuretic	12 (12.4%)
Noninvasive mechanical ventilation (NIMV)	11 (11.3%)
Oxygen support only	12 (12.4%)
Inhaler + diuretic	5 (5.2%)
Diuretic + NIMV	0 (0.0%)
Invasive mechanic ventilation (IMV)	0 (0.0%)
Treatment at the first hour	
Inhaler	40 (41.2%)
bronchodilator	15 (15.5%)
Diuretic	15 (15.5%)
NIMV	10 (10.3%)
Oxygen support only	12 (12.4%)
Inhaler + diuretic	5 (5.2%)
Diuretic + NIMV	0 (0.0%)
IMV	0 (0.0%)
Treatment at the second hour	
Inhaler	35 (36.1%)
bronchodilator	17 (17.5%)
Diuretic	18 (18.5%)
NIMV	12 (12.4%)
Oxygen support only	11 (11.3%)
Inhaler + diuretic	4 (4.1%)
Diuretic + NIMV	0 (0.0%)
IMV	0 (0.0%)
Comorbidities^a	

Table 1 (Continued.)

Heart failure	37 (38%)
Hypertension	74 (76%)
Chronic obstructive pulmonary disease	57 (58%) 36 (37%) 13 (13%) 11 (11%)
Diabetes	7 (7%)
Coronary artery disease	9 (9%) 4 (4%)
Chronic renal failure	1 (1%)
Lung malignancy	
Tuberculosis	
Other malignancies	
Interstitial lung disease	
Smoking	
Active drinker	16 (16.5%)
Ex-smoker	42 (43.3%)
Never drank	39 (40.2%)
Diagnosis of chronic hypercarbia (PaCO₂ at admission > 45 mmHg)	
COPD exacerbation	33 (34.0%) 18 (18.6%)
Heart failure	11 (11.3%)
Pneumonia	2 (2.1%)
Acute renal failure	2 (2.1%)
Outcome	
Discharged	62 (63.9%)
Chest disease ward admission	23 (23.7%) 5 (5.2%)
Cardiology ward admission	7 (7.2%)
Intensive care unit	

^a Patients may have more than one comorbid condition.

Table 3 Correlation analyses of EtCO₂ and PaCO₂ and PvCO₂ measurements.

Zeroth minute		EtCO ₂	PaCO ₂	PvCO ₂
EtCO ₂ (mmHg) <35	<i>r</i>		0.820	0.772
	<i>p</i>		<0.001	<0.001
	<i>r</i>		–	–
35–45	<i>p</i>		0.064	0.056
	<i>r</i>		0.621	0.657
	<i>p</i>		<0.001	<0.001
>45	<i>r</i>		0.844	0.803
	<i>p</i>		<0.001	<0.001
	<i>r</i>		0.820	0.891
PaCO ₂ (mmHg)	<i>p</i>	<0.001		<0.001
	<i>r</i>	0.772	0.891	
	<i>p</i>	<0.001	<0.001	
60th minute EtCO ₂	<i>r</i>		0.729	0.653
	<i>p</i>		<0.001	<0.001
	<i>r</i>		–	–
<35	<i>p</i>		0.341	0.361
	<i>r</i>		0.635	0.551
	<i>p</i>		<0.001	<0.001
35–45	<i>r</i>		0.730	0.702
	<i>p</i>		<0.001	<0.001
	<i>r</i>		0.729	0.937
>45	<i>p</i>	<0.001		<0.001
	<i>r</i>	0.653	0.937	
	<i>p</i>	<0.001	<0.001	
120th minute EtCO ₂	<i>r</i>		0.677	0.609
	<i>p</i>		<0.001	<0.001
	<i>r</i>		–	–
<35	<i>p</i>		0.323	0.336
	<i>r</i>		0.480	0.415
	<i>p</i>		<0.001	<0.001
35–45	<i>r</i>		0.667	0.563
	<i>p</i>		<0.001	<0.001
	<i>r</i>		0.677	0.876
>45	<i>p</i>	<0.001		<0.001
	<i>r</i>	0.609	0.876	
	<i>p</i>	<0.001	<0.001	

EtCO₂: End-tidal carbon dioxide; PaCO₂: Partial pressure of arterial carbon dioxide; PvCO₂: Partial pressure of venous carbon dioxide. The correlation is significant at the $p < 0.05$ level; *r*: Spearman correlation coefficient.

found between EtCO₂ and PaCO₂ and PvCO₂ measurements ($p < 0.001$ for both).

When the EtCO₂ value was above 45 mmHg, a strong positive and significant relationship was found between EtCO₂

Table 2 Comparison of end-tidal carbon dioxide (EtCO₂) values of patients with respiratory distress according to hospitalization status.

Parameters	All patients	Inpatients	Discharged	p value
EtCO ₂ at admission	35 (30–45)	46.17 ± 15.235	33.95 ± 9.31	<0.001
EtCO ₂ at the first hour	33 (28–41)	42.69 ± 14.784	32.90 ± 7.923	<0.001
EtCO ₂ at the second hour	34 (29–41.25)	42.37 ± 15.923	34.06 ± 8.400	0.006

Values were shown as mean ± SD, or median (interquartile range). $p < 0.05$ was considered statistically significant.




Figure 2 Linear correlation between end-tidal carbon dioxide (EtCO₂) and PaCO₂ levels at the admission.




Figure 3 Linear correlation between end-tidal carbon dioxide (EtCO₂) and PaCO₂ levels at the time of 60th min.

and PaCO₂ and PvCO₂ measurements ($r = 0.730$ and $r = 0.702$; $p < 0.001$ for both).

A strong positive and significant correlation was found between PaCO₂ and PvCO₂ measurement when the EtCO₂ value was below 35 mmHg ($r = 0.858$; $p < 0.001$). A strong positive and significant relationship was seen when PaCO₂ and PvCO₂ EtCO₂ values were between 35–45 mmHg ($r = 0.883$; $p < 0.001$). There was a strong positive and significant relationship between PaCO₂ and PvCO₂ measurement when the EtCO₂ value was above 45 ($r = 0.891$; $p < 0.001$).

In the last measurements at the 120th minute, a moderately positive and significant relationship was found between EtCO₂, PaCO₂ and PvCO₂ values ($r = 0.677$ and $r = 0.609$; $p < 0.001$ for both). When the EtCO₂ value was above 45 mmHg, a moderate positive and significant relationship was observed between EtCO₂ and PaCO₂ and PvCO₂ measurements ($r = 0.667$ and $r = 0.563$; $p < 0.001$ for both). When the EtCO₂ value was below 35 mmHg, a moderately positive and significant correlation was observed between PaCO₂ and PvCO₂ measurement ($r = 0.697$; $p < 0.001$). A strong positive and significant ($r = 0.791$; $p < 0.001$) relation-

ship was detected between PaCO₂ and PvCO₂ measurements when the EtCO₂ value was between 35–45 mmHg. There was a strong positive and significant relationship between PaCO₂ and PvCO₂ measurement when the EtCO₂ value was above 45 mmHg ($r = 0.870$; $p < 0.001$).

We found a strong positive and significant relationship between EtCO₂ and PaCO₂ measurements at admission ($r = 0.820$; $p < 0.001$). The variance explained by the variables on each other is 67% (Fig. 2). We found a strong positive and significant relationship between 60th-minute EtCO₂ and PaCO₂ measurement ($r = 0.729$; $p < 0.001$). The variance explained by the variables on each other is 53% (Fig. 3). A strong positive and significant relationship existed between PaCO₂ and PvCO₂ measurement at the 60th minute ($r = 0.937$; $p < 0.001$). The variance explained by the variables on each other is 87% (Fig. 4). A moderately negative and significant relationship existed between EtCO₂ and SpO₂ measurement at admission ($r = -0.516$; $p < 0.001$). There was no significant relationship between EtCO₂ and SpO₂ measurement at the 60th minute and 120th minute ($p = 0.402$ and $p = 0.771$).




Figure 4 Linear correlation between PaCO₂ and PvCO₂ levels at the time of 60th min.

Discussion

Our study found a high correlation between EtCO₂ at admission (zero minutes) and PaCO₂ and PvCO₂ measurements ($r = 0.820$ and $r = 0.772$, respectively). We also found a high correlation between PaCO₂ and PvCO₂ ($r = 0.891$; $p < 0.001$). We found a moderately positive and significant relationship between EtCO₂ and PaCO₂ and PvCO₂ measurements when the EtCO₂ value was between 35–45 mmHg ($r = 0.621$ and $r = 0.657$, respectively; $p < 0.001$ for both). When the EtCO₂ value was above 45 mmHg, a strong positive and significant relationship was found between EtCO₂ and PaCO₂ and PvCO₂ measurements ($r = 0.844$ and $r = 0.803$, respectively; $p < 0.001$ for both). Apart from this, a strong correlation was observed between PaCO₂ and PvCO₂ for all measurements. These results showed that in triage patients presenting to the emergency department with respiratory distress, if the EtCO₂ value measured with a single blow is above 45 mmHg, the patient is considered hypercapnic, and treatment can be started early. Another important conclusion is that venous blood gas measurement is a strong alternative to arterial blood gas measurement, which is an excruciating procedure.

Arterial blood gas sampling in a patient presenting with respiratory distress is often challenging for both the patient and the physician. It may need to be repeated from time to time. Many clinical studies have been designed to increase patient comfort and identify new reliable methods. One of the alternative methods is EtCO₂ measurement. Healey et al. demonstrated a high correlation between EtCO₂ and PaCO₂ measured before and after withdrawal of assisted-controlled mechanical ventilation.²¹ Plewa et al. found that the patients' EtCO₂ value obtained by the forced expiration model showed a high correlation with the PaCO₂ value. They also reported the negative predictive value of capnographic hypercapnia (EtCO₂ > 45 mmHg) in detecting arterial hypercapnia (PaCO₂ > 45 mmHg) as 95%.²² Other studies conducted with non-intubated patients have also reported a high correlation.^{1,18}

Measuring EtCO₂ in non-intubated patients has difficulties. Patient cooperation comes first among these difficulties. Although the initial panic of outpatients with respiratory distress is another difficulty for measuring EtCO₂

in the ED, it is easier than arterial blood gas sampling. Suzuki et al. their study of non-intubated patients followed up due to respiratory diseases found a high positive correlation between EtCO₂ and PaCO₂ ($r = 0.88$; $p < 0.0001$).²³ Joe et al. in their study, it was stated that there was a significant correlation between EtCO₂ measurement and PvCO₂ in patients with chronic pulmonary disease and using oxygen ($r = 0.63$) and that the PvCO₂ level of the patients could be predicted with capnography, which is a noninvasive method.²⁴ These studies support our results. EtCO₂ is a more comfortable, faster, and easily reproducible alternative to arterial blood gas sampling when starting the initial treatment of patients presenting to the ED with respiratory distress.

Venous blood gas sampling is easy. Although it is an invasive procedure, it is not as difficult for the patient and physician to repeat as arterial blood gas sampling. Therefore, understanding the relationship between PaCO₂, EtCO₂ and PvCO₂ was investigated.^{25,26} However, studies in the literature contain a partial consensus. In a systematic review, only 22.5% of included studies identified a strong correlation between arterial and venous parameters.²⁵ Predictably, different results may occur depending on patient characteristics. In their methodological study with 20 patients with various diagnoses and complaints, Lumholdt et al. investigated the mathematical adaptability of peripheral venous blood gas values to arterial blood gas values. They stated that there was a mathematically predictable relationship between PaCO₂ and PvCO₂.²⁷ A study in non-intubated patients followed up for respiratory diseases indicated a positive correlation between PaCO₂ and PvCO₂ ($r = 0.81$; $p < 0.001$).²³ Our study showed a high correlation between venous and arterial blood gas in terms of partial carbon dioxide pressure (pCO₂) in patients with respiratory distress from admission until the second hour of treatment. In light of all these findings, end-tidal carbon dioxide measurement seems to be an excellent alternative to arterial carbon dioxide measurement. In addition, venous carbon dioxide measurement is a good alternative in non-cooperative patients unsuitable for non-invasive end-tidal carbon dioxide measurement. The fact that it is less painful and can be taken from the vein simultaneously with many other blood tests makes venous carbon dioxide measurement more important.

Conflicting results have been revealed in the literature regarding whether capnography can replace PaCO₂ as a reliable tool in patients with acute respiratory distress.^{2,3,28,29} In our study, EtCO₂ values of patients admitted with acute respiratory distress at admission and in the 1st and 2nd hours after treatment were associated with hospitalization. EtCO₂ measurements were higher in hospitalized patients compared to discharged patients.

The main limitation of this study is that it was conducted in a single center with a study group consisting of patients who presented to the ED of a tertiary university hospital. The low number of cases in the city where the study was conducted caused the number of patients included in this study to be limited. Second, we did not correct pCO₂ based on body temperature in any patient. Phan et al. noted that EtCO₂ correlated better with pCO₂ after temperature correction.¹⁷ Third, the patient's respiratory rate was not recorded. When the respiratory rate increases, the dead space ratio increases, and the EtCO₂ level may decrease. Additionally, the capnometer may not detect EtCO₂ changes in case of a high respiratory rate.²⁴ The fourth limitation is that end tidal carbon dioxide measurement can only be performed in cooperative patients. Finally, conditions such as age, existing lung disease, and smoking may cause EtCO₂ values to change. Prediction ability decreases in the elderly and those with lung disease.²² We did not perform subgroup analysis on the patients. Large-scale and multi-center studies in which subgroups are studied separately and necessary corrections are made according to the patient's vital values, such as respiratory rate and temperature, are needed to confirm this alternative diagnostic method.

Conclusion

The EtCO₂ value obtained from adding an insufflation device we produced to the capnograph and spontaneously inhaling patients with respiratory distress is a powerful alternative to arterial blood gas in the ED triage of patients and the first two hours of acute treatment. It has also been shown that venous blood gas may be a good alternative. In patients with acute respiratory distress (patients who can cooperate and whose hemodynamics have not yet deteriorated), this simple, noninvasive EtCO₂ measurement is also helpful in the early prediction of hospitalization decisions.

Author's contribution

KS, MCD, MB contributed to conception; KS, MCD, MB contributed to design; MCD, EŞ, MB contributed to supervision; KS, HG, AKFK, MT, EE contributed to data collection and processing; KS, MCD contributed to analysis and interpretation; KS, MCD, EŞ, MB contributed to literature review; KS, MCD, EŞ, MB contributed to writing; MCD, MB contributed to critical review.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Declaration of interest

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Availability of data and materials

Submitted work is original and has not been published elsewhere in any language. Raw data are available for the editor on request.

Ethical statement

Ethics Committee approval was obtained from the local ethics committee (Date: March 20, 2023, Decision No: 2023/42).

Informed consent

Written consent was obtained from all patients.

Human rights

Authors declare that human rights were respected according to the Declaration of Helsinki.

The address of the institution where the study was performed

Düzce University, School of Medicine, Düzce, Türkiye.

Acknowledgements

None.

References

1. Barton CW, Wang ES. Correlation of end-tidal CO₂ measurements to arterial PaCO₂ in nonintubated patients. *Ann Emerg Med.* 1994;23(3):560-3.
2. Cinar O, Acar YA, Arziman I, Kilic E, Eyi YE, Ocal R. Can mainstream end-tidal carbon dioxide measurement accurately predict the arterial carbon dioxide level of patients with acute dyspnea in ED. *Am J Emerg Med.* 2012;30(2):358-61.
3. Doğan NÖ, Şener A, Günaydın GP, İçme F, Çelik GK, Kavaklı HŞ, et al. The accuracy of mainstream end-tidal carbon dioxide levels to predict the severity of chronic obstructive pulmonary disease exacerbations presented to the ED. *Am J Emerg Med.* 2014;32(5):408-11.
4. Manifold CA, Davids N, Villers LC, Wampler DA. Capnography for the nonintubated patient in the emergency setting. *J Emerg Med.* 2013;45(4):626-32.
5. Nagler J, Krauss B. Capnography: a valuable tool for airway management. *Emerg Med Clin North Am.* 2008;26(4):881-97, vii.
6. Yosefy C, Hay E, Nasri Y, Magen E, Reisin L. End tidal carbon dioxide as a predictor of the arterial PCO₂ in the emergency department setting. *Emerg Med J.* 2004;21(5):557-9.
7. Langan M. Availability and clinical utilization of capnography in the prehospital setting. *Conn Med.* 2011;75(4):197-201.

8. Sandroni C, De Santis P, D'Arrigo S. Capnography during cardiac arrest. *Resuscitation*. 2018;132:73–7.
9. Silvestri S, Ralls GA, Krauss B, Thundiyil J, Rothrock SG, Senn A, et al. The effectiveness of out-of-hospital use of continuous end-tidal carbon dioxide monitoring on the rate of unrecognized misplaced intubation within a regional emergency medical services system. *Ann Emerg Med*. 2005;45(5):497–503.
10. Day D. Keeping patients safe during intrahospital transport. *Crit Care Nurse*. 2010;30(4):18–32, quiz 33.
11. Hemnes AR, Newman AL, Rosenbaum B, Barrett TW, Zhou C, Rice TW, et al. Bedside end-tidal CO₂ tension as a screening tool to exclude pulmonary embolism. *Eur Respir J*. 2010;35(4):735–41.
12. Krauss B, Hess DR. Capnography for procedural sedation and analgesia in the emergency department. *Ann Emerg Med*. 2007;50(2):172–81.
13. Long B, Koyfman A, Vivirito MA. Capnography in the emergency department: a review of uses, waveforms, and limitations. *J Emerg Med*. 2017;53(6):829–42.
14. Waugh JB, Epps CA, Khodneva YA. Capnography enhances surveillance of respiratory events during procedural sedation: a meta-analysis. *J Clin Anesth*. 2011;23(3):189–96.
15. Pekdemir M, Cinar O, Yilmaz S, Yaka E, Yuksel M. Disparity between mainstream and sidestream end-tidal carbon dioxide values and arterial carbon dioxide levels. *Respir Care*. 2013;58(7):1152–6.
16. Wang W, Zhao Z, Tian X, Ma X, Xu L, Shang G. Noninvasive carbon dioxide monitoring in pediatric patients undergoing laparoscopic surgery: transcutaneous vs. end-tidal techniques. *BMC Pediatr*. 2023;23(1):20.
17. Phan CQ, Tremper KK, Lee SE, Barker SJ. Noninvasive monitoring of carbon dioxide: a comparison of the partial pressure of transcutaneous and end-tidal carbon dioxide with the partial pressure of arterial carbon dioxide. *J Clin Monit*. 1987;3(3):149–54.
18. Lenz G, Heipertz W, Epple E. Capnometry for continuous post-operative monitoring of nonintubated, spontaneously breathing patients. *J Clin Monit*. 1991;7(3):245–8.
19. Schneider AG, Eastwood GM, Bellomo R, Bailey M, Lipcsey M, Pilcher D, et al. Arterial carbon dioxide tension and outcome in patients admitted to the intensive care unit after cardiac arrest. *Resuscitation*. 2013;84(7):927–34.
20. Kozak M. What is Strong Correlation? *Teaching Statistics*. 2009;31(3):85–6.
21. Healey CJ, Fedullo AJ, Swinburne AJ, Wahl GW. Comparison of noninvasive measurements of carbon dioxide tension during withdrawal from mechanical ventilation. *Crit Care Med*. 1987;15(8):764–8.
22. Plewa MC, Sikora S, Engoren M, Tome D, Thomas J, Deuster A. Evaluation of capnography in nonintubated emergency department patients with respiratory distress. *Acad Emerg Med*. 1995;2(10):901–8.
23. Suzuki M, Fujimoto S, Sakamoto K, Tamura K, Ishii S, Iikura M, et al. Clinical usefulness of end-tidal CO₂ measured using a portable capnometer in patients with respiratory disease. *Clin Respir J*. 2023;17(2):96–104.
24. Jo T, Inomata M, Takada K, Yoshimura H, Tone M, Awano N, et al. Usefulness of measurement of end-tidal CO₂ using a portable capnometer in patients with chronic respiratory failure receiving long-term oxygen therapy. *Intern Med*. 2020;59(14):1711–20.
25. Saberian L, Sharif M, Aarabi M, Broumand B, Shafiee MA. Arterial versus venous blood gas analysis comparisons, appropriateness, and alternatives in different acid/base clinical settings: a systematic review. *Cureus*. 2023;15(7):e41707.
26. Toftegaard M, Rees SE, Andreassen S. Correlation between acid-base parameters measured in arterial blood and venous blood sampled peripherally, from vena cavae superior, and from the pulmonary artery. *Eur J Emerg Med*. 2008;15(2):86–91.
27. Lumholdt M, Damgaard KA, Christensen EF, Leutscher PDC. Mathematical arterialisation of peripheral venous blood gas for obtainment of arterial blood gas values: a methodological validation study in the clinical setting. *J Clin Monit Comput*. 2019;33(4):733–40.
28. Kartal M, Goksu E, Eray O, Isik S, Sayrac AV, Yigit OE, et al. The value of ETCO₂ measurement for COPD patients in the emergency department. *Eur J Emerg Med*. 2011;18(1):9–12.
29. Delorme S, Freund Y, Renault R, Devilliers C, Castro S, Chopin S, et al. Concordance between capnography and capnia in adults admitted for acute dyspnea in an ED. *Am J Emerg Med*. 2010;28(6):711–4.