



SCIENTIFIC LETTER

High flow Tracheal oxygen: assessment of diaphragmatic functionality by ultrasonography in adults during weaning from mechanical ventilation



Alto flujo de oxígeno traqueal: evaluación de la funcionalidad diafragmática mediante ultrasonografía en adultos durante el proceso de destete de la ventilación mecánica

High-flow oxygen therapy via nasal cannula (HFNC) has demonstrated physiological benefits for patients with acute hypoxemic respiratory failure, including improvements in oxygenation, increased End Expiratory Lung Volume (EELV), and reduced work of breathing.¹ Recent investigations have explored the potential of HFNC to facilitate weaning from invasive mechanical ventilation (iMV) in tracheostomized patients. However, these studies primarily focused on evaluating diaphragm functionality solely at the end of spontaneous breathing, without observing dynamic changes in Work of Breathing (WOB) throughout the entire Spontaneous Breathing (SB) period.^{2–4} Therefore, we conducted a case series randomized crossover study to assess dynamic changes in inspiratory effort, measured using diaphragm ultrasound, from the initiation of SB to one hour thereafter. This study aims to evaluate the effects of High-Flow Tracheal Oxygen (HFT) versus Standard Oxygen Therapy (SOT) on inspiratory effort in tracheostomized patients during the weaning phase from iMV.

We included tracheostomized adult patients who tolerated pressure support ventilation for at least two hours, in two tertiary university hospitals in Argentina. Exclu-

sion criteria were neuromuscular diseases, pregnancy, and individuals unable to undergo ultrasound diaphragm measurements. Ethical approval for the study was granted by the Institutional Review Board (#5093). Following the acquisition of informed consent, enrolled patients were randomly allocated to receive either HFT or SOT first, with a respiratory muscles resting period for 2 h in pressure support (Fig. 1).

The HFT strategy was administered with a flow of 60 L/min and a set temperature between 34 and 37 °C. The fraction of inspired oxygen was adjusted to achieve a SpO₂ of 88% or higher. In the SOT strategy, a passive humidifier device was connected to the tracheostomy cannula. Oxygen was delivered at flow rates ranging from 2 to 15 l/min to maintain SpO₂ at 88% or above.

The inspiratory effort was estimated through ultrasound assessments conducted at 5 and 60 min after iMV disconnection. Each evaluation included ultrasound measurements of diaphragmatic thickening fraction (Tfdi), diaphragmatic excursion, and Lung Ultrasound Score (LUS) to estimate lung aeration^{5,6}.

The primary outcome was to compare the change in Tfdi between 5 and 60 min after SB with HFT versus SOT. As a secondary outcome, we compared diaphragmatic excursion and pulmonary aeration between 5 and 60 min in both groups.

Thirteen patients met the inclusion criteria, but three were excluded due to difficulty in obtaining ultrasound measurements. Ten patients completed the study, with a median age of 70 (IQR 64–75). Seventy percent were male (n=7), with a median APACHE II score of 13 (IQR 10–17) and a median duration of iMV of 17 days (IQR 12–25) before inclusion. Baseline characteristics are provided in the Supplementary Material.

For the SOT strategy, the median Tfdi was 25% (IQR 13%–36%) at 5 min and 26% (IQR 21%–36%) at 60 min. During the HFT strategy, the median Tfdi was 26% (IQR 20%–36%)




Figure 1 Flow chart of the study design.

Table 1 Ultrasound measured.

	Initial measurement (5 min)	Final measurement (60 min)	Temporary changes (60–5 min)
HFT strategy			
Diaphragm excursion mm (SD)	16.6 (7.02)	15.3 (7.44)	–1.38 (8.32)
Tfdi % (SD)	0.33 (0.25)	0.29 (0.12)	–0.04 (0.31)
LUS score (point)	10.8 (7.19)	12.6 (9.18)	–1.80 (2.97)
SOT strategy			
Diaphragm excursion mm (SD)	14.6 (7.75)	16.5 (7.43)	1.89 (5.04)
Tfdi % (SD)	0.26 (0.14)	0.32 (0.18)	0.06 (0.12)
LUS score (point)	12.0 (8.08)	13.1 (9.64)	–0.44 (4.16)

HFT: High-Flow Tracheal Oxygen; SOT: Standard Oxygen Therapy; Tfdi: Diaphragmatic Thickening Fraction; LUS: Lung Ultrasound Score.

%) at 5 min and 27% (IQR 20 %–33 %) at 60 min. Dynamic changes in Tfdi between 5 and 60 min in both strategies showed nonsignificant differences ($p=0.4$).

Diaphragmatic excursion in the SOT strategy at 5 min exhibited a median of 13 mm (IQR 11–15), increasing to 15 mm (IQR 11–22) at 60 min. In the HFT strategy, the median diaphragmatic excursion was 15 mm (IQR 13–22) at the beginning, decreasing to 14 mm (IQR 10–16) after 60 min. Dynamic changes in diaphragmatic excursion between 5 and 60 min in both strategies showed nonsignificant differences ($p=0.3$).

Regarding lung aeration, the LUS score for the SOT strategy was 10 (IQR 6–21) at 5 min and 10 (IQR 6–20) at 60 min. During HFT, the median LUS score was 8 (IQR 6–17) at 5 min and 9 (IQR 7–20) at 60 min. Dynamic changes in LUS score between 5 and 60 min in both strategies showed nonsignificant differences ($p=0.4$). Table 1 describes ultrasound measures.

The application of HFT during the weaning phase from iMV did not significantly alter inspiratory effort, diaphragmatic excursion, or lung aeration compared to SOT. Previous studies assessing the impact of HFT on inspiratory effort similarly failed to demonstrate an improvement in diaphragm functionality.^{7,8} Improvements were limited to the SpO₂/FIO₂ ratio, a mild increase in positive end-expiratory pressure (PEEP), and a reduction in respiratory rate. These benefits appear to be flow-dependent, manifesting when applying flow rates exceeding 40 L/min.⁹ The absence of substantial benefits with HFT may be attributed to several factors. Firstly, the direct entry of inspiratory flow into the trachea bypasses the oropharyngeal dead space, diminishing potential effects. Secondly, the absence of glottic closure with the tracheal tube eliminates a crucial factor in maintaining PEEP.¹⁰ Lastly, the direction of inspiratory flow from the HFT tracheal device may not sufficiently increase expiratory resistances, resulting in a loss of the potential PEEP effect.

Several limitations of this study include the small sample size, which reduces statistical power. Secondly, we did not evaluate blood gas analyses, as existing literature indicates that HFT could improve arterial partial pressure of oxygen without affecting pCO₂(3) Finally, the lack of blinding among investigators could potentially introduce bias in ultrasound measurements in both modalities.

The strength of this study lies in describing the temporal changes in diaphragmatic functionality over 60 min, contrasting with other studies that evaluated shorter periods.⁸

This approach offers dynamic insights into both diaphragm functionality over time and its muscular tolerance during the weaning process from iMV, providing a comprehensive understanding of the diaphragmatic response throughout the entire duration, rather than at a specific moment.

In our population, the application of HFT during the weaning phase of iMV did not generate significant changes in inspiratory effort, in comparison to SOT. However, this preliminary finding could inform future investigations evaluating the impact of HFT in tracheostomized patients with prolonged mechanical ventilation.

Author contribution

Last name: Fernández Ceballos

First name: Ignacio

Author contribution: Literature search, data collection, study design, manuscript preparation.

First name: Iván

Last name: Huespe

Author contribution: data analysis, manuscript preparation.

Last name: Venuti

First name: María

Author contribution: Study design, data analysis.

Last name: Ferreyro

First name: Bruno

Author contribution: Study design, manuscript preparation.

Last name: Dianti

First name: Jose

Author contribution: Data analysis, writing.

Last name: Famiglietti

First name: Romina

Author contribution: Literature search, data collection.

Last name: Rivera

First name: Ana

Author contribution: data collection.

First name: Indalecio

Last name: Carboni Bisso

Author contribution: data collection.

First name: Marcos

Last name: Las Heras

Author contribution: Study design, review manuscript.

All authors have reviewed and approved the final version of the article.

Acknowledgments

The research team would like to thank María de los Angeles Magaz for translating the following manuscript into English, and Jorge Menedez for providing training and guidance in ultrasound measurements.

Declaration of generative AI and AI-assisted technologies in the writing process:

Throughout the preparation of this work, the authors utilised ChatGPT 3.5 to enhance writing quality and conciseness. Subsequently, the authors thoroughly reviewed and edited the content as necessary, assuming full responsibility for the publication's content.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.medin.2024.05.017>.

References

- Mauri T, Turrini C, Eronia N, Grasselli G, Volta CA, Bellani G, et al. Physiologic effects of high-flow nasal cannula in acute hypoxemic respiratory failure. *Am J Respir Crit Care Med*. 2017;195(9):1207–15, <http://dx.doi.org/10.1164/rccm.201605-0916OC>.
- Egbers PH, Sutt AL, Petersson JE, Bergström L, Sundman E. High-flow via a tracheostomy tube and speaking valve during weaning from mechanical ventilation and tracheostomy. *Acta Anaesthesiol Scand*. 2023;67(10):1403–13, <http://dx.doi.org/10.1111/aas.14305>.
- Corley A, Edwards M, Spooner AJ, Dunster KR, Anstey C, Fraser JF. High-flow oxygen via tracheostomy improves oxygenation in patients weaning from mechanical ventilation: a randomised crossover study. *Intensive Care Med*. 2017;43(3):465–7, <http://dx.doi.org/10.1007/s00134-016-4634-7>.
- Vadi S, Phadtare S, Shetty K. High-flow oxygen therapy via tracheostomy to liberate COVID-19-induced ARDS from invasive ventilation: a case series. *Indian J Crit Care Med*. 2021;25(6):724–8, <http://dx.doi.org/10.5005/jp-journals-10071-23858>.
- Serrano DAR, Terán PP, Blancas R, Arroyo M. Pleuropulmonary and diaphragmatic ultrasound in intensive care medicine. *Med Intensiva*. 2023;47(10):594–602, <http://dx.doi.org/10.1016/j.medine.2023.06.009>.
- Zapata L, Blancas R, Conejo-Márquez I, García-de-Acila M. Role of ultrasound in acute respiratory failure and in the weaning of mechanical ventilation. *Med Intensiva*. 2023;47(9):529–42, <http://dx.doi.org/10.1016/j.medine.2023.03.018>.
- Lersritwimanmaen P, Rittayamai N, Tscheikuna J, Brochard L. High-flow oxygen therapy in tracheostomized subjects with prolonged mechanical ventilation: a randomized crossover physiologic study. *Respir Care*. 2021;66(5):806–13, <http://dx.doi.org/10.4187/respcare.08585>.
- Stripoli T, Spadaro S, Di Mussi R, Volta CA, Terrotoli P, De Carlo F, et al. High-flow oxygen therapy in tracheostomized patients at high risk of weaning failure. *Ann Intensive Care*. 2019;9(1):4, <http://dx.doi.org/10.1186/s13613-019-0482-2>.
- Natalini D, Grieco DL, Santantonio MT, Mincione L, Toni F, Anzellotti GM, et al. Physiological effects of high-flow oxygen in tracheostomized patients. *Ann Intensive Care*. 2019;9(1):114, <http://dx.doi.org/10.1186/s13613-019-0591-y>.
- Thomas M, Joshi R, Cave G. How much PEEP does high flow deliver via tracheostomy? A literature review and bench-top experiment. *Crit Care Res Pract*. 2021;2021:6036891, <http://dx.doi.org/10.1155/2021/6036891>.

Ignacio Fernández Ceballos^{a,b,*}, Ivan Alfredo Huespe^a, María Sofía Venuti^a, Bruno Leonel Ferreyro^c, José María Dianti^{d,e}, Romina Famiglietti^a, Ana Montserrat Rivera^a, Indalecio Carboni Bisso^a, Marcos Jose Las Heras^a

^a *Critical Care Department, Hospital Italiano de Buenos Aires, Buenos Aires, Argentina*

^b *Critical Care Department, Hospital Italiano de San Justo Agustín Roca, Buenos Aires, Argentina*

^c *Department of Medicine, Division of Respiriology, Sinai Health System and University Health Network, Toronto, Canada*

^d *Interdepartmental Division of Critical Care Medicine, University of Toronto, Toronto, Ontario, Canada*

^e *Adult Intensive Care Unit, CEMIC, Buenos Aires, Argentina*

* Corresponding author.

E-mail address: ignacio.fernandez@hospitalitaliano.org.ar (I. Fernández Ceballos).

<https://doi.org/10.1016/j.medin.2024.05.017>

0210-5691/ © 2024 Elsevier España, S.L.U. and SEMICYUC. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

Motivos de rechazo de ingreso en UCI en población oncológica y su asociación con la mortalidad a seis meses



Reasons for refusal of admission to the ICU in oncological population and their association with 6-month mortality

Sr. Editor,

La incorporación de la innovación, tecnológica y farmacológica al mundo de la medicina ha modificado de forma

sustancial el proceso diagnóstico y terapéutico de un gran número de enfermedades.

Con el desarrollo de nuevas moléculas, el pronóstico de muchas neoplasias ha cambiado drásticamente en los últimos años, objetivándose un aumento en la supervivencia. Esta situación supone una dificultad al momento de valorar el ingreso de los pacientes en las Unidades de Cuidados Intensivos (UCI), ya que se debe evitar englobar a todos los pacientes con cáncer dentro de un mismo grupo y favorecer la individualización del tratamiento¹.

Al margen de la razón que motiva el ingreso en la UCI, este subgrupo de pacientes puede beneficiarse de la creación de equipos multidisciplinares en su manejo, lo cual pasa por estrechar lazos entre distintas especialidades que