



REVIEW

Alveolar recruitment maneuvers in respiratory distress syndrome[☆]

Á. Algaba^a, N. Nin^{a,b,c,d,*}, in representation of the GT-IRA of the SEMICYUC

^a Unidad de Cuidados Intensivos, Hospital Universitario de Torrejón, Madrid, Spain

^b Ciber de Enfermedades Respiratorias (CIBERES), ISCIII, Madrid, Spain

^c Fundación para la Investigación Biomédica, Hospital Universitario de Getafe, Madrid, Spain

^d Departamento de Fisiopatología, Hospital de Clínicas, UDELAR, Montevideo, Uruguay

Received 13 December 2012; accepted 30 January 2013

Available online 7 September 2013

KEYWORDS

Acute respiratory distress syndrome;
Recruitment;
Mechanical ventilation

Abstract In patients with acute respiratory distress syndrome, heterogeneity in filling of the lung parenchyma results in collapsed or distended lung areas. Protective ventilation strategies based on the use of low volumes have been shown to increase survival in this context. For opening the lung, and in addition to PEEP, recruitment maneuvers are used—this practice remained the subject of debate.

The present review offers an update on the alveolar recruitment techniques, considering the great variability that exists in the application of these maneuvers, and the different factors that influence the response to maneuvering.

© 2012 Elsevier España, S.L. and SEMICYUC. All rights reserved.

PALABRAS CLAVE

Síndrome de distrés respiratorio agudo;
Reclutamiento;
Ventilación mecánica

Maniobras de reclutamiento alveolar en el síndrome de distrés respiratorio agudo

Resumen En los pacientes con síndrome de distrés respiratorio agudo, la heterogeneidad en el llenado del parénquima pulmonar da lugar a que existan tanto áreas distendidas como colapsadas. Las estrategias de ventilación protectora basadas en el empleo de volúmenes bajos han mostrado en este contexto un aumento de supervivencia. Para abrir el pulmón, además de usar la PEEP, se emplean las maniobras de reclutamiento, todavía en debate.

En la presente revisión se analizan los fundamentos y técnicas para realizar reclutamiento alveolar, considerando la gran variabilidad que existe en cuanto a cómo aplicarlas y los distintos factores que influyen en la respuesta a las mismas.

© 2012 Elsevier España, S.L. y SEMICYUC. Todos los derechos reservados.

Background

Acute respiratory distress syndrome (ARDS) remains an important cause of severe respiratory failure, with a mortality rate of up to 30–60% according to different studies.^{1,2} It has been estimated that patients with ARDS represent up to 10–15% of all patients admitted to the Intensive Care Unit

* Please cite this article as: Algaba Á, Nin N. Maniobras de reclutamiento alveolar en el síndrome de distrés respiratorio agudo. Med Intensiva. 2013;37:355–362.

* Corresponding author.

E-mail address: [nин@торрежонсалуд.ком](mailto:nin@torrejonsalud.com) (N. Nin).

(ICU), and 20% of those who require mechanical ventilation during more than 24 h.³

The *babylung* concept refers to the great lung parenchyma heterogeneity that characterizes ARDS. Some lung regions are relatively well aerated and participate in gas exchange, while others are collapsed as a result of the inflammatory contents within the alveoli or because of the increase in interstitial pressure and the weight of the lung tissue. In this way, in the dependent pulmonary zones, with the patient in supine decubitus at dorsal level, aeration is poorer than in the non-dependent zones at sternal level. This results in very heterogeneous alveolar filling. Much of the research in relation to the treatment of ARDS has been designed to define protective ventilation strategies based on the use of low tidal volumes, which have been shown to improve patient survival.⁴ However, such volumes can favor progressive alveolar collapse due to cyclic opening and closing, which further increases the lung damage. The application of positive end-expiratory pressure (PEEP) can stabilize the alveolus by avoiding the continuous repetition of parenchymal aperture and collapse. Based on the evidence in the literature, modifications in the prescribed mechanical ventilation parameters have been introduced over the years, with the use of lesser tidal volumes and higher PEEP levels.⁵ While there is considerable agreement in calculating tidal volume, it is less clear which PEEP levels should be applied on an individual basis. A number of trials have been carried out analyzing the use of high or low PEEP in patients with ARDS—the results suggesting that high PEEP is associated with increased survival in such individuals.⁶ On the other hand, however, excessive PEEP levels can worsen the damage by distending regions that are already open, giving rise to adverse hemodynamic effects.⁷ In order to “open the lung and keep it open”,⁸ PEEP has been combined with alveolar recruitment (AR) maneuvers—the efficacy of which remains subject to debate.^{9,10}

Definition and physiopathology

Alveolar recruitment is defined as the re-expansion of previously collapsed lung areas by means of a brief and controlled increase in transpulmonary pressure.¹¹ The idea of AR is to create and maintain a collapse-free situation with the purpose of increasing the end-expiration volume and improve gas exchange.

Since the 1970s, different experimental studies have investigated the relationship between alveolar volume and pressure and alveolar shape and size, and the ways in which volume changes affect alveolar structure.¹² In 1952, Day et al. applied different pressure levels to revertatelectasis in animal lungs. These authors found that low pressures are not effective even if maintained for prolonged periods of time, while high pressures are able to open the lungs but cause damage if maintained for a long time.¹³ They concluded that a minimum pressure threshold must be exceeded in order to open an atelectatic lung, and that doing so safely requires precise control of the duration of application of such pressure.

Alveolar recruitment therefore has two fundamental components: the pressure level applied, and the time

during which it is maintained.¹⁴ The increase in transpulmonary pressure (transalveolar pressure would be a more accurate term) opens the terminal alveolar units according to their critical opening pressure, which varies greatly depending on their location. It has been seen that critical opening pressure is low in non-dependent regions, high in dependent regions, and intermediate in limiting territories.¹⁵ Based on the “open lung” theory, it is considered that the entire lung mass could be reopened in the early stages of ARDS if sufficient transalveolar pressure is applied.⁸ According to mathematical and experimental models, airway pressures of over 40 cmH₂O must be applied in order to achieve full recruitment.

In some studies, AR maneuvering has been made under radiological guidance using computed axial tomography (CAT). Gattinoni et al. studied the relationship between the percentage of potentially recruitable lung (as measured by CAT) and the clinical and physiological effects of different PEEP levels in 68 patients with ARDS.¹⁶ They found that the percentage of potentially recruitable lung varies greatly from one patient to another, with an average of 13 ± 11%, and that this parameter is related to PEEP response. As a result, it would be very useful to know the lung recruitment capacity before prescribing the ventilation parameters. Patients with greater recruitable tissue showed poorer oxygenation and compliance, a greater dead space, and increased mortality. However, it is not possible to determine the recruitment capacity of a given patient at the bedside. Costa et al. developed an impedance tomography-based algorithm for estimating recruitable alveolar collapse and hyperdistension, similar to that of CAT but without having to move the patient.¹⁷ This algorithm allowed individualized PEEP titration. Other studies have analyzed the pressure–volume curve and especially its hysteresis as a predictor of lung recruitment capacity.^{15,18} It has been seen that recruitment occurs along the entire pressure–volume curve, even above the upper inflection point.¹⁵ Hysteresis intrinsically reflects the recruited volume; as a result, increased curve hysteresis implies increased AR capacity.¹⁸

In relation to the use of complementary imaging techniques, Tomicic et al. distinguished between anatomical and functional recruitment.¹⁹ Anatomical recruitment refers to the lung tissue in which collapse is reverted, and which can be evaluated by CAT. Functional recruitment in turn is related to improvement of intrapulmonary shunting. Aerating previously collapsed lung zones does not directly imply improved gas exchange, since during partial recruitment part of the perfusion of these alveolar units may be displaced toward other collapsed units, thereby countering both effects. Increased oxygenation will depend on the changes produced in the ventilation–perfusion ratio. Independently of the effect upon oxygenation, it is considered that AR, by increasing the aerated tissue, contributes to minimize the heterogeneity of the lung and avoid cyclic opening and closing. This in turn can prevent ventilator-associated lung injury.²⁰

How alveolar recruitment is performed

The techniques used to perform AR and the results obtained vary greatly among the different studies, in terms of both

Table 1 Techniques used for alveolar recruitment and results obtained.

Author (year)	n	Recruitment technique	Timing of recruitment	Main results	Adverse effects
Amato et al. ²³ (1998)	53 (AR in 29 patients)	Within protective strategy, recruitment maneuvers: CPAP 35–40 cmH ₂ O during 40 s	Frequently at the start of the protocol and after disconnecting respirator (maximum one maneuver a day)	Decrease in mortality in protective ventilation group Greater weaning rate	None
Lapinsky et al. ²⁴ (1999)	14	30–45 cmH ₂ O during 20 s	Not specified	Brief increase in oxygenation	None of significance
Grasso et al. ²⁵ (2002)	22	40 cmH ₂ O CPAP 40 s	Not specified	Oxygenation improved only in patients with early phase ARDS and no chest wall involvement	Decrease in MAP and cardiac output in non-responders
Brower et al. ²⁶ (2003)	72	CPAP 35–40 cmH ₂ O during 30 s	Alternate days AR maneuvers/sham maneuvers	Very brief and mild improvement in saturation	Increased drop in AP One case of barotrauma
Dyhr et al. ²⁷ (2003)	8	CPAP 45 cmH ₂ O during 20 s 2 times, with 1 min between both	After aspirations	Overcomes oxygenation drop that occurs on aspirating	Not specified
Oczenski et al. ²⁸ (2004)	30	CPAP 50 cmH ₂ O during 30 s	Once	Very brief increase in PaO ₂ /FiO ₂ ratio (less than 30 min)	None of significance
Girgis et al. ²⁹ (2006)	20	CPAP 40 cmH ₂ O during 40 s	Three times followed by gradual reduction of PEEP and seeking optimum level	Improved PaO ₂ /FiO ₂ ratio	Not specified
Meade et al. ³⁰ (2008)	983	Sustained CPAP 40 cmH ₂ O during 40 s, within ventilation strategy with open lung	At start and after disconnections (up to 4 a day)	Nonsignificant decrease in mortality. Improved secondary endpoints related to hypoxemia	No significant increase in barotrauma
Pelosi et al. ³¹ (1999)	10	Three consecutive sighs/minute at 45 cmH ₂ O plateau pressure, during one hour, with protective ventilation	Ventilation protocol with several phases	Increased PaO ₂ and lung volume at end of expiration; drop in PaCO ₂ , greater effect in extrapulmonary ARDS	None of significance
Patroniti et al. ³² (2002)	13	Sighs: CPAP 20% > peak pressure or at least 35 cmH ₂ O in supportive pressure 3–5 s each min, during 1 h	Ventilation protocol with several phases	Increased oxygenation, lung volume and compliance	Not specified
Pelosi et al. ³³ (2003)	10	Three sighs 45 cmH ₂ O each min during 1 h with ventilation in prone decubitus	Ventilation protocol with several phases	Increased end-expiration volume and PaO ₂ with decreased shunt	None of significance

Table 1 (Continued)

Author (year)	n	Recruitment technique	Timing of recruitment	Main results	Adverse effects
Lim et al. ³⁴ (2001)	20	In 3 steps Vt is reduced (8 to 2 ml/kg) and PEEP is raised (10–25)	Two consecutive times with 1 min interval	Increase in PaO ₂ and static compliance	None of significance
Borges et al. ³⁶ (2006)	26	Maximum recruitment maneuvers in pressure control, guided by CAT	Upon admission	Increase in PaO ₂ /FiO ₂ , inversely related to percentage collapsed parenchyma	Transient decrease in cardiac output with no clinical repercussions
De Matos et al. ³⁸ (2012)	51	Maximum recruitment maneuvers in pressure control, guided by CAT	Upon admission and after, if Vt decreases 20% or respirator is disconnected	Sustained increase in PaO ₂ /FiO ₂ and decrease in non-aerated lung tissue measured by CAT	None of significance
Monge Garcia et al. ⁴¹ (2012)	21	Progressive increase in PEEP to 36 cmH ₂ O	Not specified	Increase in distensibility and PaO ₂ /FiO ₂	Decrease in cardiac output

CPAP: continuous positive airway pressure; FiO₂: oxygen concentration in inhaled air; AP: arterial pressure; MAP: mean arterial pressure; PaO₂: partial pressure of oxygen; PaCO₂: partial pressure of carbon dioxide; PEEP: positive end-expiratory pressure; AR: acute respiratory distress syndrome; CAT: computed axial tomography; Vt: tidal volume.

the timing and the duration of application (**Table 1**). Both conventional and alternative mechanical ventilation modes can be used.

The conventional ventilation modes can be divided into four large groups^{20–22}:

1. Sustained continuous positive airway pressure (CPAP): a certain pressure, usually 35–50 cmH₂O, is reached during 20–40 s. The most frequent combination is the application of 40 cmH₂O during 40 s. During this time the supporting pressure must be set to 0 cmH₂O in order to avoid barotrauma. This is the most widely used technique.^{23–30}
2. Sigh: increased tidal volume or PEEP during one or several respirations, with adjustment designed to reach a specific plateau pressure.^{31–33}
3. Extended sigh: this considers the interaction between pressure and time, and is characterized by a progressive increase in PEEP, together with a decrease in tidal volume during a longer period of time.^{34,35}
4. Pressure controlled ventilation, maintaining a pressure delta (usually 15 cmH₂O) to guarantee a tidal volume, with progressive increases in PEEP. Some authors have used so-called maximum recruitment maneuvers, reaching much higher pressures though in a gradual manner, and which in some cases are followed by a gradual decrease in pressure that serves to define the optimum individual PEEP following lung recruitment.^{36–38}

Likewise related to pressure controlled ventilation, "open lung" ventilation is based on early lung recruitment maneuvering, with sufficient PEEP to open the largest possible number of alveoli and ventilating with the lowest possible lung distension pressure.^{39–41}

Other techniques

- High-frequency ventilation: compared with the conventional mechanical ventilation modes, high-frequency ventilation gives rise to higher mean airway pressures, which limits cyclic alveolar closure and increases the end-expiration lung volume.⁴² Some studies have evaluated the combination of high-frequency ventilation with sustained CPAP maneuvering with a view to reinforcing its effect in AR.⁴³
- Liquid ventilation: from the purely theoretical point of view, partial lung filling with perfluorocarbon gives rise to progressive parenchyma recruitment, beginning in the dependent zones. Studies have been made in animals of the distribution of liquid and gas with different ventilation pressures.⁴⁴ This technique has not been found to offer improvement in patients.⁴⁵
- Ventilation in prone decubitus: positioning the patient in prone decubitus varies the distribution of the transpulmonary pressure gradient, giving rise to more homogeneous alveolar filling. It does not substantially modify perfusion, however, and the ventilation-perfusion ratio is therefore improved as a result.⁴⁶ Prone decubitus may be regarded as a form of recruitment in itself. If in addition some of the commented conventional ventilation techniques are used, more uniform expansion of the applied pressures will result.^{33,46,47}

Heterogeneity of the studies

A range of studies in both animals and humans have applied different AR techniques, with very diverse protocols. The studies are in fact so heterogeneous in terms of the maneuvers employed, the type of patients involved, the parameters, etc., such that it is impossible to compare the different recruitment techniques. Constantin et al. compared CPAP 40 cmH₂O during 40 s with prolonged sigh, and found the latter to afford comparatively greater oxygenation with a greater recruitment of lung tissue.⁴⁸ Mahmoud and Ammar likewise compared sustained CPAP with prolonged sigh, and reported better results and tolerance with the latter approach.⁴⁹ In 12 patients with ARDS and protective ventilation, Badet et al. applied three different forms of AR after analyzing the optimum PEEP: ventilation with the optimum PEEP level, adding sustained insufflation, and with sighs.⁵⁰ These authors recorded a greater increase in oxygenation and static compliance with the third recruitment modality, i.e., upon adding sighs. In recent years, experimental studies have found that slower and more progressive recruitment maneuvers, raising the pressure over several steps or in the form of a ramp until reaching the target level, offer better results with less hemodynamic impairment.⁵¹ In patients, pressure ventilation with pressure increments and decrements operates in this same line.⁵²

Ideally, recruitment maneuvering should be performed with the patient under sedation-relaxation, a FiO₂ of 100% and under hemodynamically stable conditions. The meta-analysis carried out by Fan et al. shows that these conditions are often not met or are not documented.²¹ Likewise, there are no studies describing the percentage of patients with ARDS in which AR is applied. All these factors greatly complicate the drawing of conclusions regarding the usefulness of AR maneuvering in patients with ARDS.

Variability of response to recruitment maneuvers

The response to AR varies according to different factors referred to both the patient and the characteristics of ARDS:

1. Origin of ARDS: Differences have been found between ARDS of pulmonary and extrapulmonary origin, with greater efficacy in extrapulmonary presentations.^{15,31,35,52} This effect appears to be due to less intraalveolar involvement, at least initially, with a predominance of interstitial edema, which appears to be associated with a greater recruitment capacity. However, occupation of the interior of the alveoli, as for example in pneumonia, implies the existence of less recruitable tissue. In contrast, Borges et al. have reported a similar recruitment capacity, independently of the cause of ARDS, on performing maximum recruitment maneuvers.³⁶
2. Evolutive phase: Some authors consider recruitment to be effective only when performed in the early phases of lung injury, since in later phases the impaired elasticity does not allow the reversion of collapse, and increases the risk of barotrauma.²⁵

3. Patient positioning: As has been commented, this factor has a strong effect upon recruitment response—the latter being greater in prone decubitus, which in itself could be regarded as a form of AR, since it increases transpulmonary pressure in the dorsal region and improves gas exchange.^{46,52}
4. Vasoactive drug treatment, by modifying cardiac output, the distribution of pulmonary blood flow and gas exchange, theoretically could also modify the response to AR.²²
5. Chest expansion capacity: Grasso et al. found AR maneuvering to be ineffective in patients with limited chest expansion.²⁵
6. Previous mechanical ventilation parameters: Ventilation with low tidal volumes can cause alveolar closure, which could be compensated by using an adequate PEEP level. It has been observed that the use of higher tidal volumes and particularly of higher PEEP values before AR is associated with a lesser response to recruitment, since maneuvering probably already starts from a recruited lung.^{35,53,54}
7. Posterior mechanical ventilation parameters: The posterior mechanical ventilation strategy, particularly as regards PEEP level, is as important as the recruitment technique used.⁵⁵

Indications

In routine practice, AR maneuvering is carried out in situations of severe hypoxemia, as a rescue measure; consequently, patients with ARDS are typically involved. There is no evidence regarding when maneuvering is indicated—in what concrete moments or with what frequency. In some studies, AR maneuvering is performed systematically, while in others it is performed only when the lung is considered to have been "de-recruited"—fundamentally after disconnecting the patient from the respiratory for some reason, such as for example the aspiration of secretions.²³

Maximum recruitment strategies and the open lung theory advocate the early application of ventilation protocols across different phases, in order to achieve adequate lung opening from the start and mechanical ventilation with individualized parameters, particularly as refers to optimum PEEP derived from decremental testing after AR.^{36–38} In this way, AR not only aims to revert a point situation of hypoxemia but moreover also forms part of the global measures for reducing the lung damage associated to mechanical ventilation.²⁰

Outside the context of Intensive Care Medicine, AR plays an important role in the operating room. In effect, AR maneuvering can be of benefit in opening atelectatic areas associated with anesthesia, particularly in obese patients, and during the immediate postoperative period of certain operations involving a high risk of respiratory complications.⁵⁶

Results

On analyzing the results of AR, it must be mentioned that most studies do not establish criteria for defining a positive response to AR maneuvering, and only describe

variable improvements in oxygenation. Grasso et al. classified responders as those patients in which the $\text{PaO}_2/\text{FiO}_2$ ratio increased at least 50% after AR maneuvering,²⁵ while Villagrá et al.⁵⁴ and Girgis et al.²⁹ accepted a much lesser increase (20%) as representing a positive response. It has been reported that maximum recruitment can be assumed with a FiO_2 of 100%, if $\text{PaO}_2 + \text{PaCO}_2 = 400$, or if the $\text{PaO}_2/\text{FiO}_2$ ratio is >350, with a collapsed lung tissue mass of less than 5%.³⁶ Instead of focusing on oxygenation-related outcomes, some studies define response in terms of lung mechanics, analyzing the effect of AR upon parameters such as compliance.

Many studies have reported improved oxygenation as a result of AR maneuvering in both basic experimentation in animals, in which different techniques are used to induce lung injury,⁵⁷ and in patients with ARDS.^{23–25,27–38,58} However, some authors have described no beneficial effects.²⁶

The increase in oxygenation is usually brief, and some studies have found the effect to fade after as little as 15–30 min²⁸—though the duration of effect is usually about 3–6 h. In 51 patients with severe ARDS, De Matos et al. prescribed pressure ventilation according to the open lung theory, using maximum recruitment techniques, and found the benefits in terms of gas exchange and respiratory mechanics to persist for several days.³⁸ The authors consider that the pressures classically reached during AR maneuvering (40–45 cmH₂O) are not enough to aerate the collapsed lung in severe ARDS, and that the capacity of AR should be reconsidered on the basis of more “aggressive” techniques, though applying them on a progressive basis.

Amato et al. reported a decrease in mortality in the group of patients subjected to AR maneuvering in the context of protective ventilation; the outcome therefore must be related to the series of adopted measures, not directly to AR.²³ The increase in oxygenation observed as a result of AR has not been found to be accompanied by any relevant impact upon the clinical outcomes. There are no differences in terms of the duration of mechanical ventilation, hospital stay or mortality on performing this technique.⁵⁸

Side effects

The increase in transpulmonary pressure is sometimes accompanied by adverse effects—the most frequently reported being hypotension and desaturation.^{25,26} In the studies that have included more complete invasive hemodynamic monitoring, decreases have been described in cardiac output, systolic volume and preload, together with a rise in heart rate.^{25,36,41} The invasive monitoring of arterial pressure is not enough to evaluate all the hemodynamic changes that occur with recruitment maneuvering, since the values may even increase temporarily.

Other complications are barotrauma, arrhythmias, hypoventilation and acidosis, and it has even been postulated that bacterial translocation can occur from within the alveoli—though the results of the different studies are contradictory in this sense.⁵⁹ In general, such effects are usually brief and of scant importance; a change in ventilation strategy is therefore usually not required.

It has been observed that when sustained insufflation maneuvers are carried out, side effects are more frequent

than with other techniques. In the case of maneuvering with pressure ventilation, there appear to be fewer cases of barotrauma or adverse hemodynamic effects, despite the fact that much higher pressures are reached in the airway, since maneuvering is applied in a more progressive manner.⁵² Gil Cano et al. analyzed the incidence of barotrauma in 100 patients ventilated using an open lung strategy.⁶⁰ They described pneumothorax in 7 patients and subcutaneous emphysema in two individuals—all of which presented primary ARDS. No changes in the mechanical ventilation parameters proved necessary.

Conclusions

On the basis of the existing evidence, AR maneuvering cannot be recommended as a general practice in clinically stable ARDS patients. Such maneuvering usually affords variable and temporary improvement in oxygenation, without beneficial effects upon mortality. The potential effects of AR depend on a series of factors referred to both the patient and the ventilation parameters before and after application of the technique. In this context, the PEEP level prescribed after maneuvering, with a view to avoiding “derecruitment”, is of crucial importance.

The more recently introduced AR maneuvers, which reach higher airway pressures in a more progressive manner, can prove useful during mechanical ventilation in seeking an optimum PEEP level; however, it is not clear whether the use of this parameter is accompanied by beneficial clinical outcomes.

Repeated AR maneuvering should not be attempted in non-responders, since it may have adverse effects.

Financial support

This study has been supported by *Ciber de Enfermedades Respiratorias*, FIS PI11/02791.

Conflict of interest

The authors declare that they have no conflicts of interest.

References

1. Villar J, Blanco J, Añón JM, Santos-Bouza A, Blanch L, Ambrós A, et al. The ALIEN study: incidence and outcome of acute respiratory distress syndrome in the era of lung protective ventilation. *Intensive Care Med.* 2011;37:1932–41.
2. Rubenfeld GD, Herridge MS. Epidemiology and outcomes of acute lung injury. *Chest.* 2007;131:554–62.
3. Frutos-Vivar F, Nin N, Esteban A. Epidemiology of acute lung injury and acute respiratory distress syndrome. *Curr Opin Crit Care.* 2004;10:1–6.
4. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The Acute Respiratory Distress Syndrome Network. *N Engl J Med.* 2000;342:1301–8.
5. Esteban A, Ferguson ND, Meade MO, Frutos F, Apezteguía C, Brochard L, et al. Evolution of mechanical ventilation in response to clinical research. *Am J Respir Crit Care Med.* 2008;177:170–7.

6. Briel M, Meade M, Mercat A, Brower RG, Talmor D, Walter SD, et al. Higher vs lower positive end-expiratory pressure in patients with acute lung injury and acute respiratory distress syndrome: systematic review and meta-analysis. *J Am Med Assoc.* 2010;303:865–73.
7. Slutsky AS, Hudson LD. PEEP or no PEEP—lung recruitment may be the solution. *N Engl J Med.* 2006;354:1839–41.
8. Lachmann B. Open up the lung and keep the lung open. *Intensive Care Med.* 1992;18:319–21.
9. Suárez Sipmann F. Utilidad de las maniobras de reclutamiento (pro). *Med Intensiva.* 2009;33:134–8.
10. Ochagavia A, Blanch L, López-Aguilar J. Utilidad de las maniobras de reclutamiento (contra). *Med Intensiva.* 2009;33:139–43.
11. Richard JC, Maggiore SM, Mercat A. Clinical review: bedside assessment of alveolar recruitment. *Crit Care.* 2004;8:163–9.
12. Klingele TG, Staub NC. Alveolar shape changes with volume in isolated, air-filled lobes of cat lung. *J Appl Physiol.* 1970;28:411–4.
13. Day R, Goodfellow AM, Apgar V, Beck GJ. Pressure-time relations in the safe correction of atelectasis in animal lungs. *Pediatrics.* 1952;10:593–602.
14. Albert SP, DiRocco J, Allen GB, Bates JH, Lafollette R, Kubiak BD, et al. The role of time and pressure on alveolar recruitment. *J Appl Physiol.* 2009;106:757–65.
15. Crotti S, Mascheroni D, Caironi P, Pelosi P, Ronzoni G, Mondino M, et al. Recruitment and derecruitment during acute respiratory failure: a clinical study. *Am J Respir Crit Care Med.* 2001;164:131–40.
16. Gattinoni L, Caironi P, Cressoni M, Chiumello D, Ranieri M, Quintel M, et al. Lung recruitment in patients with the acute respiratory distress syndrome. *N Engl J Med.* 2006;354:1775–86.
17. Costa EL, Borges JB, Melo A, Suarez-Sipmann F, Toufen C, Bohm SH, et al. Bedside estimation of recruitable alveolar collapse and hyperdistension by electrical impedance tomography. *Intensive Care Med.* 2009;35:1132–7.
18. Demory D, Arnal JM, Wysocki M, Donati S, Granier I, Corno G, et al. Recruitability of the lung estimated by the pressure volume curve hysteresis in ARDS patients. *Intensive Care Med.* 2008;34:2019–25.
19. Tomicic V, Fuentealba A, Martínez E, Graf J, Batista Borges J. Fundamentos de la ventilación mecánica en el síndrome de distrés respiratorio agudo. *Med Intensiva.* 2010;34:418–27.
20. Lapinsky SE, Mehta S. Bench-to-bedside review: recruitment and recruiting maneuvers. *Crit Care.* 2005;9:60–5.
21. Fan E, Wilcox ME, Brower RG, Stewart TE, Mehta S, Lapinsky SE, et al. Recruitment maneuvers for acute lung injury: a systematic review. *Am J Respir Crit Care Med.* 2008;178:1156–63.
22. Guerin C, Debord S, Leray V, Delannoy B, Bayle F, Bourdin G, et al. Efficacy and safety of recruitment maneuvers in acute respiratory distress syndrome. *Ann Intensive Care.* 2011;1:9.
23. Amato MB, Barbas CS, Medeiros DM, Magaldi RB, Schettino GP, Lorenzi-Filho G, et al. Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome. *N Engl J Med.* 1998;338:347–54.
24. Lapinsky SE, Aubin M, Mehta S, Boiteau P, Slutsky AS. Safety and efficacy of a sustained inflation for alveolar recruitment in adults with respiratory failure. *Intensive Care Med.* 1999;25:1297–301.
25. Grasso S, Mascia L, del Turco M, Malacarne P, Giunta F, Brochard L, et al. Effects of recruiting maneuvers in patients with acute respiratory distress syndrome ventilated with protective ventilatory strategy. *Anesthesiology.* 2002;96:795–802.
26. Brower RG, Morris A, MacIntyre N, Matthay MA, Hayden D, Thompson T, et al. Effects of recruitment maneuvers in patients with acute lung injury and acute respiratory distress syndrome ventilated with high positive end-expiratory pressure. *Crit Care Med.* 2003;31:2592–7.
27. Dyhr T, Bonde J, Larsson A. Lung recruitment manoeuvres are effective in regaining lung volume and oxygenation after open endotracheal suctioning in acute respiratory distress syndrome. *Critical Care.* 2003;7:55–62.
28. Oczenski W, Hörmann C, Keller C, Lorenzl N, Kepka A, Schwarz S, et al. Recruitment maneuvers after a positive end-expiratory pressure trial do not induce sustained effects in early adult respiratory distress syndrome. *Anesthesiology.* 2004;101:620–5.
29. Girgis K, Hamed H, Khater Y, Kacmarek RM. A decremental PEEP trial identifies the PEEP level that maintains oxygenation after lung recruitment. *Respir Care.* 2006;51:1132–9.
30. Meade MO, Cook DJ, Guyatt GH, Slutsky AS, Arabi YM, Cooper DJ, et al. Ventilation strategy using low tidal volumes, recruitment maneuvers, and high positive end-expiratory pressure for acute lung injury and acute respiratory distress syndrome: a randomized controlled trial. *J Am Med Assoc.* 2008;299:637–45.
31. Pelosi P, Cadrinher P, Bottino N, Panigada M, Carriero F, Riva E, et al. Sigh in acute respiratory distress syndrome. *Am J Respir Crit Care Med.* 1999;159:872–80.
32. Patroniti N, Foti G, Cortinovis B, Maggioni E, Bigatello LM, Cereda M, et al. Sigh improves gas exchange and lung volume in patients with acute respiratory distress syndrome undergoing pressure support ventilation. *Anesthesiology.* 2002;96:788–94.
33. Pelosi P, Bottino N, Chiumello D, Caironi P, Panigada M, Gamberoni C, et al. Sigh in supine and prone position during acute respiratory distress syndrome. *Am J Respir Crit Care Med.* 2003;167:521–7.
34. Lim CM, Koh Y, Park W, Chin JY, Shim TS, Lee SD, et al. Mechanistic scheme and effect of “extended sigh” as a recruitment maneuver in patients with acute respiratory distress syndrome: a preliminary study. *Crit Care Med.* 2001;29:1255–60.
35. Lim CM, Jung H, Koh Y, Lee JS, Shim TS, Lee SD, et al. Effect of alveolar recruitment maneuver in early acute respiratory distress syndrome according to antiderecruitment strategy, etiological category of diffuse lung injury, and body position of the patient. *Crit Care Med.* 2003;31:411–8.
36. Borges JB, Okamoto VN, Matos GF, Caramez MP, Arantes PR, Barros F, et al. Reversibility of lung collapse and hypoxemia in early acute respiratory distress syndrome. *Am J Respir Crit Care Med.* 2006;174:268–78.
37. Morán I, Blanch L, Fernández R, Fernández-Mondéjar E, Zavala E, Mancebo J. Acute physiologic effects of a stepwise recruitment maneuver in acute respiratory distress syndrome. *Minerva Anestesiol.* 2011;77:1167–75.
38. De Matos GF, Stanzani F, Passos RH, Fontana MF, Albaladejo R, Caserta RE, et al. How large is the lung recruitability in early acute respiratory distress syndrome: a prospective case series of patients monitored by computed tomography. *Crit Care.* 2012;16:R4.
39. Amato MB, Barbas CS, Medeiros DM, Schettino Gde P, Lorenzi Filho G, Kairalla RA, et al. Beneficial effects of the “open lung approach” with low distending pressures in acute respiratory distress syndrome. A prospective randomized study on mechanical ventilation. *Am J Respir Crit Care Med.* 1995;152:1835–46.
40. Papadakos PJ, Lachmann B. The open lung concept of mechanical ventilation: the role of recruitment and stabilization. *Crit Care Clin.* 2007;23:241–50.
41. Monge García MI, Gil Cano A, Gracia Romero M, Díaz Monrové JC. Cambios respiratorios y hemodinámicos durante una maniobra de reclutamiento pulmonar mediante incrementos y decrementos progresivos de PEEP. *Med Intensiva.* 2012;36:77–88.
42. Chan KP, Stewart TE, Mehta S. High-frequency oscillatory ventilation for adult patients with ARDS. *Chest.* 2007;131:1907–16.
43. Ferguson ND, Chiche J-D, Kacmarek RM, Hallett DC, Mehta S, Findlay GP, et al. Combining high-frequency oscillatory ventilation and recruitment maneuvers in adults with early acute respiratory distress syndrome: the Treatment with Oscillation

- and an Open Lung Strategy (TOOLS) Trial pilot study. *Crit Care Med.* 2005;33:479–86.
- 44. Morris K, Cox P, Frndova H, Holowka S, Babyn P. Effect of a sustained inflation on regional distribution of gas and perflurocarbon during partial liquid ventilation. *Pediatr Pulmonol.* 2007;42:204–9.
 - 45. Kacmarek RM, Wiedemann HP, Lavin PT, Wedel MK, Tütüncü AS, Slutsky AS. Partial liquid ventilation in adult patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med.* 2006;173:882–9.
 - 46. Martínez O, Nin N, Esteban A. Evidencias de la posición en decúbito prono para el tratamiento del síndrome de distres respiratorio agudo: una puesta al día. *Arch Bronconeumol.* 2009;45:291–6.
 - 47. Rival G, Patry C, Floret N, Navellou JC, Belle E, Capellier G. Prone position and recruitment manoeuvre: the combined effect improves oxygenation. *Crit Care.* 2011;15:R125.
 - 48. Constantin JM, Jaber S, Futier E, Cayot-Constantin S, Verny-Pic M, Jung B, et al. Respiratory effects of different recruitment maneuvers in acute respiratory distress syndrome. *Crit Care.* 2008;12:R50.
 - 49. Mahmoud KM, Ammar AS. A comparison between two different alveolar recruitment maneuvers in patients with acute respiratory distress syndrome. *Int J Crit Illn Inj Sci.* 2011;1:114–20.
 - 50. Badet M, Bayle F, Richard JC, Guérin C. Comparison of optimal positive end-expiratory pressure and recruitment maneuvers during lung-protective mechanical ventilation in patients with acute lung injury/acute respiratory distress syndrome. *Respir Care.* 2009;54:847–54.
 - 51. Rzezinskia AF, Oliveira GP, Santiago VR, Santos RS, Ornelas DS, Morales MM, et al. Prolonged recruitment manoeuvre improves lung function with less ultrastructural damage in experimental mild acute lung injury. *Respir Physiol Neurobiol.* 2009;169:271–81.
 - 52. Pelosi P, Gama de Abreu M, Rocco PR. New and conventional strategies for lung recruitment in acute respiratory distress syndrome. *Crit Care.* 2010;14:210.
 - 53. Foti G, Cereda M, Sparacino ME, de Marchi L, Villa F, Pesenti A. Effects of periodic lung recruitment maneuvers on gas exchange and respiratory mechanics in mechanically ventilated acute respiratory distress syndrome (ARDS) patients. *Intensive Care Med.* 2000;26:501–7.
 - 54. Villagrá A, Ochagavía A, Vatua S, Murias G, del Mar Fernández M, Lopez Aguilar J, et al. Recruitment maneuvers during lung protective ventilation in acute respiratory distress syndrome. *Am J Respir Crit Care Med.* 2002;165:165–70.
 - 55. Kacmarek RM, Villar J. Lung recruitment maneuvers during acute respiratory distress syndrome: is it useful. *Minerva Anestesiol.* 2011;77:85–9.
 - 56. Tusman G, Böhm SH, Vazquez de Anda GF, do Campo JL, Lachmann B. "Alveolar recruitment strategy" improves arterial oxygenation during general anaesthesia. *Br J Anaesth.* 1999;82:8–13.
 - 57. Kloot TE, Blanch L, Melynne Youngblood A, Weinert C, Adams AB, Marini JJ, et al. Recruitment maneuvers in three experimental models of acute lung injury. Effect on lung volume and gas exchange. *Am J Respir Crit Care Med.* 2000;161:1485–94.
 - 58. Hodgson C, Keating JL, Holland AE, Davies AR, Smirneos L, Bradley SJ, et al. Recruitment manoeuvres for adults with acute lung injury receiving mechanical ventilation. *Cochrane Database Syst Rev.* 2009;CD006667.
 - 59. Piacentini E, Villagrá A, López-Aguilar J, Blanch L. Clinical review: the implications of experimental and clinical studies of recruitment maneuvers in acute lung injury. *Critical Care.* 2004;8:115–21.
 - 60. Gil Cano A, Monge García MI, Gracia Romero M, Díaz Monrové JC. Incidencia, características y evolución del barotrauma durante la ventilación mecánica con apertura pulmonar. *Med Intensiva.* 2012;36:335–42.