

6. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. Crit Care Med. 2003;31:1250–6.
7. Egea-Guerrero JJ, Martínez-Fernández C, Rodríguez-Rodríguez A, Bohórquez-López A, Vilches-Arenas A, Pacheco-Sánchez M, et al. The utility of C-reactive protein and procalcitonin for sepsis diagnosis in critically burned patients: a preliminary study. Plast Surg (Oakv). 2015;23:239–43.

J.J. Egea-Guerrero*, A. Rodríguez-Rodríguez

U.G.C. Cuidados Críticos y Urgencias, H.U. Virgen del Rocío, IBISS/CSIC/Universidad de Sevilla, Sevilla, Spain

*Corresponding author.

E-mail addresses: jjegeaguerrero@gmail.com, juanj.egea.sspa@juntadeandalucia.es (J.J. Egea-Guerrero).

Author's reply



Respuesta de los autores

We read with interest the letter sent by JJ Egea-Guerrero et al. The authors agree to the comment of the review¹ that the routine use of biomarkers cannot be recommended in the differential diagnosis of inflammation from sepsis. The same conclusion has been achieved by Seoane et al. in a recent article.²

Egea-Guerrero et al. refer to one preliminary report about the diagnostic value of C-Reactive Protein and Procalcitonin³ in this population. The authors processed and analyzed 157 samples of 17 severe burn patients. The authors suggested that changes in Procalcitonin over time differentiated patients with sepsis and patients without sepsis only based in the ROC curves, but they did not provide data of sensitivity, specificity and likelihood ratios of different cut-off values that could help clinicians in differential diagnosis. We agree with them that it is necessary new well-designed studies to provide new insights on this topic.

Conflict of interest

None declared.

References

1. Hidalgo F, Mas D, Rubio M, García-Hierro P. Infections in critically ill burn patients. Med Intensiva. 2016;40(3):179–85.
2. Seoane L, Pértiga S, Galeiras R, Astola I, Bouza T. Procalcitonin in the burn unit and the diagnosis of infection. Burns. 2014;40:223–9.
3. Egea-Guerrero JJ, Martínez-Fernández C, Rodríguez-Rodríguez A, Bohórquez-López A, Vilches-Arenas A, Pacheco-Sánchez M, et al. The utility of C-reactive protein and procalcitonin for sepsis diagnosis in critically burned patients: a preliminary study. Plast Surg (Oakv). 2015;23:239–43.

F. Hidalgo ^{a,*}, D. Mas ^b, M. Rubio ^a

^a Department of Critical Care Medicine, Hospital Universitario de Getafe, Madrid, Spain

^b Department of Plastic Surgery, Hospital Universitario de Getafe, Madrid, Spain

* Corresponding author.

E-mail address: fhidalgos@salud.madrid.org (F. Hidalgo).