

- patients in the coronavirus disease (COVID-19). *Med Intensiva*. 2020, <http://dx.doi.org/10.1016/j.medint.2020.04.001>.
4. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, et al. Presenting characteristics comorbidities, and outcomes among 5700 patients hospitalised with COVID-19 in the New York City Area. *JAMA*. 2020, <http://dx.doi.org/10.1001/jama.2020.6775> [Epub ahead of print].
 5. Grasselli G, Zangrillo A, Zanella A, Antonelli M, Cabrini L, Castelli A, et al. Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy Region Italy. *JAMA*. 2020, <http://dx.doi.org/10.1001/jama.2020.5394> [Epub ahead of print].
 6. Blake A, Collins D, O'Connor E, Bergin C, McLaughlin AM, Martin-Lloeches I. Clinical and biochemical characteristics of patients admitted to ICU with SARS-CoV-2. *Med Intensiva*. 2020, <http://dx.doi.org/10.1016/j.medint.2020.05.003>.
 7. Abella Alvarez A, Torrejon Perez I, Enciso Calderon V, Hermosa Gelbard C, Sicilia Urban JJ, Ruiz Grinspan M, et al. ICU without walls project Effect of the early detection of patients at risk. *Med Intensiva*. 2020;37:12–8.

B. Lobo-Valbuena^{a,b,*}, Ma. García-Arias^{c,b}, R.B. Pérez^d, D.V. Delgado^e, F. Gordo^{f,b}



Support to the organization of the Intensive Care Units during the pandemic through maps created from the Clinical Information Systems[☆]

Soporte a la organización de las Unidades de Cuidados Intensivos durante la pandemia, a través de mapas creados a partir de los Sistemas de Información Clínica

Sir,

The current SARS-CoV-2 pandemic poses a major challenge for Intensive Care Units (ICUs).¹ In the last few months, the capacity of these Units has been drastically incremented, with the need to habilitate other hospital areas not initially conceived to hold critically ill patients.²

Sufficient means are clearly needed to deal with future outbreaks. In this regard, dashboards generated from the data contained in the clinical information system (CIS) offer a graphic display of relevant information on the patient condition and course. Dashboards facilitate critical patient care, offering important clinical information intuitively and in a way that is easy to interpret. They generate alerts when key parameters move beyond acceptable limits, allowing

^a Intensivista – Unidad de Cuidados Intensivos del Hospital Universitario del Henares, Coslada, Madrid, Spain

^b Grupo de Investigación en Patología Crítica, Facultad de Ciencias de la Salud, Universidad Francisco de Vitoria, Pozuelo de Alarcón, Madrid, Spain

^c Supervisora de Enfermería de la Unidad de Cuidados Intensivos del Hospital Universitario del Henares, Coslada, Madrid, Spain

^d Anestesista – Servicio de Anestesia del Hospital Universitario del Henares, Coslada, Madrid, Spain

^e Responsable de la Unidad de Apoyo a la Investigación. Facultad de Medicina, Universidad Francisco de Vitoria, Pozuelo de Alarcón, Madrid, Spain

^f Jefe de Servicio de la Unidad de Cuidados Intensivos del Hospital Universitario del Henares, Coslada, Madrid, Spain

* Corresponding author.

E-mail address: b.lobo.valbuena@gmail.com

(B. Lobo-Valbuena).

<https://doi.org/10.1016/j.medint.2020.06.020>

0210-5691/

© 2020 Elsevier España, S.L.U. y SEMICYUC. All rights reserved.

immediate corrective actions and constituting genuine clinical decision supporting tools.³ Likewise, the pandemic has evidenced the need to upgrade the traditional models and to include different forms of “telecommuting” or “remote care” in medical practice.⁴

It is clear that information technology offers an excellent opportunity to improve patient care.⁵ Specifically, CIS integrate the information from other Departments, point of care devices and laboratory test results, drug prescriptions and specific information recorded by the healthcare professionals. Nevertheless, the latter must make important efforts to obtain the required information globally and on a real-time basis in order to ensure effective and efficient organization.⁶

During the peak of the COVID-19 care crisis in our Unit, we developed a series of dashboards from the information available in the CIS that allowed us to monitor the patients on a remote basis, improving the safety and quality of care in moments of maximum healthcare burden. A recent review has shown that the use of dashboards improves efficiency in routine clinical practice, clinician satisfaction, patient safety and reliability of the working environment.⁷

The entire application has been programmed in a controlled environment, independently of the operating system, using Docker. The data are extracted in real time from the CIS through ETL processes using Python, and are displayed via website programmed with the web standards (html, JavaScript and css) using the Django framework.

We first developed a general dashboard allowing us to quickly assess the level of complexity and the life support requirements of each patient in each of the Units (mechanical ventilation, extrarenal replacement therapy or vasoactive support). This dashboard allows rapid visualization of the distribution of resources and the care burden in the Unit.

☆ Please cite this article as: Claverias L, Gómez J, Rodríguez A, Albiol J, Esteban F, Bodí M. Soporte a la organización de las Unidades de Cuidados Intensivos durante la pandemia, a través de mapas creados a partir de los Sistemas de Información Clínica. *Med Intensiva*. 2021;45:58–60.

Servei UCI-1




Figure 1 Treatment and ventilatory support dashboard. We first examine (marked in red) whether the patient is isolated due to SARS-CoV-2 positivity and the duration of UCI and hospital stay. The central panel displays the mechanical ventilation monitoring parameters in colors according to whether they are within the adequate range or not (Vt: green if within range, yellow if > 5%, orange if > 5-10%, red if > 10%; MP: green if ≤ 18 , red if > 18). Lastly, an orange color indicates whether the patient is in prone decubitus and what sedoanalgesia and/or neuromuscular relaxant is being provided in continuous perfusion. Vt: tidal volume; MaxVt: maximum tidal volume; DP: driving pressure; MP: mechanical power; CV: volume control; PEEP: positive end-expiratory pressure; RR: respiratory rate; PPlateau: plateau pressure; PPeak: peak pressure.

second dashboard in turn focuses specifically on ventilatory support (Fig. 1), exhibiting the ventilation mode, parameters and mechanics. In addition, certain parameters such as tidal volume are displayed in different colors according to whether the values fall within the advisable range or not. This allows us to quickly identify which patients are receiving a tidal volume outside the limits of 6-7 ml/kg, or whether we need to modify parameters in order to secure adequate mechanical power. This dashboard also allows us to quickly identify drug perfusions and doses, as well as which patients are in prone decubitus – facilitating the management and organization of tasks and daily pronation or supination maneuvering of these patients.

The effectiveness of dashboards is conditioned by adequate quality of the CIS data. As reported by Murillo-Cabezas et al., big data techniques can be of help in times of pandemic, though it is essential for the available data to be reliable.⁸ Our group has already shown that in order to achieve this, professional training and effort are crucial.⁹ The professionals must be well aware of the importance of adequate compilation of the data entered in the CIS – both those entered manually and those downloaded automatically from the different point of care devices and originating from other Departments (laboratory test results). It is necessary to check that the devices are properly connected and that the information they afford is reliable and generated on a real-time basis. All this will help to reach our objectives, shortening the identification and response times in situations of increased work burden, and mon-

itoring and maintaining the necessary quality and safety standards.

Financial support

Fondo de Investigación Sanitaria (Instituto de Salud Carlos III, project PI16/00491). Agència de Gestió d'Ajuts Universitaris i de Recerca 2017 SGR 00127. European Regional Development Fund.

References

1. Ferrer R. Pandemia por COVID-19: el mayor reto de la historia del intensivismo. *Med Intensiva*. 2020;44:323–4.
2. Rascado Sedes P, Ballesteros Sanz MÁ, Bodí Saera MA, Carrasco Rodríguez-Rey LF, Castellanos Ortega A, Catalán González METAL. Plan de contingencia para los servicios de medicina intensiva frente a la pandemia COVID-19. *Med Intensiva*. 2020;44:363–70 https://www.semicyuc.org/covid_19files/Plan_de_Contingencia_COVID-19.pdf
3. Wilbanks BA, Langford PA. A review of dashboards for data analytics in nursing. *CIN - Comput Informatics Nurs*. 2014;32:545–9.
4. Berwick DM. Choices for the “new Normal”. *JAMA - J Am Med Assoc*. 2020;323:2125–6.
5. Bodí M, Blanch LL, Maspons R. Los sistemas de información clínica: una oportunidad para medir valor, investigar e innovar a partir del mundo real. *Med Intensiva*. 2017;41:316–8.
6. Ballesteros Sanz MÁ, Hernández-Tejedor A, Estella Á, Jiménez Rivera JJ, González de Molina Ortiz FJ, Sandiumenge Camps

- A, et al. Recomendaciones de «hacer» y «no hacer» en el tratamiento de los pacientes críticos ante la pandemia por coronavirus causante de COVID-19 de los Grupos de Trabajo de la Sociedad Española de Medicina Intensiva, Crítica y Unidades Coronarias (SEMICYUC). *Med Intensiva*. 2020;44:371–88.
7. Khairat SS, Dukkipati A, Lauria HA, Bice T, Travers D, Carson SS. The impact of visualization dashboards on quality of care and clinician satisfaction: integrative literature review. *JMIR Hum Factors*. 2018;5:e22.
 8. Murillo-Cabezas F, Vigil-Martín E, Raimondi N, Pérez-Fernández J. Pandemia de Covid-19 y transformación digital en Cuidados Intensivos. *Med Intensiva*. 2020;1:1–11.
 9. Sirgo G, Esteban F, Gómez J, Moreno G, Rodríguez A, Blanch L, et al. Validation of the ICU-DaMa tool for automatically extracting variables for minimum dataset and quality indicators: The importance of data quality assessment. *Int J Med Inform*. 2018;112(September 2017):166–72. Available from: <https://doi.org/10.1016/j.ijmedinf.2018.02.007>

Laura Claverias^{a,b}, Josep Gómez^{a,b}, Alejandro Rodríguez^{a,b,c}, Jordi Albiol^a, Federico Esteban^{a,b}, María Bodí^{a,b,c,*}

^a Unidad de Cuidados Intensivos, Hospital Universitario Joan XXIII, Tarragona, Spain

^b Instituto de Investigación Sanitaria Pere Virgili, Rovira i Virgili University, Tarragona, Spain

^c Centro de Investigación Biomédica en Red de Enfermedades Respiratorias (CIBERES), Instituto de Salud Carlos III, Madrid, Spain

Corresponding author.

E-mail address: mbodi.hj23.ics@gencat.cat (M. Bodí).

<https://doi.org/10.1016/j.medine.2020.10.003>

2173-5727/

© 2020 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.