

J.L. Serrano-Martínez*, J.F. Machado-Casas,
M. Redondo-Orts, F. Manzano-Manzano,
J. Castaño-Pérez, J.M. Pérez-Villares
*Servicio de Medicina Intensiva, Hospital Universitario
Virgen de las Nieves, Granada, Spain*

Corresponding author.

E-mail address: jlserranouci@gmail.com

(J.L. Serrano-Martínez).

2173-5727/ © 2020 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.

Novel coronavirus (2019-nCov): do you have enough intensive care units?

Nuevo coronavirus (2019-nCov): ¿tiene suficientes unidades de cuidados intensivos?

Recently, the World Health Organization declared the novel coronavirus (2019-nCov) a global health emergency due to its global implications for the health care system and its economic impact. Italy was one of the first European countries with registered clustered cases of acute pneumonia. On February 23, 2020, the Italian government declared the first set of quarantine measures to slow the spread of the virus.¹ Estimations show that 2019-nCoV is a high-diffusion virus with a 2% fatality rate; approximately 20% all hospital admissions were directly to the ICU.² National health care systems could collapse if this spread of pneumonia continues at the current rate. This study aims to analyze official Italian data to build a predictive model.³ From February 23, 2020 to March 15, 2020, daily data from the cumulative reports of the *Protezione Civile Italiana* (Italian Civil Protection) were collected, including the number of positive subjects, hospital admissions, ICU admissions, deaths, and full recovery. Statistical programs were used for the analysis. Different models were tested, and forecast values were calculated, and the best model, with a p -value <0.05 , was considered to calculate the predicted values. The number of positive subjects (PS) follows a non-linear regression with $p < 0.001$ for the number of PS and hospital admissions, PS and ICU admissions, PS and deaths, and PS and recovered subjects. Simultaneously, the number of people admitted to hospitals follows a non-linear regression with $p < 0.001$ (Table 1). Among the 46.7% of PS admitted in hospitals, 10.0% were admitted to the ICU. The ratio of hospitalized patients to those admitted to the ICU is 22.3%, the death rate is 5%, and

the recovery rate is 8%. The relationship between hospital admissions and ICU admissions follows a linear regression, with $p < 0.001$. Recent data on 2019-n CoV present different non-linear growth patterns, besides the rapidly increasing number of PS, which are very susceptible to public health rules. It is fascinating to observe the constant ratio of hospitalized and ICU admissions. If, in the next few weeks, infections reach 1% of the Italian population, over 60.000 ICU beds will be required, which may be the breaking point for the system. These results could be confirmed and highlighted by the increasing trend of ICU admissions, and the relationship between hospitalized patients and ICU admitted subjects. The national health care system needs more time to adapt to and deal with this challenge. The 2019 n-CoV transmission probability presents the following relationship. $y = ax^3bx^2 + cx + d$. Here, y indicates infected subjects, x is the intrinsic potential reproducing number, and the constants a , b , c and d are the intercepts. With environmental strategies and adequate medical treatments, infection and death rates reduced, while recovery rates increased (Fig. 1a and b). Observing Italy's data, this equation is applicable to hospital and ICU admission, and to the rate of death and recovery. As in China, quarantine and environmental strategies have a positive, but slow effect. They can reduce the rate of infection, admissions to ICU, and death, and can change the model.⁴ Furthermore, this is a preliminary interpretation, and not the end of this *phenomenon*. It will be possible to analyze, customize, and fit the best model.⁵ However, in this context, it is important not to forget the emergency; necessary medical and surgical procedures should be guaranteed. A possible solution is to try to re-organize the mission of the hospital as happened in different and less dramatic events.⁶ This model has the potential to predict the worst-case scenario. With this knowledge, we are ready to do the best to prevent the system from reaching the breaking point and to change the 2019 nCoV curve now!

Table 1 Data from February 23 until March 15, 2020. Left panel: number of 2019-nCov positive subjects, number of hospital admissions and ICU admissions, number of deaths, and recovery. Right panel: predicted mean value calculated using non-linear regression. R^2 describes the goodness of fit, where 1 indicates a complete fit.

Day	Number of patients					Predicted number				
	2019-nCov positive	Hospital admission	ICU Admission	Death	Recovery	2019-nCov positive	Hospital admission	ICU Admission	Death	Recovery
1	132	54	26	5	0	39	51	28	0	17
2	229	101	27	6	1	224	89	27	8	0
3	322	114	35	10	1	386	125	31	12	-7
4	400	128	36	12	3	541	169	39	15	-4
5	650	248	56	12	45	705	228	52	17	9
6	888	345	64	21	46	892	312	72	20	32
7	1049	401	105	29	50	1118	427	99	25	64
8	1694	639	140	34	83	1399	583	133	35	106
9	1835	742	166	52	149	1750	787	177	50	159
10	2263	1034	229	79	160	2187	1048	229	73	220
11	2706	1346	295	107	276	2725	1374	292	104	292
12	3296	1790	351	148	414	3380	1774	366	145	373
13	3916	2394	462	197	523	4168	2255	452	198	464
14	5061	2651	567	233	589	5103	2826	550	264	564
15	6378	3557	650	366	622	6202	3496	662	345	674
16	7985	4316	733	463	733	7.495	4.151	757	474	773
17	8514	5038	877	631	1004	9.026	4.878	881	627	940
18	10590	5838	1028	827	1045	10789	5.675	1017	811	1134
19	12839	6650	1153	1016	1258	12802	6.543	1165	1031	1357
20	14995	7246	1328	1266	1439	15084	7.484	1326	1288	1610
21	17750	8372	1518	1441	1966	17653	8.500	1498	1586	1897
22	20603	9663	1672	2063	2335	20528	9.593	1684	1929	2218
R^2						0.999	0.998	0.999	0.993	0.988
p						0.001	0.001	0.001	0.001	0.001




Figure 1 (a) Reports the model curve and its fitting curve of positive subjects of 2019-nCov, hospital and ICU admissions, death, and recovery. (b) Reports the ICU admission trend with a moving averages relation.

Funding

None declared.

The Study follows Strobe Guidelines.

Database: data are available on <http://www.protezionecivile.gov.it/attivita-rischi/rischio-sanitario/emergenze/coronavirus>.

Statistic calculate were performed using Microsoft Excel® and STATA 16® program (STATA Corp LP 4905 Lakeway Drive TX 77845 USA by a physician (with statistic competence) and by an engineer: all calculate were attached as supplementary file.

Author contributions

Melegari G and Barbieri A: concept design of the study, statistic calculate, Maini G: statistic calculate and control, Giuliani E: writing of the paper, Barbieri L: graphical aspects, Baffoni P and Bertellini E: manuscript revision and final approval.

Keypoints

The study analyzes the Italian Novel Coronavirus (2019-nCov) outbreaks, searching possible predicting model, underlining the risk of the trend of phenomena.

Conflict of interest

None.

Acknowledgments

Special thanks to Dr. Giuliano Carrozzi, Epidemiology Azienda AUSL Modena, Italy.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.medin.2020.04.007](https://doi.org/10.1016/j.medin.2020.04.007)

References

- Spina S, Marrazzo F, Migliari M, Stucchi R, Sforza A, Fumagalli R. The response of Milan's Emergency Medical System to the COVID-19 outbreak in Italy. *Lancet*. 2020;395, [http://dx.doi.org/10.1016/S0140-6736\(20\)30493-1](http://dx.doi.org/10.1016/S0140-6736(20)30493-1). PE49-E50 [published online ahead of print, 2020 Feb 28].
- Wax RS, Christian MD. Practical recommendations for critical care and anesthesiology teams caring for novel coronavirus (2019-nCov) patients. *Can J Anesth*. [published online ahead of print, 2020 Feb 12], <http://dx.doi.org/10.1007/s12630-020-01591-x>.
- <http://www.protezionecivile.gov.it/attivita-rischi/rischio-sanitario/emergenze/coronavirus> [accessed 23.03.20].
- Roosa K, Lee Y, Luo R, Kirpich A, Rothenberg R, Hyman JM, et al. Real-time forecasts of the COVID-19 epidemic in China from February 5th to February 24th, 2020. *Infect Dis Model*. 2020;5:256–63, <http://dx.doi.org/10.1016/j.idm.2020.02.002>. Published 2020 Feb 14.
- Chowell G. Fitting dynamic models to epidemic outbreaks with quantified uncertainty: a primer for parameter uncertainty, identifiability, and forecasts. *Infect Dis Model*. 2017;2:379–98.
- Barbieri A, Melegari G, Lob V, Mazzali L. Response by twin Italian hub hospitals in a double seismic event: a retrospective observational investigation. *Prehosp Emerg Care*. 2018;22:353–60.

G. Melegari^{a,*}, E. Giuliani^b, G. Maini^b, L. Barbieri^b, P. Baffoni^a, E. Bertellini^a, A. Barbieri^c

^a Department of Anaesthesia and Intensive Care, Azienda Ospedaliero-Universitaria di Modena, Italy

^b University of Modena and Reggio Emilia, Italy

^c School of Anaesthesia and Intensive Care of University of Modena and Reggio Emilia, Italy

*Corresponding author.

E-mail address: melegari.gabriele@gmail.com (G. Melegari).

0210-5691/ © 2020 Elsevier España, S.L.U. y SEMICYUC. All rights reserved.