



IMAGES IN INTENSIVE MEDICINE

Should abdominal ultrasound be the first image choice test in necrotizing enterocolitis?☆



¿Debería ser la ecografía abdominal la primera prueba de elección de imagen en la enterocolitis necrosante?

A. Batista Muñoz^a, O. Otero Vaccarello^a, J. Rodriguez-Fanjul^{b,*}

^a Unidad Intensiva Neonatal, Servicio de Pediatría, Hospital Joan XXIII, Tarragona, Spain

^b Unidad Intensiva Pediátrica, Servicio de Pediatría, Hospital Joan XXIII, Tarragona, Spain

Available online 20 July 2021

Premature 28-week baby weighing 850 g at birth. Red blood cells are transfused due to prematurity anemia at the patient's 22nd day of life. Twelve hours into the transfusion the patient shows abdominal distension and vomiting of gastric content. The blood test performed showed no alterations in the blood gasometry or elevated parameters of infection. The abdominal X-ray performed showed no signs suggestive of enterocolitis (Fig. 1A), but the abdominal ultrasound performed did show lack of peristalsis and intestinal loops with 2 layers of echogenic foci in the walls (white arrows), a sign indicative of the presence of intraluminal air (Fig. 2A), and echogenic foci in the hepatic parenchyma indicative of gas bubbles inside the portal veins (white arrows), a sign suggestive of portal venous gas (Fig. 2B) oriented as a clinical sign of enterocolitis. The patient remains on a diet and antibiotic therapy is started. The ultrasound looks normal with good disease progression



Figure 1

☆ Please cite this article as: Batista Muñoz A, Otero Vaccarello O, Rodriguez-Fanjul J. ¿Debería ser la ecografía abdominal la primera prueba de elección de imagen en la enterocolitis necrosante? Med Intensiva. 2021;45:445–446.

* Corresponding author.

E-mail address: javier.rodriguez.fanjul@gmail.com
(J. Rodriguez-Fanjul).

(white asterisks) (Fig. 2C) after the completion of a 14 day course of antibiotic treatment.




Figure 2