

## medicina intensiva

medicina Internsiva

Internsiv

http://www.medintensiva.org/en/

## IMAGES IN INTENSIVE MEDICINE

## Horner's syndrome after chest drain insertion due to pneumothorax



Síndrome de Horner tras la inserción de un drenaje torácico por neumotórax

Arthur Orieux<sup>a,b,\*</sup>, Raphaël Birot<sup>a</sup>, Didier Gruson<sup>a,c</sup>, Renaud Prevel<sup>a,c</sup>

- a Service de Médecine Intensive Réanimation CHU de Bordeaux, Bordeaux, France
- <sup>b</sup> Unité INSERM UMR1034 Biology of Cardiovascular Diseases Université de Bordeaux, Bordeaux, France
- c Unité INSERM UMR1045 Centre de Recherche Cardio-Thoracique de Bordeaux Université de Bordeaux, Bordeaux, France

Available online 15 June 2024

A 27-year-old man was admitted for a first episode of spontaneous apical right pneumothorax (chest X-ray, Fig. 1A). The patient reported no relevant medical history except cannabis use. We inserted an 8.5 French pigtail catheter in the pleural space (Monaldi position). No adverse events occurred during pleural drainage (local anesthesia). Chest X-ray assessed the correct chest tube position and complete lung re-expansion (Fig. 1B). 24h after chest drain insertion, the patient presented ipsilateral palpebral ptosis with enophthalmos and myosis occurred (Fig. 1C): a

pathognomonic triad defines Horner's syndrome. A brain and cervical magnetic resonance angiography and a chest computed tomography did not reveal any lesion to explain Horner's syndrome. The right pleural drain was removed as soon as pulmonary re-expansion. At hospital discharge, the patient suffered from persistent right myosis and ptosis.

Exceptional Horner's syndrome cases could occur after a central or peripheral sympathetic pathway lesion (stellate ganglion) due to a chest tube.

E-mail address: arthur.orieux@chu-bordeaux.fr (A. Orieux).

DOI of original article: https://doi.org/10.1016/j.medin.2024.05.003

<sup>\*</sup> Corresponding author.




Figure 1