

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.medin.2021.10.005](https://doi.org/10.1016/j.medin.2021.10.005).

## Bibliografía

- Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395:1054–62, [http://dx.doi.org/10.1016/S0140-6736\(20\)30566-3](http://dx.doi.org/10.1016/S0140-6736(20)30566-3).
- Tobin MJ, Jubran A, Laghi F. Misconceptions of pathophysiology of happy hypoxemia and implications for management of COVID-19. *Respir Res*. 2020;21:249, <http://dx.doi.org/10.1186/s12931-020-01520-y>.
- Dhont S, Derom E, Van Braeckel E, Depuydt P, Lambrecht BN. The pathophysiology of 'happy' hypoxemia in COVID-19. *Respir Res*. 2020;21:198, <http://dx.doi.org/10.1186/s12931-020-01462-5>.
- Rice TW, Wheeler AP, Bernard GR, Hayden DL, Schoenfeld DA, Ware LB, et al. Comparison of the SpO<sub>2</sub>/FiO<sub>2</sub> ratio and the PaO<sub>2</sub>/FiO<sub>2</sub> ratio in patients with acute lung injury or ARDS. *Chest*. 2007;132:410–7, <http://dx.doi.org/10.1378/chest.07-0617>.
- Bilan N, Dastranji A, Behbahani AG. Comparison of the SpO<sub>2</sub>/FiO<sub>2</sub> ratio and the PaO<sub>2</sub>/FiO<sub>2</sub> ratio in patients with acute lung injury or acute respiratory distress syndrome. *J Cardiovasc Thorac Res*. 2015;7:28–31, <http://dx.doi.org/10.15171/jcvtr.2014.06>.
- Riviello ED, Kiviri W, Twagirumugabe T, Mueller A, Banner-Goodspeed VM, Officer L, et al. Hospital incidence and outcomes of the acute respiratory distress syndrome using the Kigali modification of the Berlin definition. *Am J Respir Crit Care Med*. 2016;193:52–9, <http://dx.doi.org/10.1164/rccm.201503-0584OC>.
- Tobin MJ, Laghi F, Jubran A. Why COVID-19 silent hypoxemia is baffling to physicians. *Am J Respir Crit Care Med*. 2020;202:356–60, <http://dx.doi.org/10.1164/rccm.202006-2157CP>.
- Fukuda Y, Tanaka A, Homma T, Kaneko K, Uno T, Fujiwara A, et al. Utility of SpO<sub>2</sub>/FiO<sub>2</sub> ratio for acute hypoxemic respiratory failure with bilateral opacities in the ICU. *PLOS ONE*. 2021;16:e0245927, <http://dx.doi.org/10.1371/journal.pone.0245927>.
- Choi KJ, Hong HL, Kim EJ. The association between mortality and the oxygen saturation and fraction of inhaled oxygen in patients requiring oxygen therapy due to COVID-19-associated pneumonia. *Tuberc Respir Dis (Seoul)*. 2021;84:125–33, <http://dx.doi.org/10.4046/trd.2020.0126>.

A. Kumar<sup>a</sup>, R. Aggarwal<sup>b,\*</sup>, P. Khanna<sup>a</sup>, R. Kumar<sup>a</sup>, A.K. Singh<sup>a</sup>, K.D. Soni<sup>b</sup>, A. Trikha<sup>a</sup>

<sup>a</sup> *Department of Anaesthesiology, Critical Care and Pain Medicine, AIIMS, New Delhi, India*

<sup>b</sup> *Critical and Intensive Care, JPNATC, AIIMS, India*

\* Corresponding author.

E-mail address: [pathakricha@yahoo.co.in](mailto:pathakricha@yahoo.co.in) (R. Aggarwal).

<https://doi.org/10.1016/j.medin.2021.10.005>

0210-5691/ © 2021 Elsevier España, S.L.U. y SEMICYUC. All rights reserved.

## Heterogeneity of hypoxemia severity according to pulse oximetry and blood gas analysis in COVID-19 pneumonia



## Heterogeneidad de la severidad de hipoxemia de acuerdo a oximetría de pulso y gases arteriales en neumonía COVID-19

Dear Editor,

Pneumonia is the hallmark of severe COVID-19.<sup>1</sup> Strain in healthcare systems across the world has forced countless hospitals to conduct grueling triages to decide who gets to be admitted when healthcare saturation was rampant.<sup>2</sup> As these decisions are inherently complex, numerous risk scores and predictor factors have been described to aid the attending medical team.<sup>3–5</sup> These often include clinical and laboratory values.


One commonly utilized criteria to determine patient severity is the severity of hypoxemia.<sup>6</sup> This can be assessed with arterial oxygen pressure (PaO<sub>2</sub>), PaO<sub>2</sub> to inspired fraction of oxygen (FiO<sub>2</sub>) ratio, arterial oxygen saturation (SatO<sub>2</sub>), pulse oximeter oxygen saturation (SpO<sub>2</sub>), SatO<sub>2</sub> to

FiO<sub>2</sub> ratio, SpO<sub>2</sub> to FiO<sub>2</sub> ratio, and the prescribed oxygen device.<sup>7</sup>


The use of these criteria for hypoxemia severity in non-intubated patients has been criticized given the expected high inter-patient variability in FiO<sub>2</sub>, shunt fraction, and physician's choice of oxygenation device and oxygen flow.<sup>7</sup> Therefore, relying on these criteria is suboptimal given the low comparability between different patients.

In this study, we aimed to compare the severity of hypoxemia in patients with severe COVID-19 according to oxygenation index arriving at an emergency department.

We performed a retrospective cohort study collecting information on every patient who arrived at the emergency department (ED) of a reference COVID-19 tertiary center between April 1st, 2020, and April 30th, 2021. At arrival, every patient had to go through a triage station where vital signs (including SpO<sub>2</sub>) were documented before entering the emergency department. Once in the emergency department, all patients who had low SpO<sub>2</sub> (usually < 92%) received supplemental oxygen. Only nasal cannula and non-rebreathing masks were available at our center. Arterial blood gas analysis was performed in all patients with suspected COVID-19. Generally, FiO<sub>2</sub> was estimated heuristically by the treating physician by adding to the baseline FiO<sub>2</sub> (21%) 3% for every extra liter of oxygen per minute.<sup>8</sup> For example, a patient receiving 2 l of minute of supplemental oxygen would have a calculated FiO<sub>2</sub> of 27% (21 + 3\*2).



**Figure 1** Correlation between oxygenation indexes. SpO<sub>2</sub>: oxygen saturation with pulse oximeter; PaO<sub>2</sub>: oxygen pressure in arterial blood; SatO<sub>2</sub>: oxygen saturation in arterial blood; FiO<sub>2</sub>: fraction of inspired oxygen.



**Figure 2** Variability of concordance between oxygenation indexes’ quintiles. SpO<sub>2</sub>: oxygen saturation with pulse oximeter; PaO<sub>2</sub>: oxygen pressure in arterial blood; SatO<sub>2</sub>: oxygen saturation in arterial blood; FiO<sub>2</sub>: fraction of inspired oxygen. (A) Shows relation between PaO<sub>2</sub>/FiO<sub>2</sub> and SatO<sub>2</sub>/FiO<sub>2</sub>, (B) shows relation between PaO<sub>2</sub>/FiO<sub>2</sub> and SpO<sub>2</sub>/FiO<sub>2</sub>, and (C) shows relation between SpO<sub>2</sub>/FiO<sub>2</sub> and SatO<sub>2</sub>/FiO<sub>2</sub>. Numbers represent the respective quintile (“PaO<sub>2</sub>/FiO<sub>2</sub> 1” represents the first quintile of said variable).

SpO<sub>2</sub> was obtained at ambient air, while arterial blood gas was obtained almost universally when patients received supplemental oxygen. Given the closeness between SpO<sub>2</sub> and blood gas analysis, we believe it is reasonable to assume that the clinical status of the patient is comparable between these two circumstances. We obtained data about PaO<sub>2</sub>, SatO<sub>2</sub>, SpO<sub>2</sub>, and FiO<sub>2</sub>, with which we calculated PaO<sub>2</sub>/FiO<sub>2</sub>, SatO<sub>2</sub>/FiO<sub>2</sub>, and SpO<sub>2</sub>/FiO<sub>2</sub>. All SpO<sub>2</sub> were taken at triage with a Nihon MU-631RK portable monitor, and as such, were ambient-air (FiO<sub>2</sub> 21%). SpO<sub>2</sub>/FiO<sub>2</sub> was included to better compare oxygenation indexes. Blood gas analysis was performed in the emergency department with the RADIOMETER ABL90 flex analyzer, which directly measured SatO<sub>2</sub> and PaO<sub>2</sub>. Oxygen device used at the time of the arterial blood gas analysis could not be confidently determined, so it was not included. Since the low reliability of FiO<sub>2</sub> is the most criticized aspect of using the oxygenation device as a marker of hypoxia severity, it does not affect our analysis.

We categorized a patient’s hypoxemia severity by quintiles of SpO<sub>2</sub>/FiO<sub>2</sub>, PaO<sub>2</sub>/FiO<sub>2</sub>, and SatO<sub>2</sub>/FiO<sub>2</sub> (from now on referred to as “oxygenation indexes”). Lower values indicate a higher hypoxemia severity. We determined strength of concordance between oxygenation index quintiles with chord diagrams across all three oxygenation index pairs. We calculated Spearman correlation coefficients for the three possible pairs of oxygenation indexes. We built scatterplots

and used locally weighted scatterplot smoothing regression to graphically represent the data.

All analyses were conducted with R software version 4.0.0. The study was approved by the Research in Humans ethics committee of the authors’ institution.

A total of 23,049 triage visits occurred during the study period, corresponding to 19,644 individual patients. Of these, 8123 were due to suspected COVID-19. Among 4663 patients that were finally admitted to the ED, arterial blood gas analysis was obtained for 2960 patients.

Median FiO<sub>2</sub> was 0.34 (inter-quartile range [IQR] 0.25–0.60), PaO<sub>2</sub> 68 mmHg (57–84), SatO<sub>2</sub> 94% (91–97), SpO<sub>2</sub> 83% (73–88), PaO<sub>2</sub>/FiO<sub>2</sub> 211 (124–281), SatO<sub>2</sub>/FiO<sub>2</sub> 274 (160–364), and SpO<sub>2</sub>/FiO<sub>2</sub> 395 (348–419). Correlation among oxygenation indexes is shown in Fig. 1. A strong correlation was seen between PaO<sub>2</sub>/FiO<sub>2</sub> & SpO<sub>2</sub>/FiO<sub>2</sub> (rho=0.6, p<0.001), and SatO<sub>2</sub>/FiO<sub>2</sub> & SpO<sub>2</sub>/FiO<sub>2</sub> (rho=0.65, p<0.001), while a very strong correlation was seen between PaO<sub>2</sub>/FiO<sub>2</sub> & SatO<sub>2</sub>/FiO<sub>2</sub> (rho=0.88, p<0.001). None of the variable pairs showed a linear relationship. All oxygenation indexes showed a considerable cross-over among quintiles (Fig. 2), with only 785 (26.5%) patients being in the same severity quintile across all indexes.

Our study shows there is considerable heterogeneity when classifying hypoxemia severity with different oxygenation indexes. While good correlation was observed among

the three evaluated pairs, the lower correlation between  $SpO_2/FiO_2$  and the others could be due to inaccuracy of pulse oximeters when used in patients with low oxygen saturation, physician imprecision when estimating  $FiO_2$ , and/or mixed blood in the gas analysis.<sup>9</sup> It is likely multiple factors are involved in most cases, which is consistent with the previously mentioned concerns.<sup>7</sup> For example, if solely based on  $SpO_2$  at triage, a patient could be wrongly classified as having a more severe disease, giving preference to a patient in better condition. Also, follow up could be hard if it is done only with oxygenation parameters. The morning medical team could classify the patient with a given severity with blood gas analysis, while the evening group could use a pulse oximeter and determine intubation is indicated.

Degree of hypoxemia is not a good measure when considered by itself in non-intubated patients with COVID-19 pneumonia, as considerable variation occurs depending on whether pulse oximetry or blood gas analysis is utilized.

### Data availability

Code used for the analysis is freely available at [https://github.com/isaac-nunez/Oxygenation\\_indexes](https://github.com/isaac-nunez/Oxygenation_indexes). Data will be made available upon reasonable request to the corresponding author.

### Sources of funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### Conflicts of interest

The authors have no conflicts of interest.

### Acknowledgements

The authors would like to thank the triage personnel of their institution for their invaluable work during the pandemic, as well as an anonymous reviewer for helpful comments to improve the manuscript.

### Bibliografía

- Berlin DA, Gulick RM, Martinez FJ. Severe Covid-19. *N Engl J Med*. 2020;383:2451–60.
- The Lancet. Humanitarian crises in a global pandemic. *Lancet* 2020;396:447.
- Soto-Mota A, Marfil-Garza BA, Martínez Rodríguez E, Barreto Rodríguez JO, López Romo AE, Alberti Minutti P, et al. The low-harm score for predicting mortality in patients diagnosed with COVID-19: a multicentric validation study. *J Am Coll Emerg Physicians Open*. 2020;1:1436–43.
- Núñez I, Priego-Ranero AA, García-González HB, Jiménez-Franco B, Bonilla-Hernández R, Domínguez-Cherit G, et al. Common hematological values predict unfavorable outcomes in hospitalized COVID-19 patients. *Clin Immunol*. 2021;225:108682.
- Wynants L, Van Calster B, Collins GS, Riley RD, Heinze G, Schuit E, et al. Prediction models for diagnosis and prognosis of covid-19: systematic review and critical appraisal. *BMJ*. 2020:m1328.
- Tobin MJ, Laghi F, Jubran A. Why COVID-19 silent hypoxemia is baffling to physicians. *Am J Respir Crit Care Med*. 2020;202:356–60.
- Tobin MJ, Jubran A, Laghi F.  $PaO_2/FiO_2$  ratio: the mismeasure of oxygenation in COVID-19. *Eur Respir J*. 2021;57:2100274.
- Cao Z, Luo Z, Hou A, Nie Q, Xie B, An X, et al. Volume-targeted versus pressure-limited noninvasive ventilation in subjects with acute hypercapnic respiratory failure: a multicenter randomized controlled trial. *Respir Care*. 2016;61:1440–50.
- Chan ED, Chan MM, Chan MM. Pulse oximetry: understanding its basic principles facilitates appreciation of its limitations. *Respir Med*. 2013;107:789–99.

I. Núñez<sup>a,\*</sup>, A. Soto-Mota<sup>b</sup>

<sup>a</sup> *Departamento de Medicina Interna, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Ciudad de México, Mexico*

<sup>b</sup> *Unidad de Investigación de Enfermedades Metabólicas, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Ciudad de México, Mexico*

\* Corresponding author.

E-mail address: [isaac.nunezs@incmnsz.mx](mailto:isaac.nunezs@incmnsz.mx) (I. Núñez).

<https://doi.org/10.1016/j.medin.2021.10.004>

0210-5691/ © 2021 Elsevier España, S.L.U. y SEMICYUC. All rights reserved.