



SCIENTIFIC LETTERS

Implementation of educational programs to prevent drowning. What can be done in nursery school? ☆



Implantación de programas educativos para prevenir ahogamientos. ¿Qué se puede hacer desde la escuela infantil? ☆

Dear Sir,

Drowning is regarded as a “global killer” that has not received the attention it requires.¹ In Spain, drowning is responsible for 13% of global childhood mortality due to injuries and is the second most frequent cause of accidental death, after traffic accidents.²

Drowning is preventable, and recommendations of demonstrated efficacy have been established, such as the supervision of children, the presence of lifeguards, adequate information and signaling, the protection of aquatic spaces and the use of adequate flotation devices such as life vests.^{1–4} In countries with a strong public safeguarding tradition such as Australia, educational programs have been shown to offer positive results in terms of learning and prevention.⁵ However, to the best of our knowledge, no studies in Spain have analyzed the most appropriate contents and methodology for introducing the prevention of drowning in school education. We thus sought to evaluate

a pilot childhood education program focused on the understanding, learning and memorization of measures that can prevent drowning.

A sample of 26 preschool children (5 years of age) from Pontevedra (Spain) were included in the study. All of them routinely visited beaches and swimming pools, but none had received any training or instructions referred to drowning.

A quasi-experimental study design was used. The participants were divided into two cohorts: a control group (CG; $n = 12$) that received no training, and an experimental group (EG; $n = 14$) enrolled in the drowning prevention program. The study data were anonymized and written informed consent was obtained from all the parents.

The educational material was prepared by a multidisciplinary team of physicians, childhood educators and lifeguards. An illustrated story for children based on an imaginary 5-year-old character was chosen, entitled “Xoana goes to the swimming pool and Xoana goes to the beach” (“*Xoana va a la piscina y Xoana va a la playa*”). The training program lasted one week.

Two forms were designed for the evaluation of learning: one for each scenario (beach and swimming pool) (Fig. 1). Each contained safety elements (life vest, green flag, lifeguard, supervising adult, safety barrier) and potentially hazardous elements (diving, red flag, yellow flag, eating before swimming or using a float), in compliance with the corresponding scientific recommendations.^{1,2} Both



Figure 1 Evaluation form at T2 corresponding to the experimental group on the beach (prevention stickers added) and in the swimming pool (rounded safety elements).

☆ Please cite this article as: Barcala-Furelos R, Carbia-Rodríguez P, Peixoto-Pino L, Abelairas-Gómez C, Rodríguez-Núñez A. Implantación de programas educativos para prevenir ahogamientos. ¿Qué se puede hacer desde la escuela infantil? Med Intensiva. 2019;43:180–182.

Table 1 Results referred to the evaluation of knowledge about the prevention of drowning in the different study phases.

		<i>n</i> = 26 CG, <i>n</i> = 12; (EG, <i>n</i> = 14)								
Swimming pool-T0		Swimming pool-T1		Swimming pool-T2		ANOVA (Friedman test)				
Mean (SD)	95%CI	Mean (SD)	95%CI	Mean (SD)	95%CI	a vs. b	a vs. c	b vs. c		
CG	27.14 (16.84)	17.42–36.86	40.00 (29.71)	27.49–61.80	48.57 (20.33)	36.83–60.31	0.02	0.003	0.10	
EG	6.67 (27.41)	–24.08–10.75	96.67 (11.55)	89.33–104.00	81.67 (19.92)	69.01–94.33	0.001	0.001	0.03	
Sig.	0.03 ^a		<0.001 ^b		0.001 ^b					
Beach-T0		Beach-T1		Beach-T2						
Mean (SD)	95%CI	Mean (SD)	95%CI	Mean (SD)	95%CI	a vs. b	a vs. c	b vs. c		
CG	37.50 (23.51)	23.92–51.08	44.64 (29.71)	27.49–61.80	48.21 (20.72)	36.25–60.18	0.26	0.03	1.00	
EG	25.00 (15.08)	15.42–34.58	97.92 (7.22)	93.33–102.50	97.92 (7.22)	93.33–102.50	0.001	0.001	1.00	
Sig.	0.12 ^a		<0.001 ^b		<0.001 ^b					
GAP-T0		GAP-T1		GAP-T2						
Mean (SD)	95%CI	Mean (SD)	95%CI	Mean (SD)	95%CI	a vs. b	a vs. c	b vs. c		
CG	32.32 (16.63)	22.72–41.92	42.32 (22.39)	29.39–55.25	48.39 (17.75)	38.15–58.64	0.02	0.004	0.13	
EG	15.83 (17.03)	5.01–26.66	97.29 (6.52)	93.15–101.44	89.79 (10.68)	83.00–96.58	0.001	0.001	0.09	
Sig.	0.02 ^a		<0.001 ^b		<0.001 ^b					

ANOVA: analysis of variance; SD: standard deviation; CG: control group; EG: experimental group; 95%CI: 95% confidence interval; Sig.: statistical significance; T0: test 0 pre-training; T1: test 1 post-training; T2: test 2 one month post-training; GAP: global assessment of aquatic prevention.

^a Student *t*-test
^b Mann-Whitney *U*-test

between- (CG versus EG) and within-group analyses (CG versus CG and EG versus EG) were made at three timepoints: before training (T0), at the end of training (T1), and after one month as a memory check (T2).

The dependent variable was the result (%) referred to the right answers minus the wrong answers in each test. The following equations were used: ($\Sigma\%$ right answers – $\Sigma\%$ wrong answers = % knowledge of prevention in swimming pool) and ($\Sigma\%$ right answers – $\Sigma\%$ wrong answers = % knowledge of prevention on beach), while for the global assessment of aquatic prevention we used: ($[\%$ knowledge of prevention in swimming pool + % knowledge of prevention on beach]/2). The SPSS version 20 statistical package for Mac (IBM Corp., Armonk, NY, USA) was used for the statistical analysis. The following statistical tests were used: (1) normal data distribution was checked with the Shapiro-Wilk test; (2) within-group analyses were based on the Mann-Whitney *U*-test; and (3) between-group comparisons were made using the Student *t*-test for independent samples or the Friedman test in the absence of a normal distribution. Statistical significance was considered for $p < 0.05$ in all cases.

The level of knowledge of these young children was seen to be low at T0 referred to both the swimming pool and the beach (percentage right answers between 6 and 37%).

After training, the EG improved significantly at T1 versus T0 both in the swimming pool and on the beach ($p < 0.001$). In relation to global aquatic prevention, the EG exceeded the CG by 55% (97.29 ± 7.22 versus 42.32 ± 22.39 , respectively; $p < 0.001$). After one month (T2), the EG continued to

yield better results in both the swimming pool ($p = 0.001$) and on the beach ($p < 0.001$), and in relation to global aquatic prevention, the EG exceeded the CG by 41% (89.79 ± 10.68 versus 48.39 ± 17.15 , respectively; $p < 0.001$). In turn, the within-group analysis showed clear improvement between T0 and T1, with learning stabilization at T2 (Table 1).

Different studies indicate that the introduction of drowning prevention strategies in the educational system has positive effects in terms of recognizing drowning triggering factors at vulnerable ages.^{5,6} While preliminary, our data show how a simple but well oriented intervention is able to ensure learning and retaining of what has been learned over the middle term in children as young as 5 years of age. The data obtained should serve to support the design and implementation of broad and systematic school programs for reducing the incidence of drowning in our setting.

References

1. World Health Organization. Preventing drowning: an implementation guide. Geneva: WHO; 2017. Available from: <http://apps.who.int/iris/bitstream/10665/255196/1/9789241511933-eng.pdf?ua=1> [accessed 29.05.17].
2. Rubio B, Yagüe F, Benítez MT, Esparza MJ, González JC, Sánchez F, et al. Recomendaciones sobre la prevención de ahogamientos. *An Pediatr (Barc)*. 2015;82, <http://dx.doi.org/10.1016/j.anpedi.2014.06.010>, 43.e1–43.e5.
3. Szpilman D, Webber J, Quan L, Bierens J, Morizot-Leite L, Langendorfer SJ, et al. Creating a drowning chain of survival.

Resuscitation. 2014;85:1149–52, <http://dx.doi.org/10.1016/j.resuscitation.2014.05.034>.

4. Szpilman D, Tipton M, Sempritt J, Webber J, Bierens J, Dawes P, et al. Drowning timeline: a new systematic model of the drowning process. *Am J Emerg Med*. 2016;34:2224–6, <http://dx.doi.org/10.1016/j.ajem.2016.07.063>.
5. Wilks J, Kanasa H, Pendergast D, Clark K. Beach safety education for primary school children. *Int J Inj Contr Saf Promot*. 2017;24:283–92, <http://dx.doi.org/10.1080/17457300.2016.1170043>.
6. Turgut T, Yaman M, Turgut A. Educating children on water safety for drowning prevention. *Soc Indic Res*. 2016;129:787–801.

R. Barcala-Furelos^{a,b,*}, P. Carbia-Rodríguez^a,
L. Peixoto-Pino^c, C. Abelairas-Gómez^{b,c},
A. Rodríguez-Núñez^{b,d,e}

^a Grupo de Investigación REMOSS, Facultad de Ciencias de la Educación y el Deporte, Universidad de Vigo, Vigo, Pontevedra, Spain

^b Grupo de Investigación CLINURSID, Departamento de Enfermería, Universidade de Santiago de Compostela, A Coruña, Spain

^c Facultad de Ciencias de la Educación, Universidad de Santiago de Compostela, Santiago de Compostela, A Coruña, Spain

^d Área de Pediatría, Servicio de Críticos y Urgencias Pediátricas, Hospital Clínico Universitario de Santiago, Instituto de Investigación Sanitaria de Santiago de Compostela (IDIS), Santiago de Compostela, A Coruña, Spain

^e Departamento de Enfermería, Universidade de Santiago, Santiago de Compostela, A Coruña, Spain

* Corresponding author.

E-mail address: roberto.barcala@uvigo.es

(R. Barcala-Furelos).

2173-5727/

© 2017 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.

Susac's syndrome. A case of autoimmune encephalitis[☆]



Síndrome de Susac. A propósito de las encefalitis autoinmunes

Dear Sir,

Encephalitis is a potentially fatal inflammatory disorder of the central nervous system (CNS). Although the diagnosis is complex, early identification of the disorder is important, since it can condition the prognosis. In our setting, most cases of encephalitis are of viral origin, though a number of autoimmune presentations may also be observed¹ that should be taken into account in order to start early treatment.

We present the case of a 40-year-old male with no disease history of relevance who reported to the emergency room repeatedly because of 15–20 days of varying neurological manifestations in the form of headache, instability or confusion starting 24 h after a disabling acute headache episode. The brain CAT study revealed no alterations. In view of the variability of the clinical manifestations and the personal conflicts of the patient, the condition was taken to represent a dissociative disorder, and neuroleptic and antidepressive treatment was started. However, the patient continued to suffer disconnection from the environment and disorientation, heteroaggressive behavior, and oppositionist and behavioral disorders. The patient was admitted to the psychiatric ward with akinetic mutism. The neurological exploration revealed no alterations (though the patient

was totally uncooperative), and the laboratory test results (including serological testing for syphilis and HIV infection) proved normal. On the third day his condition worsened, with no response to stimuli, the appearance of flaccid tetraparesis and a left positive Babinski reflex.

On day four the patient developed high fever, respiratory sounds suggestive of bronchoaspiration, and finally tonic-clonic seizures. Admission to our Intensive Care Unit (ICU) was thus decided. The patient was admitted under postcritical condition, with fever (38.6 °C) and SatO₂ 75% with oxygen therapy (reservoir); intubation was decided, and mechanical ventilation was started. There were few crepitations in the right lung base, and the rest of the exploration yielded no findings of relevance. The routine laboratory test parameters were normal, with the exception of CPK 1800 U/l, and the chest X-rays showed a weak right basal infiltrate. Repeat CAT showed no alterations, and lumbar puncture yielded clear cerebrospinal fluid (CSF), emerging without pressure, containing 13 cells (with 92% monocytes), no glucose depletion, and a protein concentration of 189.7 mg/dl. Aciclovir and broad spectrum antibiotic treatment were prescribed.

The hemodynamic and respiratory alterations quickly reverted; the blood and CSF cultures proved negative; and the usual flora was identified in the tracheal aspirate. Magnetic resonance imaging (MRI) revealed numerous foci with enhanced signal intensity in T2-FLAIR images. The lesions were of small size, tended to merge, and presented a cotton-like appearance; they were mostly restricted in size in diffusion sequencing and did not increase in intensity with contrast administration. The lesions were located bilaterally in the basal ganglia, corpus callosum (body and splenium), the medial regions of both thalami, the deep periventricular white matter, the frontal subcortical zone, the cortico-subcortical region at right parasylvian level and the left anteromedial temporal region – with involvement of the cerebellar peduncles and right hemi-mesencephalon (Fig. 1).

The epidemiological context required us to discard encephalitis caused by west Nile virus (the patient worked

[☆] Please cite this article as: Mora López D, Tristancho Garzon A, Guzmán Llorente M, Jiménez Conde C, Montero Urbina A, Oliva Fernández P. Síndrome de Susac. A propósito de las encefalitis autoinmunes. *Med Intensiva*. 2019;43:182–184.