



ORIGINAL ARTICLE

Intubation in the Intensive Care Unit: C-MAC video laryngoscope versus Macintosh laryngoscope



S. Dey^a, D. Pradhan^{b,*}, P. Saikia^c, P. Bhattacharyya^d, H. Khandelwal^e, K.N. Adarsha^f

^a All India Institute of Medical Sciences, Raipur, India

^b School of Medical Sciences and Research, Sharda University, New Delhi, India

^c Guwahati Medical College, Guwahati, India

^d North Eastern Indira Gandhi Regional Institute of Health and Medical Sciences, Shillong, India

^e Shri Guru Ram Rai Institute of Medical and Health Sciences, Dehradun, India

^f Apollo Hospital, Bannerghatta Road, Bengaluru, India

Received 25 July 2019; accepted 7 October 2019

KEYWORDS

Video laryngoscope;
Intubation;
Visualization of the
glottis

Abstract

Purpose: Various modifications of the Macintosh blade and direct laryngoscopy have been incorporated into practice to improve the intubation success rate and avoid complications while ensuring patient safety. This study evaluates the usefulness of two different direct laryngoscopy methods used by operators with various level of experience in the Intensive Care Unit.

Material and methods: In a single centre prospective study, C-MAC and Macintosh laryngoscopes were compared in terms of laryngoscopy and intubation outcomes such as glottic visualization, number of intubation attempts, intubation success and satisfaction score.

Results: During the one-year study period, 263 patients were evaluated and data of 218 patients were analyzed. The rate of successful first attempt intubation was higher in the video laryngoscope group (VL) (84% vs 57%; $P < 0.001$). A significantly greater number of patients in the Macintosh laryngoscopy group had difficult visualization of the glottis in terms of the modified Cormack and Lehane classification and Percentage of Glottic Opening scale.

Conclusion: The use of video laryngoscope for intubation in ICU settings results in better visualization of the glottis and a higher incidence of successful intubation attempts.

© 2019 Elsevier España, S.L.U. y SEMICYUC. All rights reserved.

* Corresponding author.

E-mail address: dmkcg85@gmail.com (D. Pradhan).

PALABRAS CLAVE

Videolaringoscopia;
Intubación;
Visualización de la
glotis

Intubación en la unidad de cuidados intensivos: videolaringoscopia C-MAC frente a laringoscopia Macintosh

Resumen

Objetivo: En la práctica clínica se han incorporado diversas modificaciones a la hoja de Macintosh y del laringoscopia directa para mejorar la tasa de éxito de la intubación y evitar complicaciones, a la vez que se garantiza la seguridad del paciente. En este estudio evaluamos la utilidad de los 2 métodos de laringoscopia directa empleados por profesionales sanitarios con distintos grados de experiencia en la unidad de cuidados intensivos.

Materiales y métodos: En un estudio prospectivo y unicéntrico se compararon los laringoscopios de Macintosh y C-MAC en términos de deslances de la laringoscopia y la intubación, como la visualización de la glotis, el número de intentos de intubación, el éxito de la intubación y la puntuación de la satisfacción.

Resultados: Durante el periodo del estudio de un año se evaluaron 263 pacientes y se analizaron los datos de 218 pacientes. La tasa de intubación con éxito al primer intento fue más elevada en el grupo de videolaringoscopia (84% frente al 57%, $p < 0,001$). Un número significativamente más alto de pacientes del grupo del laringoscopia de Macintosh presentó dificultades de visualización de la glotis de acuerdo con la clasificación modificada de Cormack y Lehane, y según la escala de porcentaje de apertura de la glotis.

Conclusión: El uso del videolaringoscopia para la intubación en el entorno de la unidad de cuidados intensivos se asocia con una mejor visualización de la glotis y una tasa más alta de intentos de intubación exitosos.

© 2019 Elsevier España, S.L.U. y SEMICYUC. Todos los derechos reservados.

Introduction

Intubation and mechanical ventilation are essential and sometime urgent in various life threatening conditions in the Intensive Care Unit (ICU). Laryngoscopy and intubation are highly challenging and not free from complications such as hypoxia, bradycardia, hypotension, damage to upper airways, aspiration and cardiac arrest.¹ Various attempts have been made to improve success rate and minimize complications, which include use of neuromuscular blockade, care bundles, training on simulators.²

In the last decade, many airway equipments including various types of video laryngoscope (VL) were being developed and compared with the conventional Macintosh laryngoscopy (ML). Better visualization of glottic opening may result in the easy and quick endotracheal intubation while reducing complication rates. Recent systematic review found evidence supporting decreased incidences of intubation failure in the context of difficult airway in operation theatre by using video laryngoscopy. There are positive and negative reports regarding the benefits of video laryngoscopy over direct laryngoscopy in critically ill patients in an ICU.¹⁻³ From Indian subcontinent, no studies have been reported about the comparison between VL and ML in ICU settings. This study was conducted to test the hypothesis that C-MAC video laryngoscope increases the success rate of first pass orotracheal intubation in intensive care setting as compared to Macintosh laryngoscope.

Materials and methods

This prospective, randomized, comparative study was conducted in a 32-bed multidisciplinary adult ICU of a

tertiary-care teaching hospital. Following approval from Institute Ethics Committee, the study was conducted over a period of eighteen months (from 1st January 2017 till 30th June 2018). Informed consents were obtained from patients or their legal representatives and data relevant to intubation were collected in a prescribed format.

All ICU patients who needed elective endotracheal intubation were included in the study except patients of age less than 18 years, pregnant and lactating women, patients with facial trauma including burn injury, suspected or confirmed cervical spine injury, lack of time for randomization and inclusion due to ongoing resuscitative efforts, unable to obtain informed consent. Following inclusion into the study, sequential sealed opaque envelope technique was used for allocation of laryngoscopy method, either Macintosh Laryngoscope (ML group), blade size 3 or 4 or Karl Storz C-MAC video laryngoscope (VL group), Macintosh shaped blades size 3 or 4 (Karl Storz GmbH & Co, KG, Tuttlingen, Germany). C-MAC VL blade is a Macintosh type with a small camera and light source at the distal one third of the blade. The removable electronic module, interface between front-end of VL and the portable monitor unit, fits into the receptacle of laryngoscope and after intubation they can be separated for the preparation of the laryngoscope for the next intubation.

All anaesthesiologists working in this ICU were being oriented to video laryngoscopy (using C-MAC VL) on manikin at the time of joining in the department. Intubating anaesthesiologists (laryngoscopist) had experience of minimum fifty videolaryngoscopies using C-MAC VL. Laryngoscopists were categorized (based on exposure) into junior (up to three years), senior (more than 3–8 years) and consultant (more than 8 years) based on their years of anaesthesia experience.




Figure 1 Patient flow diagram.

All intubations were performed in the presence of two laryngoscopists, one of them being either senior or consultant anaesthesiologist.

Preoxygenation was done for at least 3 min by using either bag valve mask with oxygen flow of 15 l per minute or non-invasive ventilation using 100% fraction of inspired oxygen. The medications for induction were intravenous (IV) fentanyl (1–2 µg/kg) in every patient, with either propofol (1.5–2.0 mg/kg) IV or thiopentone (3–5 mg/kg) IV. Neuromuscular blockade was obtained by using succinylcholine (1–1.5 mg/kg) IV (except patients with hyperkalemia, burn injury older than 24 h) or Rocuronium (0.9 mg/kg) IV. Laryngoscopies were performed using the method as per random allocation, after three failed attempts of intubations alternative techniques were used and subsequently they were not included in the analysis. Stylet was used as when required and external laryngeal manipulations were performed as per laryngoscopist's instruction. Alternative techniques such as gum elastic bougie, laryngeal mask airway (LMA), intubating LMA, intubation endoscope, cricothyrotomy set were used based on the in-house difficult airway algorithm. An intubation attempt was defined as introduction of laryngoscope and its subsequent removal with or without endotracheal tube (ETT) placement. Correct placement of ETT in first attempt by an individual laryngoscopist was defined as first-pass successful intubation. Successful placement of ETT was confirmed by both auscultation and mainstream capnography (normal appearing waveform over four or more breathing cycles).

The laryngoscopist documented the laryngoscopic view by using the modified Cormack & Lehane (C&L) classification and Percentage of Glottic Opening scale (POGO) and other intubation characteristics such as the indication for the intubation, number of intubation attempts, intubation success,

and ease of intubation in relation to the video laryngoscopic view and satisfaction scores.^{4,5}

Macintosh laryngoscopy (ML) has up to 69% success rate during first-pass orotracheal intubation, where as the success rate is 79% patients if video laryngoscope (VL) is used.^{1,3,6,7} Assuming 80% success rate of first-pass orotracheal intubation during VL, with type I error set at 5% and type II error set at 20%, 104 patients were needed in each group (i.e. 208 patients in total), to detect a difference of 18.5% in success rate.

Baseline and demographic data were expressed as the mean ± standard deviation for Gaussian variables. The comparison of the two proportions was performed with the use of the chi-square test or Fischer's exact test when appropriate. The comparisons of means and medians were performed using Student's *t* test and Mann–Whitney test, respectively. The differences were considered statistically significant if $P < 0.05$. Data were analyzed using MedCalc software (version 16.0, Ostend, Belgium).

Results

A total of 263 patients were evaluated, 248 patients were randomized into two groups (VL – videolaryngoscopy and ML – Macintosh laryngoscopy) and data of 218 patients were available for the final analysis (Fig. 1). Both the groups were comparable in terms of their demographic variables (Table 1). Sequential Organ Failure Assessment II score and the indications for intubation were similar between the two groups (Table 1).

Successful first-pass intubation was significantly higher in VL group (84% vs 57%, $P < 0.001$) (Table 2). Reciprocally, higher number of patients in ML group needed two or more than two attempts for successful intubation ($P < 0.05$). The

Table 1 Baseline characteristics of study population.

	Macintosh laryngoscopy (n = 110)	C-MAC video laryngoscopy (n = 108)	P-value
Age in years, mean (SD)	45.8 (16.2)	48.3 (16.8)	0.26
Gender (male/female)	67/43	63/45	0.92
Body mass index, mean (SD)	24.8 (7.6)	23.9 (6.8)	0.36
Sequential organ failure assessment score, mean (SD)	9.1 (3.9)	8.8 (4.2)	0.58
Stylet use	41/110 (37.3)	71/108 (65.7)	<0.0001
Stylet use in first attempt	12/63 (19.0)	55/91 (60.4)	<0.0001
Stylet use in second attempt	21/27 (77.8)	13/14 (92.8)	0.23
Stylet use after second attempt	20/20 (100%)	3/3 (100%)	–
Indication for intubation			
Neurological failure	56 (50.9%)	58 (53.7%)	0.68
Respiratory failure	38 (34.54%)	34 (31.5%)	0.63
Circulatory failure	4 (3.6%)	5 (4.63%)	0.70
Others	12 (10.9%)	11 (10.18%)	0.86

Table 2 Intubation characteristics (number, percentage).

	Macintosh	CMAC	P-value
Cormack and Lehane grade			
Grade 1	49 (44.54%)	91 (84.25%)	<0.0001
Grade 2	28 (25.45%)	7 (6.48%)	0.0003
Grade 3	21 (19.09%)	6 (5.55%)	0.0047
Grade 4	12 (10.90%)	4 (3.70%)	0.0752
Percentage of glottic opening	54 ± 23	81 ± 11	<0.0001
Successful first-pass intubation			
Intention-to-treat analysis	63/124 (50.8)	91/124 (73.37)	0.0003
Per-protocol analysis	63/110 (57.27)	91/108 (84.26)	<0.0001
No. of intubation attempts			
1	63 (57.27%)	91 (84.26%)	<0.0001
2	27 (24.54%)	14 (12.96%)	0.044
>2	20 (18.18%)	3 (2.78%)	0.0005
Laryngoscopist's satisfaction score			
0	3 (2.73%)	1 (0.93%)	0.63
1	3 (2.73%)	3 (2.78%)	0.69
2	5 (4.55%)	6 (5.56%)	0.97
3	21 (19.09%)	14 (12.96%)	0.29
4	68 (61.81%)	84 (77.78%)	0.015

incidence of difficult intubation, at least two failed intubation attempts, was 18% and 3% in the ML and VL groups, respectively (Table 2). Stylet use (Table 1) was higher in VL group as compared to ML group during first (VL vs ML, 65.7% vs 37.3%, $P < 0.0001$) and second attempt intubation (VL vs ML, 92.8% vs 77.8%, $P = 0.23$).

Based on modified C&L classification, higher number of patient in ML group had difficult visualization of the glottis in comparison to VL group. The glottis could not be

visualized in 33 (30%) and 10 (9%) patients (C&L grade 3 or 4), was only partially visualized in 28 (25%) and 7 (6%) patients (C&L grade 2) and was fully visualized in 49 (44%) and 91 (84%) patients (C&L grade 1) in ML and VL groups, respectively. POGO scale showed significant improvement in glottis view in VL group compared to ML group (81 vs 54, $P < 0.0001$) (Table 1). Junior laryngoscopists could successfully intubate higher number of patients in VL group in first attempt (57.9% vs 27.5%, VL vs ML, $P = 0.007$). Intubation

Table 3 Comparison of intubation attempts (number, percentage).

	Macintosh	C-MAC	P-value
<i>Successful intubation in one attempt</i>			
Junior	11 (27.5%)	22 (57.89%)	0.007
Senior	41 (70.69%)	37 (72.55%)	0.8307
Consultant	11 (91.67%)	18 (94.74%)	0.7388
<i>Successful intubation in two attempt</i>			
Junior	16 (40.0%)	12 (31.58)	0.4414
Senior	12 (20.69)	11 (21.57%)	0.9110
Consultant	1 (8.33%)	1 (5.26%)	0.7388
<i>Successful intubation in more than two attempts</i>			
Junior	13 (32.5%)	4 (10.53%)	0.0196
Senior	5 (8.62%)	3 (5.9%)	0.5888
Consultant	0 (0)	0 (0)	0

success rates were similar in both the groups when experience of the anaesthesiologist was concerned, except in the subgroup as mentioned above (Table 3).

Discussion

Endotracheal intubation in critically ill patients differs from intubation in the well controlled operation theatre environment in many ways such as: illness and instability of patients, limitations in terms of time spent preparation, patient positioning, equipments, medications, preoxygenation, operator experience, optimum glottic visualization. In the present study, use of the C-MAC® videolaryngoscope significantly improved percentage of glottis opening during airway management in the ICU. During video laryngoscopy, looking around the corner improves the optical access, but there can be difficulty in negotiating the tip of the tube into larynx with increased risk of contact with the anterior larynx wall. Increased usage of stylet in VL group during all attempts were the result of this phenomenon, which has been described earlier with the use of several video laryngoscopes.⁸

In patients with history of difficult laryngoscopy, C-MAC laryngoscopy had better glottis view in 94% of patients.⁹ Patients with predicted difficult airway had better optical access and more successful first attempt intubation during C-MAC laryngoscopy as compared to Macintosh laryngoscopy.¹⁰ There has been mixed result from various studies when VL was compared with ML in ICU setting. A recent multi-centre randomized controlled trial (RCT) (MACMAN trial) conducted seven ICUs in France found no improvement in the frequency of successful first-pass intubation by using VL compared to ML.² The outcome difference between present study and MACMAN trial can be attributed to single centre vs multicenter, type of video laryngoscope, observer bias, laryngoscopist's background and skill level and experience with the VL. Various studies¹¹⁻¹⁴ including two meta-analyses^{8,15} had reported increased first pass intubation success rate of VL over ML in non-ICU settings, but there were methodological flaws and heterogeneity. At the same time, evidences from other studies including two RCTs in ICU had shown no

improvement in the success rate of video laryngoscope over direct one during first-pass intubation.^{9,10}

A recent systematic review, which included intubations done in operation theatre, had reported reduced intubation failure, easier intubation in patients with known or predicted difficult airway and likely improvement of glottic view and reduction in number of laryngoscopy attempts. There was lack of evidence to support the ability of VL to reduce the time required for intubation and decrease incidences of hypoxia or respiratory complications.¹⁶ In this study, though VL group had significantly higher successful first-pass intubation, but in terms of overall successful intubation, both the groups were similar.

Intubation at locations other than operating room is commonly met with many difficulties. The incidence of difficult laryngoscopies reported for the operating room (5%) is significantly less frequent than in other settings.¹⁷ Various studies have stated that the occurrence of difficult intubations in critically ill patients ranges from 10 to 22%.¹⁸⁻²⁰ In our study, the rate of difficult laryngoscopy (C&L grade 3 and 4) using ML was 30%. We attribute the involvement of anesthesiologists with different years of experience to the higher incidence of difficult airway in the current study. In earlier studies, VL had a better glottic visualization but not higher first-pass successful intubation and higher usage of stylet or gum elastic bougie.¹⁻³ In the present study, we reported better glottic visualization, laryngoscopist's satisfaction, higher successful first-pass intubation and increased stylet usage in VL group as compared to ML group. Similar to the result of MACMAN trial,³ we had a significantly higher usage of stylet in the VL group as compared to ML group. Gum elastic bougie as compared to stylet improved the success rate during Macintosh laryngoscopy in cases with poor glottic visualization.²¹ At the same time, gum elastic bougie has been found to be as efficient as stylet during video laryngoscopy.²² Routine use of stylet during video laryngoscopy had not improved the rate of successful first-pass intubation and French guidelines recommends avoidance of stylet and use of gum elastic bougie in case of difficult orotracheal intubation.²³

In terms of experience, juniors had a significantly higher successful first-pass intubation by using VL, but the difference was not significant in the hands of senior and consultant

anaesthesiologists. In an earlier prospective study conducted in ICU, there were no significant differences in successful intubation attempt when experience was taken into account by using video and direct laryngoscopy.¹ The finding from current study indicates that in the hand of less experienced laryngoscopists, video laryngoscope may increase the success rate of laryngoscopy and intubation in intensive care setup. But further studies are required to compare the time needed for intubation and rates of complication between the different laryngoscopies. Various factors that can confound include the study population, place of use, use and type of neuromuscular blockade, laryngoscopist characteristics, type of VL, difficult mask ventilation and intubation, involvement of independent trained data collector.

Limitations of our study were: (a) only Macintosh blade of C-MAC not the more curved D-blade was compared with direct laryngoscopy was used, (b) predictors of difficult laryngoscopy were not taken into account, (c) neuromuscular monitoring was not used to assess the adequacy of muscle relaxation, (d) time required for successful intubation and complications of intubation such as changes in oxygenation and haemodynamic parameters were not taken into account, (e) number of intubations performed by various laryngoscopists were not the same, juniors performed significantly higher number of laryngoscopies, and (f) use of subjective methods of assessment such as Cormack and Lehane grade, POGO score and laryngoscopist's satisfaction always introduce observer's bias. Future studies may plan alternative objective ways of assessment to minimize such bias.

Conclusion

Use of VL not only improved glottis visualization, but also increased the rate of successful first-pass intubation and especially in less experienced hands in ICU. Despite equivocal support for video and direct laryngoscopy, further studies involving laryngoscopists of different expertise are required to evaluate usefulness various available video laryngoscopes as compared to Macintosh laryngoscope.

Funding

None to disclose.

Authors' contribution

Samarjit Dey, conception and design of the study, data acquisition.

Debasis Pradhan, conception and design, analysis and interpretation of data.

Priyam Saikia, analysis and interpretation of data, draft of the article, final approval.

Prithwis Bhattacharyya, conception and design, critical review of the intellectual content and final approval.

Hariom Khandelwal, data acquisition, draft of the article.

Adarsha KN, data acquisition, draft of the article.

Conflicts of interest

None.

Acknowledgements

This work would not have been possible without the help and commitments of all the nursing and medical staffs of Department of Anaesthesiology and Critical Care, NEIGRIHMS, Shillong, India.

References

1. Noppens RR, Geimer S, Eisel N, David M, Piepho T. Endotracheal intubation using the C-MAC® video laryngoscope or the Macintosh laryngoscope: a prospective, comparative study in the ICU. *Crit Care*. 2012;16:R103.
2. Lascarrou JB, Boisrame-Helms J, Bailly A, Le Thuaut A, Kamel T, Mercier E, et al. Video laryngoscopy vs direct laryngoscopy on successful first-pass orotracheal intubation among ICU patients: a randomized clinical trial. *JAMA*. 2017;317:483–93.
3. Goksu E, Kilic T, Yildiz G, Unal A, Kartal M. Comparison of the C-MAC video laryngoscope to the Macintosh laryngoscope for intubation of blunt trauma patients in the ED. *Turk J Emerg Med*. 2016;16:53–6.
4. Yentis SM, Lee DJ. Evaluation of an improved scoring system for the grading of direct laryngoscopy. *Anaesthesia*. 1998;53:1041–4.
5. Ochroch EA, Hollander JE, Kush S, Shofer FS, Levitan RM. Assessment of laryngeal view: percentage of glottic opening score vs Cormack and Lehane grading. *Can J Anaesth*. 1999;46:987–90.
6. De Jong A, Clavieras N, Conseil M, Coisel Y, Moury PH, Pouzeratte Y, et al. Implementation of a combo videolaryngoscope for intubation in critically ill patients: a before–after comparative study. *Intensive Care Med*. 2013;39:2144–52.
7. Wang HE, Reitz SR, Hostler D, Yealy DM. Defining the learning curve for paramedic student endotracheal intubation. *Prehosp Emerg Care*. 2005;9:156–62.
8. Griesdale DE, Liu D, McKinney J, Choi PT. Glidescope® video-laryngoscopy versus direct laryngoscopy for endotracheal intubation: a systematic review and meta-analysis. *Can J Anesth*. 2012;59:41–52.
9. Griesdale DE, Chau A, Isac G, Ayas N, Foster D, Irwin C, et al., Canadian Critical Care Trials Group. Video-laryngoscopy versus direct laryngoscopy in critically ill patients: a pilot randomized trial. *Can J Anesth*. 2012;59:1032–9.
10. Janz DR, Semler MW, Lentz RJ, Matthews DT, Assad TR, Norman BC, et al. Randomized trial of video laryngoscopy for endotracheal intubation of critically ill adults. *Crit Care Med*. 2016;44:1980.
11. Lakticova V, Koenig SJ, Narasimhan M, Mayo PH. Video laryngoscopy is associated with increased first pass success and decreased rate of esophageal intubations during urgent endotracheal intubation in a medical intensive care unit when compared to direct laryngoscopy. *J Intensive Care Med*. 2015;30:44–8.
12. Kory P, Guevarra K, Mathew JP, Hegde A, Mayo PH. The impact of video laryngoscopy use during urgent endotracheal intubation in the critically ill. *Anesth Analg*. 2013;117:144–9.
13. Mosier JM, Whitmore SP, Bloom JW, Snyder LS, Graham LA, Carr GE, et al. Video laryngoscopy improves intubation success and reduces esophageal intubations compared to direct laryngoscopy in the medical intensive care unit. *Crit Care*. 2013;17:R237.

14. Hypes CD, Stolz U, Sakles JC, Joshi RR, Natt B, Malo J, et al. Video laryngoscopy improves odds of first-attempt success at intubation in the intensive care unit. A propensity-matched analysis. *Ann Am Thorac Soc.* 2016;13:382–90.
15. De Jong A, Molinari N, Conseil M, Coisel Y, Pouzeratte Y, Belafia F, et al. Video laryngoscopy versus direct laryngoscopy for orotracheal intubation in the intensive care unit: a systematic review and meta-analysis. *Intensive Care Med.* 2014;40:629–39.
16. Lewis SR, Butler AR, Parker J, Cook TM, Schofield-Robinson OJ, Smith AF. Videolaryngoscopy versus direct laryngoscopy for adult patients requiring tracheal intubation: a Cochrane Systematic Review. *Br J Anaesth.* 2017;119:369–83.
17. Hagberg CA. Benumof's airway management. Philadelphia, USA: Mosby; 2007.
18. Jaber S, Amraoui J, Lefrant JY, Arich C, Cohendy R, Landreau L, et al. Clinical practice and risk factors for immediate complications of endotracheal intubation in the intensive care unit: a prospective, multiple-center study. *Crit Care Med.* 2006;34:2355–61.
19. Martin LD, Mhyre JM, Shanks AM, Tremper KK, Kheterpal S. 3,423 emergency tracheal intubations at a University Hospital: airway outcomes and complications. *J Am Soc Anesthesiol.* 2011;114:42–8.
20. Le ST, Wolter P, Rusterholtz T, Harlay M, Gayol S, Sauder P, et al. Complications of difficult tracheal intubations in a critical care unit. *Ann Fr Anesth Reanim.* 2000;19:719–24.
21. Noguchi T, Koga K, Shiga Y, Shigematsu A. The gum elastic bougie eases tracheal intubation while applying cricoid pressure compared to a stylet. *Can J Anaesth.* 2003;50:712.
22. Batuwitige B, McDonald A, Nishikawa K, Lythgoe D, Mercer S, Charters P. Comparison between bougies and stylets for simulated tracheal intubation with the C-MAC D-blade videolaryngoscope. *Eur J Anaesthesiol.* 2015;32:400–5.
23. Langeron O, Bourgain JL, Laccoureye O, Legras A, Orliaguet G. Difficulty airway algorithms and management: question 5. French Society of Anesthesia and Reanimation. *Ann Anesth Reanim.* 2008;27:41.