



ORIGINAL

Usefulness of diaphragmatic ultrasound in predicting extubation success[☆]



F. Varón-Vega^{a,b,c}, Á. Hernández^a, M. López^a, E. Cáceres^a, L.F. Giraldo-Cadavid^{d,e}, A.M. Uribe-Hernandez^{a,b,d,*}, S. Crevoisier^f

^a Unidad de Cuidado Intensivo Médico, Fundación Neumológica Colombiana, Bogotá, Colombia

^b Fundación Cardioinfantil-Instituto de Cardiología, Bogotá, Colombia

^c Departamento de Anestesiología y Cuidados Intensivos, Universidad de Navarra, Pamplona, Spain

^d Departamento de Investigación, Fundación Neumológica Colombiana, Bogotá, Colombia

^e Departamento de Epidemiología y de Medicina Interna, Universidad de La Sabana, Chía, Colombia

^f Medicina Crítica y Cuidado Intensivo, Universidad de La Sabana, Chía, Colombia

Received 22 July 2019; accepted 24 October 2019

Available online 5 May 2020

KEYWORDS

Critical illness;
Airway extubation;
Diaphragm/diagnostic imaging;
Intensive care units;
Predictive value of tests;
Respiration;
Artificial;
Sensitivity and specificity;
Ultrasonography;
Ventilator weaning;
Weaning from ventilation

Abstract

Objective: To evaluate the diagnostic accuracy of diaphragmatic ultrasound in predicting extubation success.

Design: A diagnostic accuracy study was carried out.

Scope: Intensive Care Unit of an Academic hospital in the city of Bogotá (Colombia).

Patients or participants: A consecutive sample of patients >18 years of age subjected to invasive mechanical ventilation for >48 h.

Interventions: Diaphragmatic ultrasound evaluation at the end of spontaneous ventilation testing.

Main variables of interest: Diaphragmatic excursion (DE, cm), inspiration time (TPIA_{dia}, s), diaphragm contraction speed (DE/TPIA_{dia}, cm/s) and total time (T_{tot}, s) were evaluated, together with thickening fraction (TF_{di}, %).

Results: A total of 84 patients were included, 79.8% (n=67) with successful extubation and 20.2% (n=17) with failed extubation. The variable with the best discriminatory capacity in predicting extubation success was diaphragm contraction speed, with AUC-ROC 0.70 (p=0.008).

Conclusions: Diaphragm contraction speed exhibited acceptable discriminatory capacity. Ultrasound could be part of a multifactorial approach in the extubation process.

© 2019 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.

[☆] Please cite this article as: Varón-Vega F, Hernández Á, López M, Cáceres E, Giraldo-Cadavid LF, Uribe-Hernandez AM, et al. Utilidad de la ecografía diafragmática para predecir el éxito en la extubación. Med Intensiva. 2021;45:226–233.

* Corresponding author.

E-mail address: auribe@neumologica.org (A.M. Uribe-Hernandez).

PALABRAS CLAVE

Enfermedad crítica;
Extubación de la vía aérea;
Diafragma/diagnóstico por imagen;
Unidades de cuidados intensivos;
Valor predictivo de las pruebas;
Respiración; Artificial;
Sensibilidad y especificidad;
Ultrasonografía;
Destete del ventilador;
Retirada de la ventilación

Utilidad de la ecografía diafragmática para predecir el éxito en la extubación**Resumen**

Objetivo: Evaluar la exactitud diagnóstica de la ecografía diafragmática para predecir el éxito en la extubación.

Diseño: Estudio de exactitud diagnóstica.

Ámbito: Unidad de Cuidado Intensivo Médico de un hospital académico de la ciudad de Bogotá (Colombia).

Pacientes o participantes: Muestra consecutiva de pacientes mayores de 18 años con ventilación mecánica invasiva durante más de 48 h.

Intervenciones: Evaluación ecográfica diafragmática al finalizar la prueba de ventilación espontánea.

Variables de interés principales: Se evaluó la excursión diafragmática (ED, cm), el tiempo de inspiración (TPIA_{dia}, s), la velocidad de contracción del diafragma (ED/TPIA_{dia}, cm/s), el tiempo total (T_{tot}, s) y la fracción de engrosamiento (TF_{di}, %).

Resultados: Se incluyeron 84 pacientes, el 79.8% (n=67) con extubación exitosa y el 20.2% (n=17) con extubación fallida. La variable con mejor capacidad discriminadora para predecir éxito en la extubación fue la velocidad de contracción, con un AUC-ROC de 0,70 (p=0,008).

Conclusiones: La velocidad de contracción diafragmática mostró una capacidad discriminadora aceptable. La ultrasonografía podría formar parte de un abordaje multifactorial en el proceso de extubación.

© 2019 Elsevier España, S.L.U. y SEMICYUC. Todos los derechos reservados.

Introduction

The need for mechanical ventilation (MV) is one of the main reasons for admission to the Intensive Care Unit (ICU)¹. Despite its benefits, however, the complications of MV are an important source of patient morbidity–mortality^{2–4}. Establishing the optimum moment for withdrawal of ventilatory support remains one of the greatest challenges for the treating professional team, since late extubation is directly associated to an increased incidence of in-hospital infections, including ventilator-associated pneumonia (VAP), as well as to increased costs, diaphragm dysfunction, worsened quality of life over the middle term, and a longer stay in the ICU and in hospital in general^{5–7}. In contrast, early extubation resulting in a need for reintubation has been associated to a 25–50% increase in patient mortality^{8,9}.

The heterogeneity of the patients admitted to the ICU implies that the causes of extubation failure are also multiple¹⁰; diaphragm dysfunction appears to be implicated in up to 50% of all failed extubations¹¹. This situation is related to the structural and functional changes observed in the muscle fibers after the start of ventilatory support^{12,13}. On the basis of the above, one of the cornerstones of patient management is the facilitation of early rehabilitation^{1,14}. To date, no reference parameters have been able to predict extubation success. The most widely used clinical parameters are the rapid shallow breathing index (RSBI), vital capacity (VC) and peak inspiratory pressure (PI_{max}), among others^{15,16}. There is great variability in the cut-off points and diagnostic precision of these parameters¹¹, and none of them reflect the integrity of diaphragm structure and function.

In this context, in recent years ultrasound at the patient bedside (point of care) has become one of the tools of choice in the ICU due to its accessibility and low cost. It allows us to assess structure and function quantitatively and qualitatively before, during and after extubation¹⁷. A range of ultrasound parameters have been studied to date: diaphragmatic excursion (DE), thickening fraction (TF_{di}), contraction velocity (V)^{18–20} and even variations in rapid shallow breathing index (respiratory frequency/DE)²¹. The cut-off points of these parameters are likewise diverse, with great variability in performing the test.

The present study was carried out to evaluate the diagnostic accuracy of diaphragmatic ultrasound at the patient bedside in predicting extubation success.

Patients and methods

A prospective, observational cohort study on diagnostic accuracy was carried out.

Study population

The study was carried out in the ICU of an academic institution in the city of Bogotá (Colombia), with the consecutive inclusion of all patients over 18 years of age subjected to invasive MV for over 48 h and who met the following inclusion criteria: (1) resolution of the cause of respiratory failure; (2) hemodynamic stability; (3) metabolic equilibrium; (4) optimum level of consciousness; and (5) indication of spontaneous breathing test (SBT) (Table 1).

The exclusion criteria were: (1) neuromuscular disease; (2) previous diaphragmatic paralysis; (3) use of

Table 1 Inclusion criteria.

Blood gas parameters	PaO ₂ ≥ 55 mmHg with FiO ₂ < 40% PEEP < 8 cm H ₂ O PaO ₂ /FiO ₂ > 175 PaO ₂ /PAO ₂ > 0.3 pH ≥ 7.35 ≤ 7.48
Hemodynamic stability	Norepinephrine <0.1 µg/kg/min Dopamine <5 µg/kg/min Dobutamine <5 µg/kg/min Hemoglobin ≥7 g/dl
Level of consciousness	Glasgow coma score ≥12 CAM-ICU negative
Spontaneous breathing test (SBT)	Modality: T-tube: patients – pressure support Daily evaluation by medical staff/respiratory therapy to define moment of SBT 30 min duration
Metabolic equilibrium	pH ≥ 7.35 ≤ 7.48 Temperature ≤38 °C Serum sodium, potassium, phosphorus in normal ranges

neuromuscular blockers during admission to the Unit; (4) pneumothorax or pneumomediastinum; and (5) pregnancy.

The following data were compiled at baseline: patient age and gender, cause of respiratory failure, duration of MV, and arterial gas and laboratory test values before extubation.

Measurements

The decision to perform the spontaneous breathing test (SBT) was assessed daily by the supervising medical team and the respiratory therapy group of the Unit. After the 30 min of the SBT, diaphragmatic function was assessed by ultrasound, with calculation of the rapid shallow breathing index as part of the standard evaluation for establishing extubation.

The diaphragmatic measurements were carried out by intensivists trained in ultrasound in the critical care setting, using a Sonocare ultrasound system (Sonosite EDGE 03VRYF). A 1–5 MHz transducer was used for the M-mode evaluation of diaphragmatic excursion (DE, cm), time to peak inspiratory amplitude (TPIA_{dia}, s), contraction velocity of the diaphragm (DE/TPIA_{dia}, cm/s) and total time (T_{tot}, s). The thickening fraction (TF_{di}, %) was evaluated with a 6–13 MHz transducer in M-mode (Fig. 1 and Table 2). Diaphragm dysfunction was defined as DE < 1 cm or paradoxical motion¹⁸.

The measurements were made only in the right half of the diaphragm, with the patient in the semi-sitting position (headrest raised 45 degrees). The transducer was positioned just below the ribcage, between the clavicular midline and the anterior axillary line. The ultrasound beam was directed cephalad, perpendicular to the posterior third of the diaphragm. Three operators performed the ultrasound explorations in the ICU, distributed as follows: 45 explorations made by an intensivist during the morning shift, and 20 explorations each performed by two intensivists in the afternoon. Before the study, a 12-h training session with an expert radiologist was held to ensure standardization of the ultrasound measurements.

Before extubation, all patients were reconnected to their previous ventilation mode during 1 h²².

Study objectives

The primary study objective was to determine the accuracy of diaphragmatic ultrasound as a predictor of the success of weaning from MV. Successful extubation was defined as the capacity to maintain spontaneous breathing for over 48 h without ventilatory assistance after extubation. Failed extubation in turn was defined as the need for patient reintubation in under 48 h⁹.

As secondary objective, we evaluated the differences in extubation success or failure in relation to the different demographic, clinical and ultrasound parameters, and diaphragm dysfunction (defined as DE < 1 cm or paradoxical motion)¹⁸.

Ethical aspects

The study protocol was approved by the local Ethics Committee (Ref. no.: 205 of 2014). The study was considered to pose minimal risks for patients according to resolution 8430 of 1993 of the Colombian Ministry of Health. Informed consent was obtained from all the participants.

Statistical analysis

Convenience non-probability sampling was performed, calculating a sample size of 84 patients based on an estimated prevalence of 20% for extubation failure, with a sensitivity of 90% and a specificity of 86%, a 95% confidence interval with an area under the receiver operating characteristic curve (AUC-ROC) of 0.15, an alpha error of 0.05, and a statistical power of 80%^{23,24}.

Central tendency and dispersion measures were used for the quantitative variables, and frequencies and percentages for the qualitative variables.




Figure 1 Ultrasound measures used to assess the success of extubation. Mean in M-mode. 1.1: Thickening fraction (TF_{di}, %), expiratory thickness (A), inspiratory thickness (B). 1.2: Measurement of diaphragmatic excursion (a) (DE, cm), time to peak inspiratory amplitude (b) (TPIA_{dia}, s), diaphragmatic contraction velocity (DE/TPIA_{dia} [cm/s]). 1.3: Time to peak inspiratory amplitude (a) (TPIA_{dia}, s), total time (b) (s).

Table 2 Diaphragmatic measurements.

Measurement	Evaluation M-mode
Diaphragmatic excursion (DE, cm)	Excursion amplitude from start of contraction to maximum inspiration
Time to peak inspiratory amplitude (TPIA _{dia} , s)	Time from start of diaphragmatic contraction to maximum inspiration
Diaphragmatic contraction velocity (cm/s)	Diaphragmatic excursion (DE)/time to peak inspiratory amplitude (TPIA _{dia})
Total time (s)	Inspiratory time + expiratory time
Thickening fraction (TF _{di} , %)	Diaphragmatic thickness at end of inspiration – diaphragmatic thickness at end of expiration / diaphragmatic thickness at end of expiration × 100

The patients were divided into two groups according to the primary outcome (extubation success or failure). The chi-squared test was used for the bivariate comparison of categorical variables. The Student *t*-test in turn was used for the comparison of continuous variables exhibiting a normal distribution, while parameters with a non-normal distribution were contrasted by means of the Mann–Whitney *U*-test. The quantitative variables with a non-parametric distribution that included follow-up time were subjected to negative binomial regression or Poisson analysis, depending on whether the standard deviation (SD) was greater or smaller than the mean of such variables.

We calculated the operator characteristics of each of the ultrasound measures to predict extubation success or failure, and ROC curves were plotted to establish the diagnostic accuracy of each of the ultrasound parameters. The point of maximum discriminating capacity was selected, based on the Youden index.

The AUC-ROC was interpreted as follows²⁵: =0.5, no discriminating capacity; >0.7–0.79: acceptable discriminating capacity; >0.8–0.89: excellent discriminating capacity; >0.9: outstanding discriminating capacity.

The data were analyzed using the SPSS version 20 statistical package and MedCalc version 19.

Results

A total of 84 patients were included in the study, and no losses were recorded. The general characteristics of the study sample are described in Table 3. The median patient age was 58 years (range 35–51), with a female predominance (56%). Most of the patients (88%) presented medical conditions as the indication of MV, with an APACHE II severity score of 21 (17–28). The method of choice for SBT was the T-tube technique (85.7%), versus pressure support (14.3%).

Successful extubation was achieved in 79.8% of the patients (*n* = 67), and extubation failed in the remaining 20.2% of the cases (*n* = 17). The comparison of results between both groups is shown in Table 3. There were no significant differences between the groups in terms of the demographic or clinical characteristics. However, the patients with failed extubation presented APACHE II scores that were slightly higher than those recorded in the patients with successful extubation. The rapid shallow breathing index was also similar in both groups, with slightly higher scores among the patients with successful extubation versus those with failed extubation: 48 (36–64) and 40 (32–62), respectively. The SBT choice likewise showed no significant differences. The duration of MV and of ICU stay was slightly longer in the failed extubation group, though the differences failed to reach statistical significance.

Table 3 General characteristics of the study sample. Total population, extubation success and failure groups.

	Total	Success n = 67 (79.8%)	Failure n = 17 (20.2%)	p
Age (years), median (IQR)	58 (35–51)	58 (34–72)	59 (49–64)	0.79
Gender, n (%)				0.4
Female	47 (56)	39 (83)	8 (17)	
Male	37 (44)	28 (75.7)	9 (24.3)	
Type of patient, n (%)				0.39
Medical	74 (88)	58 (78.4)	16 (21.6)	
Surgical	10 (12)	9 (90)	1 (10)	
Cause of respiratory failure, n (%)				
Pulmonary sepsis	21 (25)	17 (80.9)	4 (19.1)	0.87
Extrapulmonary sepsis	26 (31)	20 (76.9)	6 (23.1)	0.56
Neurological	19 (22.6)	15(78.9)	4 (21.1)	0.92
Postoperative	6 (7.1)	6 (100)	0 (0)	0.20
Others	12 (14.3)	9 (75)	3 (25)	0.65
APACHE II, median (IQR)	21 (17–28)	19 (17–28)	21 (18–25)	0.52
Rapid shallow breathing index, median (IQR)	47 (36–63)	48 (36–64)	40 (32–62)	0.38
Weaning test, n (%)				0.73
T-tube	72 (85.7)	57 (79.2)	15 (20.8)	
Pressure support	12 (14.3)	10 (83.3)	2 (16.7)	
Diaphragmatic measurements				
DE (cm), median (IQR)	2.18 (1.6–2.75)	2.22 (1.66–2.75)	2.02 (1.63–2.31)	0.44
Excursion <10 mm, n (%)	1 (1.2)	0 (0)	1 (100)	0.046
DE/TPIA _{dia} (cm/s), median (IQR)	2.74 (1.90–3.33)	2.90 (2.00–4.01)	2.02 (1.49–2.80)	0.013
TPIA _{dia} (s), median (IQR)	0.79 (0.64–1.02)	0.80 (0.67–0.95)	0.77 (0.62–1.08)	0.76
T _{tot} (s), mean (DE)	2.96 (0.65)	2.97 (0.65)	2.93 (0.65)	0.84
TF _{di} (%), median (IQR)	31 (24–45)	32 (27–47)	30 (21–35)	0.10
TF _{di} > 30%, n (%)	47 (56)	39 (83)	8 (17)	0.40
Outcomes, median (IQR)				
MV time, days	5 (3–10)	5 (3–9)	7 (4–11)	0.98
Weaning time, days	2 (1–3)	2 (1–3)	3 (1–3)	0.65
ICU stay (days), median (IQR)	10 (7–17)	10 (6–17)	11 (8–16)	0.69

APACHE II: Acute Physiology and Chronic Health Evaluation II; SD: standard deviation; DE: diaphragmatic excursion; IQR: interquartile range; TF_{di}: thickening fraction; TPIA_{dia}: time to peak inspiratory amplitude; T_{tot}: total time; ICU: Intensive Care Unit; V: contraction velocity; MV: mechanical ventilation.

Table 4 Operator characteristics.

Contraction velocity (cm/s)	Sensitivity	Specificity	PPV	NPV	LR+	LR–
>1.74	85	42	85	41	1.45	0.36
>2.90	46.27	88.24	94	29	3.93	0.61
>4.3	19.4	88.24	87	22	1.62	0.91

LR+: positive likelihood ratio; LR–: negative likelihood ratio; NPV: negative predictive value; PPV: positive predictive value.

Of the different ultrasound parameters, differences were only observed for contraction velocity and diaphragm dysfunction (DE <1 cm); the latter was only present in 1.2% of the total cases. The ROC curve for diaphragm contraction velocity is shown in Fig. 2. This variable presented AUC 0.70 (p=0.008; 95% CI: 0.58–0.79). Three cut-off points were calculated (shown in Table 4). The point of maximum discriminating capacity according to the

Youden index was >2.9 cm/s, with a sensitivity of 4.27% (95% CI: 0.34–0.59) and a specificity of 88.24% (95% CI: 0.50–0.93). Values of over 1.74 cm/s were associated to greater sensitivity (85% [95% CI: 0.74–0.93]) and lesser specificity (41% [95% CI: 0.19–0.67]). On the other hand, thresholds of over 4.3 cm/s showed a sensitivity of 19.4% (95% CI: 0.11–0.30) and a specificity of 88.24% (95% CI: 0.64–0.99).




Figure 2 Receiver operating characteristic curve for contraction velocity and extubation success. AUC 0.70 ($p=0.008$ [95% CI: 0.58–0.79]).

Discussion

The main finding of our investigation was an acceptable discriminating capacity (AUC 0.70; $p=0.008$ [95% CI: 0.58–0.79]) in predicting extubation success or failure on assessing diaphragmatic contraction velocity as isolated marker in critical patients admitted to the ICU.

A range of ultrasound parameters have been evaluated in the MV weaning process²⁶. The most widely studied are DE and TF_{di} . Two meta-analyses have compiled the available evidence. The first included 19 observational studies with a total of 1068 patients. The DE values were between 10 and 27 mm and the TF_{di} values between 20 and 36%. The analysis of the ROC curve for TF_{di} yielded AUC 0.87, while in the case of DE the lack of data and their heterogeneity only allowed the estimation of a cumulative specificity of 75% and a sensitivity of 80%²⁷. The second meta-analysis, published in 2018, evaluated 13 observational studies with a total of 742 patients. The findings were similar to those of the previous meta-analysis, with good performance being observed for both DE and TF_{di} , with AUC 0.859 and 0.838, respectively²⁸. Nevertheless, both meta-analyses were characterized by heterogeneity in defining extubation failure, in the selection of patients, the indication of intubation, and the selection of cut-off points. In our study, neither DE nor TF_{di} showed differences between the extubation success and failure groups. The only variable exhibiting statistically significant differences was diaphragmatic contraction velocity. This variable is taken to be an indirect measure of diaphragm contraction strength. In healthy individuals, the normal value is estimated to be 1.3 ± 0.4 cm/s. The role of this variable has not been widely studied to date, though higher values appear to be related to an increased probability of successful extubation²⁹. We recorded values of >2.9 cm/s (2.00–4.01) in the successful extubation group

versus >2.02 cm/s in the failed extubation group (1.49–2.80) ($p=0.013$). A number of cut-off points were analyzed with a view to optimizing the discriminating capacity of the test. A velocity >1.74 cm/s showed high sensitivity. However, the low associated specificity could indicate a greater number of patients at risk of reintubation. On the other hand, thresholds above 4.3 cm/s (sensitivity 19.4% and specificity 88.24%) would limit the start of weaning in clinical practice. The maximum discriminating capacity was established at >2.9 cm/s, with only acceptable overall performance (AUC 0.70; $p=0.008$). To date, three studies have defined velocity as a differential marker in the extubation process. The first study described much lower thresholds than those established in our study (>0.8 cm/s), with a sensitivity of 100%, a specificity of 86.67% and an outstanding discriminating capacity (AUC 0.93)²⁹. The second study, with slightly higher thresholds (0.92 cm/s), reported a sensitivity of 100%, with low specificity (45%) in discriminating extubation success (AUC 0.66)³⁰. Lastly, the third study evaluated $TPIA_{dia}$ as derived variable ($DE/TPIA_{dia}$), with a discriminating capacity similar to that found in our study (AUC 0.71)¹¹.

The optimum time for suspending MV remains a challenge for multidisciplinary teams in the ICU. The need for reintubation is one of the most feared complications, because of the associated increase in patient mortality⁸. The incidence of extubation failure reported in the literature ranges from 10 to 25%³¹, which coincides with the figure recorded in our study (20.2%).

Muscle trauma with consequent ventilator-induced diaphragm dysfunction (VIDD) is one of the main causes of failed ventilation withdrawal³². In some cases it may go unnoticed; active search on the part of the clinician is therefore essential in this regard³³. To date, the parameter used to evaluate diaphragm function has been direct fluoroscopy, though this technique is of limited applicability in the intensive care setting because of the problems posed by having to transfer critical patients outside the Unit¹¹. For this reason diaphragmatic ultrasound has been found to be very useful as a tool that can be used at the patient bedside. In our study, diaphragm dysfunction was only observed in 1.2% of the cases, in contrast to other series in the literature that have reported a prevalence of 23–36%¹². Ultrasound definition also has limitations, and the technique has not been prospectively contrasted versus the reference standards. Furthermore, the cut-off points described in the literature are heterogeneous. Nevertheless, we consider that the low frequency of diaphragm dysfunction recorded in our study is attributable to the presence of a strict early rehabilitation program that starts upon patient admission to the Unit.

Prolonged MV is one of the consequences of diaphragm dysfunction³⁴. Our results showed a relatively short period of MV, with a median of 5 days, and with no differences between the extubation success and failure groups. Likewise, the ventilation weaning period corresponded to 40% of the ventilation time, in coincidence with the data found in the literature³⁵. The total duration of stay in the Unit was slightly longer in the failure group, though statistical significance was not reached, probably because of statistical power limitations related to the sample size involved.

Taking into account that no clinical or imaging parameter considered isolatedly has been able to predict the outcome of the extubation process, the heterogeneity of the

critically ill makes it necessary to adopt a multimodal approach to ventilation withdrawal³⁶. The multiple systems involved should be evaluated qualitatively and quantitatively as part of the extubation process. Ultrasound could be a tool accompanying traditional evaluation, and in this sense the results of our study with regard to the rapid shallow breathing index (<105 in both groups) confirm its scant clinical usefulness when considered isolatedly. At present there are two main scenarios for ultrasound utilization at the critical patient bedside: (a) application as an evaluation and follow-up strategy in order to avoid muscle trauma^{32,37} and (b) use as part of multifunctional cardiovascular, pulmonary and pleural assessment³⁸. We consider the definition of new integrating indices contemplating systematic ultrasound evaluation to be a priority with a view to improving the outcomes of the extubation process.

The main strength of our study is the identification of contraction velocity as an independent variable for estimating the success of extubation. Despite only acceptable performance in the ROC curves, it deserves becoming the focus of future research.

Our study has a number of limitations. Ultrasound is operator-dependent. Inter- and intra-observer variability in the ventilation weaning scenario has been evaluated only in relation to the measurement of the diaphragmatic thickening fraction³⁹. Our measurements were made by different operators without conducting concordance studies between them; the results of the other ultrasound variables therefore require validation. On the other hand, the lack of distinction between the types of ventilation withdrawal (easy, difficult or prolonged⁹) could modify ultrasound performance as a prognostic tool. In turn, the severity of disease of the patients included in the study, assessed by means of the APACHE II score, was considered to be high; the findings therefore might not be extrapolatable to other ICUs of lesser complexity.

Authorship

Fabio Varon-Vega, Angela Hernandez, Luis Fernando Giraldo: study design and structure.

Mauricio Lopez and Edgar Caceres: data collection.

Fabio Varon-Vega, Ana Maria Uribe, Luis Fernando Giraldo and Stephanie Crevoisier: data analysis and drafting of the manuscript.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Acknowledgement

Thanks are due to the medical and nursing and respiratory therapy staff of the medical ICU of the *Fundación Neumológica Colombiana*.

References

- Girard TD, Alhazzani W, Kress JP, Ouellette DR, Schmidt GA, Truwit JD, et al. An Official American Thoracic Society/American College of Chest Physicians Clinical Practice Guideline: Liberation from mechanical ventilation in critically ill adults rehabilitation protocols, ventilator liberation protocols, and cuff leak tests. *Am J Respir Crit Care Med*. 2017;195:120–33, <http://dx.doi.org/10.1164/rccm.201610-2075ST>.
- Katira BH. Ventilator-induced lung injury: classic and novel concepts. *Respir Care*. 2019;64:629–37, <http://dx.doi.org/10.4187/respcare.07055>.
- Baid H. Patient safety: identifying and managing complications of mechanical ventilation. *Crit Care Nurs Clin North Am*. 2016;28:451–62, <http://dx.doi.org/10.1016/j.cnc.2016.07.005>.
- Melsen WG, Rovers MM, Groenwold RH, Bergmans DC, Camus C, Bauer TT, et al. Attributable mortality of ventilator-associated pneumonia: a meta-analysis of individual patient data from randomised prevention studies. *Lancet Infect Dis*. 2013;13:665–71, [http://dx.doi.org/10.1016/S1473-3099\(13\)70081-1](http://dx.doi.org/10.1016/S1473-3099(13)70081-1).
- Jaber S, Quintard H, Cinotti R, Asehnoune K, Arnal JM, Guitton C, et al. Risk factors and outcomes for airway failure versus non-airway failure in the intensive care unit: a multicenter observational study of 1514 extubation procedures. *Crit Care*. 2018;22:236, <http://dx.doi.org/10.1186/s13054-018-2150-6>.
- Kim WY, Lim CM. Ventilator-induced diaphragmatic dysfunction: diagnosis and role of pharmacological agents. *Respir Care*. 2017;62:1485–91, <http://dx.doi.org/10.4187/respcare.05622>.
- Busico M, Intile D, Sívori M, Irastrorza N, Alvarez AL, Quintana J, et al. Factores de riesgo relacionados al empeoramiento de la calidad de vida (QOL) en pacientes que recibieron ventilación mecánica. Estudio prospectivo multicéntrico. *Med Intensiva*. 2016;40:422–30, <http://dx.doi.org/10.1016/j.medin.2016.01.002>.
- Thille AW, Cortés-Puch I, Esteban A. Weaning from the ventilator and extubation in ICU. *Curr Opin Crit Care*. 2013;19:57–64, <http://dx.doi.org/10.1097/MCC.0b013e32835c5095>.
- Boles JM, Bion J, Connors A, Herridge M, Marsh B, Melot C, et al. Weaning from mechanical ventilation. *Eur Respir J*. 2007;29:1033–56, <http://dx.doi.org/10.1183/09031936.00010206>.
- Heunks L, van der Hoeven JG. Clinical review: The ABC of weaning failure – a structured approach. *Crit Care*. 2010;14:245, <http://dx.doi.org/10.1186/cc9296>.
- Theerawit P, Eksombatchai D, Sutherasan Y, Suwatanapongched T, Kiatboonsri C, Kiatboonsri S. Diaphragmatic parameters by ultrasonography for predicting weaning outcomes. *BMC Pulm Med*. 2018;18:175, <http://dx.doi.org/10.1186/s12890-018-0739-9>.
- Dres M, Demoule A. Diaphragm dysfunction during weaning from mechanical ventilation: an underestimated phenomenon with clinical implications. *Crit Care*. 2018;22:73, <http://dx.doi.org/10.1186/s13054-018-1992-2>.
- Supinski GS, Callahan LA. How important is diaphragm function as a determinant of outcomes for MICU patients in respiratory failure? *Physiology*. 2015;30:336–7, <http://dx.doi.org/10.1152/physiol.00025.2015>.
- Pun BT, Balas MC, Barnes-Daly MA, Thompson JL, Aldrich JM, Barr J, et al. Caring for critically ill patients with the ABCDEF bundle: results of the ICU liberation collaborative in over 15,000 adults. *Crit Care Med*. 2019;47:3–14, <http://dx.doi.org/10.1097/CCM.0000000000003482>.
- Nemer SN, Barbas CS, Caldeira JB, Cárias TC, Santos RG, Almeida LC, et al. A new integrative weaning index of discontinuation from mechanical ventilation. *Crit Care*. 2009;13:R152, <http://dx.doi.org/10.1186/cc8051>.
- González-Castro A, Suárez-Lopez V, Gómez-Marcos V, González-Fernandez C, Iglesias-Posadilla D, Burón-Mediavilla J, et al. Valor de la fracción de espacio muerto (Vd/Vt) como predictor de éxito en la extubación.

- Med Intensiva. 2011;35:529–38, <http://dx.doi.org/10.1016/j.medin.2011.05.016>.
17. Palkar A, Mayo P, Singh K, Koenig S, Narasimhan M, Singh A, et al. Serial diaphragm ultrasonography to predict successful discontinuation of mechanical ventilation. *Lung*. 2018;196:363–8, <http://dx.doi.org/10.1007/s00408-018-0106-x>.
 18. Kim WY, Suh HJ, Hong SB, Koh Y, Lim CM. Diaphragm dysfunction assessed by ultrasonography: influence on weaning from mechanical ventilation. *Crit Care Med*. 2011;39:2627–30, <http://dx.doi.org/10.1097/CCM.0b013e3182266408>.
 19. Rittayamai N, Hemvimon S, Chierakul N. The evolution of diaphragm activity and function determined by ultrasound during spontaneous breathing trials. *J Crit Care*. 2019;51:133–8, <http://dx.doi.org/10.1016/j.jcrc.2019.02.016>.
 20. Matamis D, Soilemezi E, Tsagourias M, Akoumianaki E, Dimassi S, Boroli F, et al. Sonographic evaluation of the diaphragm in critically ill patients. Technique and clinical applications. *Intensive Care Med*. 2013;39:801–10, <http://dx.doi.org/10.1007/s00134-013-2823-1>.
 21. Abbas A, Embarak S, Walaa M, Lutfy SM. Role of diaphragmatic rapid shallow breathing index in predicting weaning outcome in patients with acute exacerbation of COPD. *Int J COPD*. 2018;13:1655–61, <http://dx.doi.org/10.2147/COPD.S161691>.
 22. Fernandez MM, González-Castro A, Magret M, Bouza MT, Ibañez M, García C, et al. Reconnection to mechanical ventilation for 1 h after a successful spontaneous breathing trial reduces reintubation in critically ill patients: a multicenter randomized controlled trial. *Intensive Care Med*. 2017;43:1660–7, <http://dx.doi.org/10.1007/s00134-017-4911-0>.
 23. DiNino E, Gartman EJ, Sethi JM, McCool FD. Diaphragm ultrasound as a predictor of successful extubation from mechanical ventilation. *Thorax*. 2014;69:423–7, <http://dx.doi.org/10.1136/thoraxjnl-2013-204111>.
 24. Machin D, Campbell M, Tan S, Tan S. *Sample size tables for clinical studies*. 3rd ed. Wiley-Blackwell; 2009.
 25. Hosmer DW, Lemeshow S. *Assessing the fit of the model*. In: Hosmer DW, Lemeshow S, editors. *Applied logistic regression*. 2nd ed. New York, NY: John Wiley & Sons, Inc.; 2000. p. 162.
 26. Vetrugno L, Guadagnin GM, Barbariol F, Langiano N, Zangrillo A, Bove T. Ultrasound imaging for diaphragm dysfunction: a narrative literature review. *J Cardiothorac Vasc Anesth*. 2019;33:2525–36, <http://dx.doi.org/10.1053/j.jvca.2019.01.003>.
 27. Llamas-Álvarez AM, Tenza-Lozano EM, Latour-Pérez J. Diaphragm and lung ultrasound to predict weaning outcome: systematic review and meta-analysis. *Chest*. 2017;152:1140–50, <http://dx.doi.org/10.1016/j.chest.2017.08.028>.
 28. Li C, Li X, Han H, Cui H, Wang G, Wang Z. Diaphragmatic ultrasonography for predicting ventilator weaning: a meta-analysis. *Medicine (Baltimore)*. 2018;97:e10968, <http://dx.doi.org/10.1097/MD.0000000000010968>.
 29. Banerjee A, Mehrotra G. Comparison of lung ultrasound-based weaning indices with rapid shallow breathing index: are they helpful? *Indian J Crit Care Med*. 2018;22:435–40, http://dx.doi.org/10.4103/ijccm.IJCCM.331_17.
 30. Palkar A, Narasimhan M, Greenberg H, Singh K, Koenig S, Mayo P, et al. Diaphragm excursion-time index: a new parameter using ultrasonography to predict extubation outcome. *Chest*. 2018;153:1213–20, <http://dx.doi.org/10.1016/j.chest.2018.01.007>.
 31. Subirà C, Hernández G, Vázquez A, Rodríguez-García R, Gonzalez-Castro A, García C, et al. Effect of pressure support vs T-piece ventilation strategies during spontaneous breathing trials on successful extubation among patients receiving mechanical ventilation. *JAMA*. 2019;321:2175–82, <http://dx.doi.org/10.1001/jama.2019.7234>.
 32. Schepens T, Dres M, Heunks L, Goligher EC. Diaphragm-protective mechanical ventilation. *Curr Opin Crit Care*. 2019;25:77–85, <http://dx.doi.org/10.1097/MCC.0000000000000578>.
 33. Dres M, Dubé BP, Mayaux J, Delemazure J, Reuter D, Brochard L, et al. Coexistence and impact of limb muscle and diaphragm weakness at time of liberation from mechanical ventilation in medical intensive care unit patients. *Am J Respir Crit Care Med*. 2017;195:57–66, <http://dx.doi.org/10.1164/rccm.201602-0367OC>.
 34. Carson SS. Outcomes of prolonged mechanical ventilation. *Curr Opin Crit Care*. 2006;12:405–11, <http://dx.doi.org/10.1097/01.ccx.0000244118.08753.dc>.
 35. Lu Z, Xu Q, Yuan Y, Zhang G, Guo F, Ge H. Diaphragmatic dysfunction is characterized by increased duration of mechanical ventilation in subjects with prolonged weaning. *Respir Care*. 2016;61:1316–22, <http://dx.doi.org/10.4187/respcare.04746>.
 36. Heunks LM, Doorduyn J, van der Hoeven JG. Monitoring and preventing diaphragm injury. *Curr Opin Crit Care*. 2015;21:34–41, <http://dx.doi.org/10.1097/MCC.000000000000168>.
 37. Jung B, Gleeton D, Daurat A, Conseil M, Mahul M, Rao G, et al. Conséquences de la ventilation mécanique sur le diaphragme. *Rev Mal Respir*. 2015;32:370–80, <http://dx.doi.org/10.1016/j.rmr.2014.08.013>.
 38. Mayo P, Volpicelli G, Lerolle N, Schreiber A, Doelken P, Vieillard-Baron A. Ultrasonography evaluation during the weaning process: the heart, the diaphragm, the pleura and the lung. *Intensive Care Med*. 2016;42:1107–17, <http://dx.doi.org/10.1007/s00134-016-4245-3>.
 39. Goligher EC, Laghi F, Detsky ME, Farias P, Murray A, Brace D, et al. Measuring diaphragm thickness with ultrasound in mechanically ventilated patients: feasibility, reproducibility and validity. *Intensive Care Med*. 2015;41:642–9, <http://dx.doi.org/10.1007/s00134-015-3687-3>.