



ORIGINAL ARTICLE

Comparison of severity score models based on different sepsis definitions to predict in-hospital mortality among sepsis patients in the Intensive Care Unit



T. Songsangjinda^a, B. Khwannimit^{b,*}

^a Department of Internal Medicine, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand

^b Division of Critical Care Medicine, Department of Internal Medicine, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand

Received 6 September 2018; accepted 2 December 2018

KEYWORDS

Septic shock;
Sequential organ failure assessment;
Predisposition, insult, response, organ dysfunction;
Systemic inflammatory response syndrome;
Outcome

Abstract

Objective: A comparison is made of the accuracy between severity models, based on different sepsis definitions (systemic inflammatory response syndrome (SIRS), predisposition, insult, response, organ dysfunction (PIRO), and sequential organ failure assessment (SOFA) concepts), in predicting outcomes among sepsis patients.

Design: A retrospective study was carried out.

Setting: The study was conducted in the Intensive Care Unit (ICU) of a university teaching hospital.

Patients: Septic patients admitted to the ICU during 2007–2016.

Main variables of interest: The primary outcome was in-hospital mortality, with ICU mortality being the secondary outcome.

Results: A total of 2152 septic patient were identified, with ICU and in-hospital mortality rates of 33.3% and 45.9%, respectively. The Moreno PIRO (AUC, 95%CI) (0.835; 0.818–0.852) showed the highest discriminating capacity, followed by SOFA (0.828; 0.811–0.846), qSOFA (0.792; 0.775–0.809), Rubulotta PIRO (0.708; 0.687–0.730), Howell PIRO (0.706; 0.685–0.728) and SIRS (0.578; 0.556–0.600). The AUC of the SOFA score was comparable to that of the Moreno PIRO ($p=0.43$), though the AUCs of both of these scores were significantly higher than those of the other scores ($p < 0.001$ for all other comparisons). However, the SOFA score showed the best discriminating capacity in predicting ICU mortality (0.838; 0.820–0.855), followed by Moreno PIRO (0.804; 0.785–0.823) and qSOFA (0.787; 0.770–0.805). The accuracy of the qSOFA in predicting ICU mortality was comparable to that of the Moreno PIRO score ($p=0.15$).

Conclusion: The SOFA score and Moreno PIRO score showed the best accuracy in predicting in-hospital mortality among septic patients admitted to the ICU.

© 2018 Elsevier España, S.L.U. y SEMICYUC. All rights reserved.

* Corresponding author.

E-mail address: kbordin@medicine.psu.ac.th (B. Khwannimit).

PALABRAS CLAVE

Shock séptico;
Evaluación del fallo
orgánico secuencial;
Predisposición,
lesión, respuesta,
disfunción orgánica;
Síndrome de
respuesta
inflamatoria
sistémica;
Resultado

Comparación de los modelos de puntuación de la intensidad basados en diferentes definiciones de la sepsis en cuanto a la predicción de la mortalidad intrahospitalaria de los pacientes con sepsis ingresados en la Unidad de Cuidados Intensivos

Resumen

Objetivo: Comparar la precisión entre varios modelos de intensidad, basándose en diferentes definiciones de la sepsis (síndrome de respuesta inflamatoria sistémica [SIRS, por sus siglas en inglés], predisposición, infección, respuesta, disfunción orgánica [PIRO por sus siglas en inglés] y puntuación de la evaluación del fallo orgánico secuencial [SOFA por sus siglas en inglés]) para predecir los desenlaces en los pacientes con sepsis.

Diseño: Estudio retrospectivo.

Ámbito: El estudio se llevó a cabo en la unidad de cuidados intensivos (UCI) de un hospital universitario.

Pacientes: Enfermos con sepsis ingresados en la UCI durante 2007-2016.

Variables de interés principales: El desenlace principal fue la mortalidad hospitalaria, mientras que la mortalidad en la UCI fue el desenlace secundario.

Resultados: Se identificó un total de 2.152 pacientes con sepsis, con unas tasas de mortalidad en la UCI e intrahospitalaria del 33,3 y del 45,9%, respectivamente. El modelo Moreno-PIRO (AUC, IC del 95%) (0,835; 0,818-0,852) fue el que presentó una mayor capacidad de discriminación, seguido del modelo SOFA (0,828; 0,811-0,846), modelo qSOFA (0,792; 0,775-0,809), modelo Rubulotta-PIRO (0,708; 0,687-0,730), modelo Howell-PIRO (0,706; 0,685-0,728) y modelo SIRS (0,578; 0,556-0,600). El AUC de la puntuación SOFA fue comparable al de Moreno-PIRO ($p = 0,43$), si bien el AUC de ambas puntuaciones fue significativamente superior al de otras puntuaciones ($p < 0,001$ para todas las demás comparaciones). Sin embargo, la puntuación SOFA es la que presenta la mayor capacidad de discriminación para predecir la mortalidad en la UCI (0,838; 0,820-0,855), seguida de Moreno-PIRO (0,804; 0,785-0,823) y qSOFA (0,787; 0,770-0,805). La precisión de la puntuación qSOFA en cuanto a la predicción de la mortalidad en la UCI fue comparable a la de la puntuación Moreno-PIRO ($p = 0,15$).

Conclusión La puntuación SOFA y la puntuación Moreno-PIRO mostraron la mejor precisión en la predicción de la mortalidad intrahospitalaria entre pacientes sépticos ingresados en la UCI.

© 2018 Elsevier España, S.L.U. y SEMICYUC. Todos los derechos reservados.

Introduction

Sepsis and septic shock are currently a leading problem among intensive care unit (ICU) patients worldwide.¹⁻⁵ The ICU as well as in-hospital mortality owing to sepsis and septic shock have also been significantly higher than other critical illness conditions over the past few decades.¹⁻⁵ Until now, there have been three sepsis definitions endorsed by international societies, according to update knowledge in sepsis pathogenesis and pathophysiology. The first sepsis definition (Sepsis-1) was established in 1992, and focused on sepsis as a syndrome of infection, with systemic inflammatory response syndrome (SIRS concept).⁶ The second definition of sepsis by International Sepsis Definitions in 2001 (Sepsis-2) was also based on the SIRS concept, but tried to revise and included a list of more signs and symptoms, more so than SIRS.⁷ Signs and symptoms of sepsis according to Sepsis-2 definition was also classified into four categories which were: Predisposition, Insult/Infection, Response, Organ dysfunction; this in turn introduced the PIRO concept as a staging system of sepsis.⁷ However, the severity scoring system of sepsis was not unified, and so some authors tried to develop an individual score to predict mortality, based on the PIRO concept for both general sepsis

patients,⁸⁻¹⁴ and a more specific group such as; hospital acquired pneumonia.^{15,16} Unfortunately, the PIRO scores are not widely used nowadays owing to their complexity.

The sepsis definition was change again in 2016, by the Third International Consensus Definitions for Sepsis and Septic shock (Sepsis-3), the changed sepsis into an organ dysfunction syndrome, rather than SIRS, as the organ failure concept is better in mortality prediction.¹⁷ This definition reintroduces Sequential Organ Failure Assessment (SOFA)¹⁸ score into clinical use. The more simplified score or quick SOFA (qSOFA) was introduced for clinician alert for sepsis.¹⁷ The later score was validated by Seymour et al. and reported better accuracy of discrimination in term of mortality other than SIRS, especially in non-ICU encounter situations.¹⁹ The external validity was also conducted by other authors, who also reported better performance.²⁰⁻²⁴

Proof of accuracy, between the different sepsis concepts on the same population, may be useful for further development of a unified sepsis score. Thus, our study aimed to compare the accuracy between severity scores from 3 different definition concepts (SIRS, PIRO and SOFA-based scores), in term of in-hospital mortality prediction among sepsis patients admitted to an intensive care unit.

Methods

We conducted a retrospective cohort study analyzing data over a 10-year period (from January 1, 2007 to December 31, 2016), using electronic records along with the sepsis and severity scores database of a single medical ICU of a teaching hospital of Prince of Songkla University, Thailand. The protocol was approved by the institutional ethics committee on January 7, 2014 (Approval letter 57-366-14-4).

Population

Patients 18 years of age or older, who was admitted to our medical ICU with a diagnosis of severe sepsis and septic shock were enrolled regardless of the infection source or acquisition type. We defined sepsis terminology following the criteria of Sepsis-2 definition, according to the starting study period. Patients whose diagnosis was only sepsis using the terminology in Sepsis-2 definition without organ dysfunction were excluded. Organ failure was defined by an increase in the SOFA score of at least 2 points in each organ system. Patients for whom the length of stay was less than 4 h were also excluded, because it was too early to evaluate for change in either physiologic response or organ dysfunction.

Data Collection

Data for all components of SIRS,⁶ Moreno PIRO,⁸ Rubulotta PIRO,⁹ Howell PIRO,¹⁰ and qSOFA¹⁹ (detail in Tables S1–S6), as described by the original articles, were collected from the electronic database and Hospital Information System (HIS) of our hospital. Database of the studying ICU was manually collected by a well-trained research assistant in our unit.^{3,25} Parameters of patient predisposition (underlying diseases, age, gender, chronic illness *etc.*) were based on the admission data. Severity of illness was defined using Acute Physiology and Chronic Health Evaluation II (APACHE-II) score.²⁶ APACHE II and SOFA scores were derived from sepsis and severity scores database. Parameters of infection (infection source and acquisition type) were based on data from a final summary of hospital admission. Worst values of parameters within the first 24 h of ICU admission, related to physiological change and laboratory data were used to calculate the scores. The primary outcome was all causes of in-hospital mortality. The secondary outcome was ICU mortality. All parameters were electronically recorded using Epidata 3.1.

Statistical analysis

Baseline characteristic data were analyzed and represented with mean \pm SD, or median with interquartile range (IQR), according to data type and distribution. Student *T*-test was used for normal distribution data, and Wilcoxon's rank sum test for non-normal distribution data. We used the area under receiver operating characteristic curve (AUC) to determine discrimination among scores. Discrimination is an ability of the score to distinguish between survivors and non-survivors. Pair-wise comparison between the AUC




Figure 1 Enrollment and exclusion of study population.

of each score using method by the DeLong et al.,²⁷ and used Bonferroni correction for multiple comparisons of the AUC. Stata version 11 was used for all of statistical analysis. A *p*-value of less than 0.05 is accepted as a statistical significance. Serum lactate measurement, which is a component in Howell PIRO score,¹⁰ were available in our center only after 2011 owing to resource limitation before that period, causing 42.4% of missing data in serum lactate in our database. We also decided to perform sensitivity analysis for Howell PIRO score using the subgroup from the database collected after 2011.

Results

There were 6986 patients in the database during the 10-year study period in our ICU. A total of 2234 patients had been diagnosed with severe sepsis and septic shock, using the Sepsis-2 criteria. After exclusion of 82 entries, which were diagnosed as sepsis without organ failure, 2152 patients were included into the primary analysis (Fig. 1). The mean APACHE-II score was 23.9 ± 9.8 . The ICU and hospital mortality was 33.3% and 45.9%, respectively. The baseline characteristics of the study population are presented in Table 1.

Mortality tended to be higher according to the score increasing, in all scores, except SIRS (Figures S1–S3). The Moreno PIRO score had the highest AUC for in-hospital mortality (0.835, 95%CI 0.818–0.852), followed by SOFA score (0.828, 95%CI 0.811–0.846), qSOFA (0.792, 95%CI 0.775–0.809), Rubulotta PIRO score (0.708, 95%CI 0.687–0.730), Howell PIRO score (0.706, 95%CI 0.685–0.728), and SIRS (0.578, 95%CI 0.556–0.600). However, the AUC of SOFA score was not statistically different with Moreno PIRO score (*p* = 0.43). The AUC of SOFA and Moreno PIRO scores were significantly higher than that of other scores (Table 2 and Fig. 2). Nevertheless, the APACHE II score provide the higher AUC (0.883, 95%CI 0.868–0.897) than SOFA and other PIRO scores.

Table 1 Patient characteristics data.

Variables	All patients (n=2152)	Survivors (n=1164)	Non-survivors (n=988)	p-Value
Age, year (Median, IQR)	61 (45–74)	58 (44–74)	62 (46–75)	0.11
Male (%)	1216 (56.5)	686 (56.4)	570 (46.9)	0.29
Presence of at least one chronic illness (%)	901 (41.9)	376 (32.3)	525 (53.1)	<0.001
Hematologic malignancy (%)	239 (11.1)	74 (6.3)	165 (16.7)	<0.001
Immunosuppressive agent using (%)	166 (7.7)	86 (7.4)	80 (8.1)	0.54
Liver cirrhosis (%)	110 (5.1)	31 (2.7)	79 (8.0)	<0.001
Septic shock (%)	1775 (82.5)	853 (73.3)	922 (93.3)	<0.001
Positive blood culture	652 (30.3)	325 (27.9)	327 (33.1)	0.94
Community acquired infection (%)	1423 (66.1)	907 (77.9)	516 (52.2)	<0.001
Source of infection				
Respiratory tract infection (%)	1067 (49.9)	541 (46.5)	526 (53.2)	0.65
Gastrointestinal tract infection (%)	303 (14.2)	161 (13.8)	142 (14.4)	0.72
Urinary tract infection (%)	191 (8.9)	137 (11.8)	54 (5.5)	<0.001
ICU length of stay, day (Median, IQR)	4 (2–8)	4 (2–8)	4 (1–9)	0.95
Scores				
SIRS	3.3 ± 0.82	3.2 ± 0.86	3.4 ± 0.73	<0.001
Moreno PIRO	34 (25–43)	27 (21–34)	42 (35–49)	<0.001
Rubulotta PIRO	6 (4–7)	5 (4–6)	7 (5–8)	<0.001
Howell PIRO	14 (12–17)	14 (11–16)	16 (14–18)	<0.001
SOFA	9 (6–12)	6 (4–9)	12 (9–14)	<0.001
qSOFA	2.4 ± 0.66	2.0 ± 0.61	3.8 ± 0.19	<0.001

PIRO: Predisposition, Insult/Infection, Response, Organ dysfunction; qSOFA: quick sequential organ failure assessment; SIRS: systemic inflammatory response syndrome; SOFA: sequential organ failure assessment.

Table 2 Area under receiving operating characteristic (AUC) and p-value comparison between sepsis based-scores for in-hospital mortality.

Scores (AUC, 95%CI)	SIRS	Moreno	Rubulotta	Howell	SOFA
SIRS (0.578, 0.556–0.600)					
Moreno PIRO (0.835, 0.818–0.852)	<0.001				
Rubulotta PIRO (0.708, 0.687–0.730)	<0.001	<0.001			
Howell PIRO (0.706, 0.685–0.728)	<0.001	<0.001	0.74		
SOFA (0.828, 0.811–0.846)	<0.001	0.43	<0.001	<0.001	
qSOFA (0.792, 0.775–0.809)	<0.001	<0.001	<0.001	<0.001	<0.001

PIRO: Predisposition, Insult/Infection, Response, Organ dysfunction; qSOFA: quick sequential organ failure assessment; SIRS: systemic inflammatory response syndrome; SOFA: sequential organ failure assessment.




Figure 2 Receiving operating characteristic curves (ROC) of severity scores for predict in-hospital mortality.

The SOFA score present the best discrimination with an AUC with 95%CI of 0.838, 0.820–0.855 to predict ICU mortality. There were no differences of AUCs between Moreno PIRO score and qSOFA (AUC with 95%CI 0.804, 0.785–0.823 and 0.787, 0.770–0.805, respectively), but their performance were statistically higher than other scores to predict ICU mortality (Table S7). The AUCs of each score to predict the ICU mortality are shown in Figure S4 and Table S7.

The discrimination of each score for subgroups analysis is shown in Table 3. There was similar predictive ability of SOFA and Moreno PIRO to predict hospital mortality in all subgroup of sepsis patients, as well as, the AUCs of both scores were statistically higher than SIRS and other PIRO scores to predict hospital mortality in all subgroup of our sepsis patients.

Table 3 Comparison of the area under receiver operating characteristic curve (AUC) for sepsis based-scores for predict in-hospital mortality.

Subgroup of patients	SIRS	Moreno PIRO	Rubulotta PIRO	Howell PIRO	SOFA	qSOFA
Severe sepsis (n = 377)	0.586 (0.516–0.656)	0.769 (0.702–0.836)	0.689 (0.612–0.768)	0.646 (0.569–0.724)	0.769 (0.695–0.840)	0.713 (0.647–0.779)
Septic shock (n = 1775)	0.558 (0.533–0.582)	0.821 (0.803–0.840)	0.678 (0.654–0.703)	0.690 (0.666–0.714)	0.820 (0.800–0.839)	0.782 (0.763–0.802)
Respiratory tract infection (n = 1076)	0.580 (0.548–0.611)	0.835 (0.811–0.858)	0.712 (0.681–0.742)	0.704 (0.674–0.735)	0.847 (0.824–0.870)	0.781 (0.756–0.806)
Gastrointestinal tract infection (n = 303)	0.551 (0.490–0.611)	0.876 (0.837–0.914)	0.714 (0.657–0.771)	0.726 (0.670–0.782)	0.848 (0.805–0.891)	0.819 (0.775–0.864)
Year 2007–2011 (n = 1077)	0.561 (0.530–0.592)	0.805 (0.779–0.830)	0.712 (0.682–0.742)	0.695 (0.664–0.726)	0.799 (0.752–0.816)	0.737 (0.710–0.764)
Year 2012–2016 (n = 1075)	0.592 (0.560–0.623)	0.868 (0.847–0.889)	0.705 (0.674–0.735)	0.720 (0.690–0.751)	0.882 (0.862–0.902)	0.845 (0.822–0.865)

PIRO: Predisposition, Insult/Infection, Response, Organ dysfunction; qSOFA: quick sequential organ failure assessment; SIRS: systemic inflammatory response syndrome; SOFA: sequential organ failure assessment.

However, there were only 1239 patients having had their serum lactate values, so as to be included into subgroup analysis. In subgroup analysis, the AUC of Howell PIRO score was higher than the whole population (0.724 vs. 0.706). However, the performance of Howell PIRO to predict primary outcome was lower than Moreno PIRO and SOFA score and similarly to Rubulotta PIRO, as same as the entire population (Table S8).

Discussion

To the best of our knowledge, this is the first comparison of severity scores derived from all sepsis definition concepts. We found both the SOFA and Moreno PIRO score had the highest mortality discrimination, additionally they were also higher than the value reported in original articles of both scores (AUC of SOFA and Moreno PIRO score in original article was 0.74¹⁹ and 0.772,⁸ respectively). This finding may be owing to the higher mortality in our center, compared with worldwide epidemiologic data.² Although, we hypothesized that both SOFA and Moreno PIRO scores still preserve their good performance, even in a more severe sepsis patients. This statement may not be true for other scores, as their AUC did not result in as good results, as in their original articles.^{9,10,19} Actually, components of organ dysfunction in the Moreno PIRO score⁸ was derived from the original SOFA, hence, may explain why the Moreno PIRO score is better than other PIRO scores. Even though these results may not directly conclude that the organ failure concept is better than other concepts for sepsis, we found that scores that, focus around organ failure seem to have better mortality discrimination than other scores.

In this study, we aimed to compare between different sepsis definitions. Thus, we excluded patients without organ dysfunction, in order to unify the study population into the

same terminology, which preserves severe sepsis in Sepsis-1 and Sepsis-2 definition is actually the same as sepsis in Sepsis-3 definition. Fortunately, sepsis patients without organ dysfunction were the minority our study population (only 3.6%), and may not have affected the primary outcome.

Previous studies from other investigators had conflicted results. MacDonald et al. compared Howell PIRO, SOFA, and Mortality in ED Sepsis (MEDS) in 240 sepsis patients, who visited an emergency department, and found that the Howell PIRO score performed better than the SOFA score, and similarly to the MEDS score to predict 30-day mortality.²⁸ We hypothesized that the SOFA score may have better performance in an ICU setting, rather than in an emergency department setting, which is too early to detect changes in organ functions during the diseases course. Thus, the PIRO concept that includes predisposition factors in the score should be used in this setting. The study also included infections with no organ dysfunction (18% of study population), which is not considered to be sepsis nowadays. Another study from de Groot et al. also found that the performance of Howell PIRO, qSOFA, MEDS, Modified Early Warning Score (MEWS), National Early Warning Score (NEWS) in an emergency department setting were poor and the AUC ranged from 0.56 to 0.64 in older patients, significantly lower than the good AUC range from 0.72 to 0.86 in younger patients (lower than 70 years). The qSOFA had a lower AUC than that of the Howell PIRO score.²⁹ Although some of the study populations in this study were within the older age group, most of them were younger than 70 years of age. The difference in both the study population and setting may explain the different results. The study of de Groot et al. also reported lower mortality (9.5%)²⁹ compare with ours, and qSOFA discrimination may not as good as our results. Tafelski et al. compared the accuracy between three PIRO scores, based on a model of the critically-ill in an ICU, and they found

that Moreno PIRO and Howell PIRO were statistically different compared with Rubulotta PIRO.³⁰ Most of the patients in this study were in the organ dysfunction group, much like our study, but hospital mortality was 16.2% lower than our results. Different PIRO scores may have different accuracy dependant on population settings and sepsis severity.

Our findings also support that among the more severe sepsis patients, use of SOFA score, which focus on organ dysfunction has more advantage in mortality prediction. Moreover, SOFA score is more simplified and easier to use than Moreno PIRO score, even though all of them have a comparable discrimination performance. We support the use of the SOFA score more so, than the Moreno PIRO score for risk prediction in ICU sepsis patients.

The limitation of our study was that it only included sepsis patients admitted into a medical ICU of a university teaching hospital; these results may some limitation in application in other sepsis patients in other ICU or sepsis in wards or emergency department. Secondly, our results had a higher prevalence of septic shock and high mortality rate; these findings may not be generalized to less severely-sepsis patients. More studies are required to clarify prognostic determination among these different phenotypes of sepsis patients. Thirdly, the accuracy of these scores for predicting mortality in sepsis patients diagnosed according to Sepsis-2 criteria may not be the same as those diagnosed using the new Sepsis-3.

Conclusion

Severity scores derived from the Sepsis-3 concept that consists of organ dysfunction and organ failure assessment have better discrimination performance than other concepts. Our findings support SOFA score for risk prediction in medical-based ICU sepsis patients.

Authors contribution

The study concept and design was created by all authors. TS and BK collected data and performed the statistical analysis. TS drafted the manuscript. BK has reviewed and revised manuscript. All authors read and approved the final manuscript.

Conflict of interest

None.

Acknowledgements

This study was supported by a research grant from the Faculty of Medicine, Prince of Songkla University. The authors would like to acknowledge Mr. Andrew Tait, of the International Affairs Department, Faculty of Medicine, Prince of Songkla University for his help in editing the manuscript.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.medin.2018.12.004](https://doi.org/10.1016/j.medin.2018.12.004).

References

1. Vincent JL, Marshall JC, Namendys-Silva SA, Francois B, Martin-Loeches I, Lipman J, et al. Assessment of the worldwide burden of critical illness: the Intensive Care Over Nations (ICON) audit. *Lancet Respir Med*. 2014;2:380–6.
2. Vincent JL, Rello J, Marshall J, Silva E, Anzueto A, Martin CD, et al. International study of the prevalence and outcomes of infection in intensive care units. *JAMA*. 2009;302:2323–9.
3. Khwannimit B, Bhurayanontachai R. The epidemiology of, and risk factors for, mortality from severe sepsis and septic shock in a tertiary-care university hospital setting. *Epidemiol Infect*. 2009;137:1333–41.
4. Vincent JL, Sakr Y, Sprung CL, Ranieri VM, Reinhart K, Gerlach H, et al. Sepsis in European intensive care units: results of the SOAP study. *Crit Care Med*. 2006;34:344–53.
5. Vincent JL. EPIC II: sepsis around the world. *Minerva Anesthesiol*. 2008;74:293–6.
6. Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest*. 1992;101:1644–55.
7. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med*. 2003;31:1250–6.
8. Moreno RP, Metnitz B, Adler L, Hoechtl A, Bauer P, Metnitz PG, et al. Sepsis mortality prediction based on predisposition, infection and response. *Intensive Care Med*. 2008;34:496–504.
9. Rubulotta F, Marshall JC, Ramsay G, Nelson D, Levy M, Williams M. Predisposition, insult/infection, response, and organ dysfunction: a new model for staging severe sepsis. *Crit Care Med*. 2009;37:1329–35.
10. Howell MD, Talmor D, Schuetz P, Hunziker S, Jones AE, Shapiro NI. Proof of principle: the predisposition, infection, response, organ failure sepsis staging system. *Crit Care Med*. 2011;39:322–7.
11. Cardoso T, Teixeira-Pinto A, Rodrigues PP, Aragao I, Costa-Pereira A, Sarmiento AE. Predisposition, insult/infection, response and organ dysfunction (PIRO): a pilot clinical staging system for hospital mortality in patients with infection. *PLoS ONE*. 2013;8:e70806.
12. Chen YX, Li CS. Evaluation of community-acquired sepsis by PIRO system in the emergency department. *Intern Emerg Med*. 2013;8:521–7.
13. Granja C, Povoas P, Lobo C, Teixeira-Pinto A, Carneiro A, Costa-Pereira A. The predisposition, infection, response and organ failure (Piro) sepsis classification system: results of hospital mortality using a novel concept and methodological approach. *PLoS ONE*. 2013;8:e53885.
14. Yeh CF, Chen KF, Ye JJ, Huang CT. Derivation of a clinical prediction rule for bloodstream infection mortality of patients visiting the emergency department based on predisposition, infection, response, and organ dysfunction concept. *J Microbiol Immunol Infect*. 2014;47:469–77.
15. Lisboa T, Diaz E, Sa-Borges M, Socias A, Sole-Violan J, Rodriguez A, et al. The ventilator-associated pneumonia PIRO score: a tool for predicting ICU mortality and health-care resources use in ventilator-associated pneumonia. *Chest*. 2008;134:1208–16.

16. Rello J, Rodriguez A, Lisboa T, Gallego M, Lujan M, Wunderink R. PIR0 score for community-acquired pneumonia: a new prediction rule for assessment of severity in intensive care unit patients with community-acquired pneumonia. *Crit Care Med*. 2009;37:456–62.
17. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016;315:801–10.
18. Vincent JL, Moreno R, Takala J, Willatts S, De Mendonca A, Bruining H, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med*. 1996;22:707–10.
19. Seymour CW, Liu VX, Iwashyna TJ, Brunkhorst FM, Rea TD, Scherag A, et al. Assessment of clinical criteria for sepsis: for the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016;315:762–74.
20. Forward E, Konecny P, Burston J, Adhikari S, Doolan H, Jensen T. Predictive validity of the qSOFA criteria for sepsis in non-ICU inpatients. *Intensive Care Med*. 2017;43:945–6.
21. Guirgis FW, Puskarich MA, Smotherman C, Sterling SA, Gautam S, Moore FA, et al. Development of a simple sequential organ failure assessment score for risk assessment of emergency department patients with sepsis. *J Intensive Care Med*. 2017, 885066617741284.
22. Ho KM, Lan NS. Combining quick Sequential Organ Failure Assessment with plasma lactate concentration is comparable to standard Sequential Organ Failure Assessment score in predicting mortality of patients with and without suspected infection. *J Crit Care*. 2017;38:1–5.
23. Raith EP, Udy AA, Bailey M, McGloughlin S, MacIsaac C, Bellomo R, et al. Prognostic accuracy of the SOFA score, SIRS criteria, and qSOFA score for in-hospital mortality among adults with suspected infection admitted to the Intensive Care Unit. *JAMA*. 2017;317:290–300.
24. Khwannimit B, Bhurayanontachai R, Vattanavanit V. Comparison of the performance of SOFA, qSOFA and SIRS for predicting mortality and organ failure among sepsis patients admitted to the intensive care unit in a middle-income country. *J Crit Care*. 2017;44:156–60.
25. Khwannimit B, Bhurayanontachai R. Validation of predisposition, infection, response and organ dysfunction score compared with standard severity scores in predicting hospital outcome in septic shock patients. *Minerva Anesthesiol*. 2013;79:257–63.
26. Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. *Crit Care Med*. 1985;13:818–29.
27. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics*. 1988;44:837–45.
28. Macdonald SP, Arendts G, Fatovich DM, Brown SG. Comparison of PIRO, SOFA, and MEDS scores for predicting mortality in emergency department patients with severe sepsis and septic shock. *Acad Emerg Med*. 2014;21:1257–63.
29. de Groot B, Stolwijk F, Warmerdam M, Lucke JA, Singh GK, Abbas M, et al. The most commonly used disease severity scores are inappropriate for risk stratification of older emergency department sepsis patients: an observational multi-centre study. *Scand J Trauma Resusc Emerg Med*. 2017;25:91.
30. Tafelski S, Nachtigall I, Stengel S, Wernecke K, Spies C. Comparison of three models for sepsis patient discrimination according to PIRO: predisposition, infection, response and organ dysfunction. *Minerva Anesthesiol*. 2015;81:264–71.