

- Intensive and Critical Care Medicine and Coronary Units (SEMICYUC), the Spanish Society of Otorhinolaryngology and Head and Neck Surgery (SEORL-CCC) and the Spanish Society of Anesthesiology and Resuscitation (SEDAR) on tracheotomy in patients with COVID-19 infection. *Med Intensiva*. 2020;S0210-S5691, <http://dx.doi.org/10.1016/j.medin.2020.05.002>.
2. Foster P, Cheung T, Craft P, Baran K, Kryskow M, Knowles R, et al. Novel approach to reduce transmission of COVID-19 during tracheostomy. *J Am Coll Surg*. 2020. April 10;S1072-7515(20)30322-30327. doi:<https://doi.org/10.1016/j.jamcollsurg.2020.04.014>. Epub ahead of print.
  3. Shidham VB, Frisch NK, Layfield LJ. Severe acute respiratory syndrome coronavirus 2 (the cause of COVID 19) in different types of clinical specimens and implications for cytopathology specimen: an editorial review with recommendations. *Cytojournal*. 2020;17:7, doi:[/Cytojournal\\_24\\_2020](https://doi.org/10.1016/j.cytojournal.2020.04.014).
  4. Zuo MZ, Huang YG, Ma WH, Xue ZG, Zhang JQ, Gong YH, et al. Expert recommendations for tracheal intubation in critically ill patients with novel coronavirus disease 2019. *Chin Med Sci J*. 2020. February 27. doi: [10.24920/003724](https://doi.org/10.24920/003724). Epub ahead of print.
  5. Canelli R, Connor CW, Gonzalez M, Nozari A, Ortega R. Barrier enclosure during endotracheal intubation. *N Engl J Med*. 2020;382:1957-8, doi: <https://doi.org/10.1056/NEJMc2007589>
- G. Pérez Acosta, D. González Romero, L. Santana-Cabrera\*
- Servicio de Medicina Intensiva del Complejo Hospitalario Universitario Insular Materno Infantil de Las Palmas de Gran Canaria, Spain*
- \*Corresponding author.  
E-mail address: [lsancabx@gobiernodecanarias.org](mailto:lsancabx@gobiernodecanarias.org) (L. Santana-Cabrera).
- 2173-5727/ © 2020 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.

## Reply to “Consensus document on tracheotomy in patients with COVID-19”<sup>☆</sup>



### Respuesta a «Documento de consenso de la traqueotomía en pacientes con COVID-19»

Dear Editor:

We wish to thank the warm welcome to “Consensus document on tracheostomy in patients with COVID-19”<sup>1</sup> and congratulate the authors on their quest for tools to reduce the risk of health professionals while managing these patients.

We agree that we should be extra-cautious with patients infected with SARS-CoV-2, especially while performing high-risk procedures. Specific recommendations on the use of personal protection equipment (PPE) for every particular clinical situation have been established. The risk of contagion involved in procedures like intubation, tracheostomy or bronchoscopy and the reduced availability of PPE has triggered the design of devices that act as barriers to reduce the risk of contagion.

The so-called “aerosol box” was initially designed to intubate patients with COVID-19 in an attempt to reduce the spread of unwanted aerosols.<sup>2</sup> Its use has been controversial ever since with limitations like complexity in cases of difficult airways, limited movements for operators and technicians, difficulties placing or repositioning the device in emergency cases (even damaging the patient), reduced visibility, possibility of cross contamination if the device has not been properly disinfected or risk of contagion during its retrieval.<sup>3</sup> Other barrier devices like plastic screens have

been used with greater maneuverability and visualization or suction systems to reduce the viral load and spread of aerosols.

From these experiences different prototypes of devices have been developed. Their ergonomics, shape, size, and type of material have improved. Their initial design has changed and adapted to the actual clinical needs. Also, some of the limitations described above have been solved. The use of more flexible curve-shaped materials has improved visibility and avoided cleaning issues. New access ports have been added that technicians and assistants can use or sealing machines have been introduced to reduce the risk of viral particle spread or systems to increase the stability of the devices.<sup>4</sup>

The use of all these barriers has extended to procedures like tracheostomy (as the authors say) and other experiences.<sup>5</sup> Actually, they improve the safety of one of the most risky and common procedures often performed during these months in patients with COVID-19.

Despite the continuous improvement in the design of these devices, the apparent greater safety and confidence of health professionals should not minimize the precautions that still need to be observed in the use of PPE and other standard protection measures. Therefore, health professionals should be properly trained in the use of these devices. Simulation can be an excellent tool to validate these devices during the training process of health professionals.

This necessary innovation in times of high tension and healthcare system collapse should consolidate in the near future to make sure that these medical devices are regulated, manufactured, assessed, and used according to common quality and safety standards and without new risks. To that end, multidisciplinary teams should provide their clinical experience and technological know-how to the design and manufacture processes of these devices.

Their implementation into the routine clinical practice should be accompanied by clinical results that prove their effectiveness reducing the rates of infection or the quantitative viral loads in PPE and in the environment without

<sup>☆</sup> Please cite this article as: Martín Delgado MC, Bernal-Sprekelsen M, Villalonga Vadel R. Respuesta a «Documento de consenso de la traqueotomía en pacientes con COVID-19». *Med Intensiva*. 2021;45:254-255.

compromising the safety of the procedures for which they were designed in the first place.

## Funding

The authors confirm that no funding was received while conducting this study.

## References

- Martín Delgado MC, Avilés-Jurado FX, Álvarez Escudero J, Aldecoa Álvarez-Santuyano C, de Haro López C, Díaz de Cerio Canduela P, et al. Consensus document of the Spanish Society of Intensive and Critical Care Medicine and Coronary Units (SEMICYUC), the Spanish Society of Otorhinolaryngology and Head and Neck Surgery (SEORL-CCC) and the Spanish Society of Anesthesiology and Resuscitation (SEDAR) on tracheotomy in patients with COVID-19 infection. *Med Intensiva*. 2020, <http://dx.doi.org/10.1016/j.medin.2020.05.002>.
- Canelli R, Connor CW, Gonzalez M, Nozari A, Ortega R. Barrier enclosure during endotracheal intubation. *N Engl J Med*. 2020;382, <http://dx.doi.org/10.1056/NEJMc2007589>.
- Rosenblatt WH, Sherman JD. More on barrier enclosure during endotracheal intubation. *N Engl J Med*. 2020;382:e69, <http://dx.doi.org/10.1056/NEJMc2012960>.
- Bai JW, Ravi A, Notario L, Choi M. Opening the discussion on a closed intubation box. *Trends Anaesth Crit Care*. 2020, <http://dx.doi.org/10.1016/j.tacc.2020.06.004>.
- Alves Filho W, Pontes Grangeiro Teles TS, Studart da Fonseca MR, Farias Pereira Filho FJ, Monteiro Pereira G, Moura Pontes AB, et al. Barrier device prototype for open tracheotomy during COVID-19 pandemic. *Auris Nasus Larynx*. 2020;S0385–8146, <http://dx.doi.org/10.1016/j.anl.2020.05.003>, 30119-X.

M.C. Martín Delgado<sup>a,b,\*</sup>, M. Bernal-Sprekelsen<sup>c,d</sup>, R. Villalonga Vadell<sup>e,f</sup>

<sup>a</sup> *Servicio de Medicina Intensiva, Hospital Universitario de Torrejón, Torrejón de Ardoz, Madrid, Spain*

<sup>b</sup> *Universidad Francisco de Vitoria, Pozuelo de Alarcón, Madrid, Spain*

<sup>c</sup> *Vicepresidente de la SEORL-CCC, Madrid, Spain*

<sup>d</sup> *Servicio de ORL, Hospital Clínic, Barcelona, Spain*

<sup>e</sup> *Vicepresidenta de la Comisión Nacional de Anestesiología, Reanimación y Terapéutica del Dolor, Spain*

<sup>f</sup> *Presidenta de la Sociedad Catalana de Anestesiología, Reanimación y Terapéutica del Dolor (SCARTD), Spain*

\* Corresponding author.

E-mail address: [mcmartindelgado@gmail.com](mailto:mcmartindelgado@gmail.com) (M.C. Martín Delgado).

2173-5727/ © 2020 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.