

results, with the limitations already described, pave the way for actually working on recommendations and protocols to facilitate the decision-making process in this setting. These tools have proven capable of preventing up to 40% of all inappropriate ICU admissions among these patients.⁵ Also, we think collaborative work to develop advanced directives in this type of patients is of paramount importance. However, in other settings, after controlling malignancy these advanced directives did not anticipate the reason for ICU admission, disease severity, age, and decision to start life-support therapies in patients with and without such directives. However, the presence of an advanced directive may have helped guide prior decisions on the duration of therapy and state of resuscitation.⁶

In this context, back in 2017, SEOM and SEMICYUC signed an agreement to improve the quality of care of cancer patients with critical complications. This initiative aims at contributing to the decision-making process, standardizing criteria, reducing subjectivity, generating channels of communication, and going deeper into the ethical and scientific aspects of these situations. Cancer drugs, chemosensitive tumors, presence of mutations predictive of response to targeted therapies, possibility of long-term tumor control with immunotherapy, and other specific situations should make us be able to agree on what patients are eligible for ICU admissions based on their distinctive characteristics.⁷

Funding

No funding was received whatsoever.

References

1. Sociedad Española de Medicina Intensiva y Unidades Coronarias (SEMICYUC). Convenio de colaboración SEMI-

CYUC y SEOM. [consultado 5 Jul 2019]. Available at: <https://semicyuc.org/convenios-de-colaboracion/>.

2. Nilsson J, Blomberg C, Holgersson G, Carlsson T, Bergqvist M, Bergström S. End-of-life care: where do cancer patients want to die? A systematic review. *Asia Pac J Clin Oncol*. 2017;13:356–64.
3. Thiéry G, Azoulay E, Darmon M, Ciroidi M, De Miranda S, Lévy V, et al. Outcome of cancer patients considered for intensive care unit admission: a hospitalwide prospective study. *J Clin Oncol*. 2005;23:4406–13.
4. Díaz-Díaz D, Villanova Martínez M, Palencia Herrejón E. Oncological patients admitted to an intensive care unit. Analysis of predictors of in-hospital mortality. *Med Intensiva*. 2018;42:346–53.
5. Nasir S, Muthiah M, Ryder K, Clark K, Niell H, Weir A. ICU deaths in patients with advanced cancer: reasonable criteria to decrease potentially inappropriate admissions and lack of benefit of advance planning discussions. *Am J Hosp Palliat Care*. 2017;34:173–9.
6. Kish Wallace S, Martin CG, Shaw AD, Price KJ. Influence of an advance directive on the initiation of life support technology in critically ill cancer patients. *Crit Care Med*. 2001;29:2294–8.
7. Carmona-Bayonas A, Gordo F, Beato C, Castaño Pérez J, Jiménez-Fonseca P, Virizuela Echaburu J, et al. Intensive care in cancer patients in the age of immunotherapy and molecular therapies: commitment of the SEOM-SEMICYUC. *Med Intensiva*. 2018;42:363–9.

P. Escudero-Acha^a, Y. Peñasco^{a,b}, J.C. Rodríguez-Borregan^{a,b}, E. Cuenca Fito^a, L. Ferreira Freire^b, A. González-Castro^{a,*}

^a Servicio de Medicina Intensiva, Hospital Universitario Marqués de Valdecilla, Santander, Cantabria, Spain

^b Servicio de Oncología, Hospital Universitario Marqués de Valdecilla, Santander, Cantabria, Spain

Corresponding author.

E-mail address: e409@humv.es (A. González-Castro).

2173-5727/ © 2019 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.

Analysis of SARS-CoV-2 in the air of an ICU dedicated to covid-19 patients[☆]



Análisis de SARS-CoV-2 en el aire de una UCI dedicada a pacientes covid-19

Infection due to SARS-CoV-2 produces pneumonia and acute respiratory distress syndrome in the context of a disease known as COVID-19.¹ In only a few months, the virus spread throughout the world, infecting millions of people and causing hundreds of thousands of deaths.² Contagion takes place through respiratory droplets (>5 μm) and contact with con-

taminated objects (fomites).^{3,4} Recently, the World Health Organization (WHO) has acknowledged that airborne transmission through aerosols (<5 μm) that remain in the air for hours is also a possible contagion route, due to the inhalation of viral particles that are deposited within the distal airway.^{5–7} Airborne transmission is of great relevance to public health and the protection of healthcare professionals, and in this regard the measures of caution need to be modified to avoid contagion—incrementing the required 2 m safety distance, particularly in closed areas or spaces.

Intensive Care Units (ICUs) have periodic air renewal mechanisms, and during the pandemic, many Units were equipped with negative pressure systems. However, other places such as homes, restaurants, public transport or even hospital wards have no such safe ventilation systems. Some studies have reported that SARS-CoV-2 is able to remain in the air generated by aerosols for up to three hours,⁷ with demonstration of the presence of the viral genome in the air and filters of hospitals.

[☆] Please cite this article as: Escudero D, Barrera JA, Balboa S, Viñas S, Martín G, Boga JA. Análisis de SARS-CoV-2 en el aire de una UCI dedicada a pacientes covid-19. *Med Intensiva*. 2021;45:247–250.

Table 1 Respiratory support, days of admission, and microbiological characteristics of the patients with COVID-19 in boxes where the air was analyzed.

Patients	Respiratory support	Days of admission	PCR	IgM	IgG
Box 33	Nasal cannula 2l/min	63 days ICU	—	+	+
Box 37	Mechanical ventilation. Pressure assist	39 days ICU	+	+	+
Box 38	FiO ₂ 0.3 PEEP 7 Mechanical ventilation	46 days ICU	—	+	+
Box 50–2 measurements (plate/swab and filter)	FiO ₂ 0.6 PEEP 8 Oxygen mask. FiO ₂ 0.5	42 days ICU	—	+	+
Box 56	Mechanical ventilation FiO ₂ 0.4 PEEP 10	36 days ICU	—	+	+
Pneumology ward room	Nasal cannula 1l/min	One day	—	+	+

FiO₂: fraction of inspired oxygen; PCR: polymerase chain reaction; PEEP: positive end-expiratory pressure; ICU: Intensive Care Unit.

In Spain, the SARS-CoV-2 contagion rate among healthcare professionals is the highest in the world, followed by Italy (10%) and China (3,8%).⁸ According to the Spanish Ministry of Health, on 9 July 2020, the number of infected healthcare professionals totaled 52,643, representing over 22% of all cases in the country.⁹ Those working in the ICU are at a high risk of contagion due to the great environmental exposure to SARS-CoV-2 and the use of techniques that generate aerosols.^{5,10,11} Thus, one of our major concerns was to improve the safety measures and minimize the risk of contagion among the healthcare professionals. In this context, we decided to analyze the presence of SARS-CoV-2 in the air of two ICUs and in the pneumology ward dedicated to the treatment of patients with COVID-19. The study was carried out in late May 2020 in 5 different boxes of the ICU (Table 1), placing the extraction equipment on the floor, close to the head of the patient and as far as possible from the air outlet, collecting the air samples during a time of 2–4 h. The entire ICU was equipped with negative air pressure (–10 Pa), and the air was renewed at a rate of 15–20 cycles/h. The hospital ward had no such air renewal system. The air volume in the ICU boxes is variable, since their dimensions differ—though the average is 51 m³, i.e., 51,000 l, and the hospital ward rooms have a volume of 54.4 m³. The air samples were obtained using two different methods: (1) SAS Bioser Mod. Microbio 0111302 sampling equipment with an air flow of 500 l/300 s and a Rodac plate measuring 55 mm in diameter from which samples were subsequently obtained with pre-humidified swabs. With this system the estimated volume of air passing through the plate in one hour is 5,967 l; and (2) A filtration ramp with a polyethersulfone membrane filter (FILTER-LAB®) of pore size 0.1 μm and measuring 47 mm in diameter, connected to the hospital vacuum system by means of a 60 kPa vacuumeter. There is a small variability in vacuum of the different ICU boxes, ranging between 55–60 kPa; as a result, calculation of the air passing through the plate is only an estimate. Considering 55 kPa, tubing measuring 8 mm in diameter, and the different loading losses, the calculated sampled air volume is 2,160 l/h. Thus, with the studied time period of 2–4 h

Table 2 Air samples collected, obtainment method and SARS-CoV-2 PCR test results.

Air samples	SARS-CoV-2 PCR
- Box 33 ICU: Obtainment by plate/swab system	—
- Sampling time 2 h	
- Box 37 ICU: Obtainment by filter	—
- Sampling time 3 h	
- Box 38 ICU: Obtainment by filter	—
- Sampling time 2 h	
- Box 50 ICU:	—
- Obtainment by filter. Sampling time 4 h	
- Obtainment by plate/swab. Sampling time 2 h	
- Box 56 ICU: Obtainment by filter	—
- Sampling time 3 h	
- Pneumology ward room:	—
Obtainment by filter	
- Sampling time 3 h	

PCR: polymerase chain reaction; SARS-CoV-2: severe acute respiratory syndrome coronavirus-2; ICU: Intensive Care Unit.

and with the estimated flow rate, we analyzed 4,320 l in two hours and 8,640 in four hours—this in most cases representing 16.9% of the air in the ICU box.

The presence of SARS-CoV-2 was analyzed by detecting the viral genome using multiple quantitative RT-PCR. The nucleic acids were purified by the MagNa Pure 96 System (Roche, Geneva, Switzerland) from the swab transport medium or the eluate obtained after incubating the filters for half an hour at 37 °C. The extracts were subjected to an amplification reaction supplemented with a mixture of TaqMan® MGB primers and probes with two targets: the ORF1ab and N genes. The amplifications and their posterior analysis were performed using the Applied Biosystems® 7500 Real-time PCR System (ABI, Foster City, CA, USA).

A total of 7 air samples were obtained (Table 2). In no case did quantitative RT-PCR testing detect the SARS-CoV-2 genome in the samples obtained with the two described methods. In the present study we therefore could not demonstrate the presence of SARS-CoV-2 in the air of either the ICU or the hospital ward. These findings are consistent with those of authors such as Cheng et al.,¹² who did not detect coronavirus in 8 samples of air collected at 10 cm from the chin of a patient with COVID-19. However, other studies have detected viral RNA in between 35–68% of the air samples analyzed.^{13–15} In the study published by Guo et al.,¹⁴ the aerosols reached a distance of 4 m and proved positive in 40.6% of the air samples of the box in which the patient was located, and in 12.5% of the samples in the working areas. The ICU described by Guo et al. was equipped with air renewal at a rate of 16 cycles per hour, though no mention was made of whether the Unit was under negative pressure or not. Santarpia et al.¹⁵, in an isolation unit for patients in quarantine and in a hospitalization area of the University Medical Center of Nebraska, measured the presence of SARS-CoV-2 in the air of the room, and recorded a 63.2% positivity rate with a mean concentration of 2.86 copies/l of air. The air samples taken from the corridors in turn yielded a positivity rate of 66.7% with a mean concentration of 2.59 copies/l of air. Posteriorly, and in order to calculate infective capacity, Vero E6 cells were cultured with these samples, though the tests indicated no viral replication. The finding of SARS-CoV-2 RNA does not report on the presence of viable viruses and thus of their infective capacity, though contamination of the air would indicate that aerosol transmission of SARS-CoV-2 is plausible. The mentioned study did not specify air renewal (cycles per hour) or whether there was negative pressure in the rooms. In our ICU, all the boxes were equipped with negative air pressure (–10 Pa), and the air was renewed at a rate of 15–20 cycles/h, which could explain the absence of SARS-CoV-2 RNA in our study.

Following lockdown, a total of 237 ICU workers—including the cleaning staff—underwent PCR testing for SARS-CoV-2 and serological testing with the ELISA technique. All these tests proved negative.

With regard to the limitations of our study, mention must be made of the sample number, the fact that the screened volume of air represented only a part of the total volume, and the fact that air renewal and negative pressure in the ICU facilitate the elimination of SARS-CoV-2. Nevertheless, it is interesting to note that to our knowledge, this is the first study on the presence of coronavirus in the air of an ICU in Spain, and that few studies of this kind can be found in the literature.

The safety of healthcare professionals must be viewed as a priority concern in order to avoid collapse of the health systems and impede disease transmission from the hospitals to the rest of the community. In this regard it is particularly important to control the presence of SARS-CoV-2 on both the exposed environmental surfaces of the ICU and in the air, with a view to minimizing contagion risk, and to adopt strict safety measures during all maneuvers and treatments capable of generating aerosols in the ICU.

Authorship

Dolores Escudero designed the study and wrote the manuscript.

José Antonio Boga, Gabriel Martín and Juan A. Barrera participated in preparation and correction of the manuscript.

Salvador Balboa and Silvia Viñas performed collection of the samples and reviewed the manuscript.

Financial support

This study has received no financial support.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Acknowledgements

We wish to express our support of the patients who have suffered COVID-19 and our appreciation of their families for their courage in facing the disease under extreme conditions.

Thanks are due to Dr. Sergio Pérez-Holanda and to Mr. Josu Jiménez-Idoeta of Hospital Universitario Central de Asturias, for their permanent support in all the ICU projects.

References

- Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med.* 2020;382:727–33, <http://dx.doi.org/10.1056/NEJMoa2001017>.
- [Consulted 22 Jul 2010] Available in <https://coronavirus.jhu.edu/map.html>.
- Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med.* 2020;382:1199–207, <http://dx.doi.org/10.1056/NEJMoa2001316>.
- Peng X, Xu X, Li Y, Cheng L, Zhou X, Ren B. Transmission routes of 2019-nCoV and controls in dental practice. *Int J Oral Sci.* 2020;12:9, <http://dx.doi.org/10.1038/s41368-020-0075-9>.
- Wilson NM, Norton A, Young FP, Collins DW. Airborne transmission of severe acute respiratory syndrome coronavirus-2 to healthcare workers: a narrative review. *Anaesthesia.* 2020;75:1086–95, <http://dx.doi.org/10.1111/anae.15093>.
- Dancer SJ, Tang JW, Marr LC, Miller S, Morawska L, Jimenez JL. Putting a balance on the aerosolization debate around SARS-CoV-2. *J Hosp Infect.* 2020;105:569–70, <http://dx.doi.org/10.1016/j.jhin.2020.05.014>.
- Van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med.* 2020;382:1564–7, <http://dx.doi.org/10.1056/NEJMc2004973>.
- [Consulted 8 May 2010] Available in: <https://www.ecdc.europa.eu/sites/default/files/documents/covid-19-rapid-risk-assessment-coronavirus-disease-2019-ninth-update-23-april-2020.pdf>.
- [Consulted 8 May 2010] Available in: <https://www.rtve.es/noticias/20200710/profesionales-sanitarios-contagiados-covid-19-superan-50000/2014047.shtml>.

10. Ong SWX, Tan YK, Chia PY, Lee TH, Ng OT, Wong MSY, et al. Air, surface environmental, and personal protective equipment contamination by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) from a symptomatic patient. *JAMA*. 2020;326:1610–2, <http://dx.doi.org/10.1001/jama.2020.3227>.
11. Bahl P, Doolan C, de Silva C, Chughtai AA, Bourouiba L, MacIntyre CR. Airborne or droplet precautions for health workers treating COVID-19? *J Infect Dis*. 2020, <http://dx.doi.org/10.1093/infdis/jiaa189>, jiaa189.
12. Cheng VC, Wong SC, Chen JH, Yip CC, Chuang VW, Tsang OT, et al. Escalating infection control response to the rapidly evolving epidemiology of the Coronavirus disease 2019 (COVID-19) due to SARS-CoV-2 in Hong Kong. *Infect Control Hosp Epidemiol*. 2020;41:493–8.
13. Zhou J, Otter J, Price JR, Cimpeanu C, Garcia DM, Kinross J, et al. Investigating SARS-CoV-2 surface and air contamination in an acute healthcare setting during the peak of the COVID-19 pandemic in London. *Clin Infect Dis*. 2020, <http://dx.doi.org/10.1093/cid/ciaa905>, ciaa905.
14. Guo ZD, Wang ZY, Zang SF, Li X, Li L, Li C, et al. Aerosol and surface distribution of severe acute respiratory syndrome coronavirus 2 in hospital wards, Wuhan, China, 2020. *Emerg Infect Dis*. 2020;26, <http://dx.doi.org/10.3201/eid2607.200885>.
15. Santarpia JI, Rivera DN, Herrera V, Morwitzer MJ, Creager H, Santarpia GW, et al. Transmission potential of SARS-CoV-2 in viral shedding observed at the university of Nebraska medical center. *medRxiv*. 2020;3:20039446, <http://dx.doi.org/10.1101/2020.03.23.20039446>.

D. Escudero^{a,b,*}, J.A. Barrera^c, S. Balboa^{a,b}, S. Viñas^{a,b}, G. Martín^d, J.A. Boga^{b,d}

^a Servicio de Medicina Intensiva, Hospital Universitario Central de Asturias, Oviedo, Asturias, Spain

^b Grupo de Investigación Microbiología Traslacional del Instituto de Investigación Sanitaria del Principado de Asturias, Spain

^c Anaqua SL y Laboratorios Innoagral SL, Sevilla, Spain

^d Servicio de Microbiología, Hospital Universitario Central de Asturias, Oviedo, Asturias, Spain

Corresponding author.

E-mail address: lolaescudero@telefonica.net (D. Escudero).

2173-5727/ © 2020 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.

Survey of the neurocritical patient care in Spain. Part 1: Trauma of the central nervous system[☆]



Encuesta de atención al paciente neurocrítico en España. Parte 1: Traumatismos del sistema nervioso central

Sir,

With the aim of minimizing variability in critical patient care, the Spanish Society of Intensive and Critical Care Medicine and Coronary Units (*Sociedad Española de Medicina Intensiva, Crítica y Unidades Coronarias* [SEMICYUC]) has recently published its recommendations on “what to do” and “what not to do”, endorsed by the corresponding Working Groups (WGs).^{1,2} The management of patients with central nervous system (CNS) disease is a clear example reflecting the need to standardize the treatment protocols.³

From the Neurointensive and Trauma Care WG of the SEMICYUC, we decided to conduct a survey of neurocritical patient care with the aim of knowing the characteristics of the Departments of Intensive Care Medicine (DICMs) that manage such patients and the availability of the different techniques, as well as of analyzing certain controversial aspects in the clinical management of the disorders involved (traumatic brain injury [TBI], vertebral and spinal

cord injury, non-traumatic subarachnoid hemorrhage [SAH] and acute hemorrhagic and ischemic cerebrovascular disease [ACVD]). Scientific endorsement of the SEMICYUC was obtained. No approval from the Clinical Research Ethics Committee was considered necessary, in view of the voluntary and anonymous nature of participation in the study, which moreover did not involve the compilation of patient information. The survey was submitted to the Heads of Department on 7 November 2018, allowing an open reply period of three months. In the event of a duplicate response, that obtained first was entered in the analysis of results. The data obtained were subjected to descriptive statistical analysis, with calculation of the number (percentage) or median (interquartile range [IQR]). The present study reports the findings referred to patients with CNS trauma.

We received 45 replies with four duplicates; the valid responses thus totaled 41 (response rate 22.3%). The responding centers are indicated in Annex A (Supplementary material). The survey was answered by Heads of Department (36.6%), Section chiefs (19.5%) and staff physicians (43.9%). The hospitals were largely public (90.9%), university-related (85.4%), reference centers in Neurosurgery (80.5%) and of third level (61%), with training of residents in Intensive Care Medicine (75.6%). The median (IQR) number of hospital beds was 650 (480), while the number of DICM beds was 19 (14). A total of 68.3% of the centers monitored quality indicators.

The availability of imaging techniques and interventional radiology resources is summarized in Table 1. With regard to point-of-care techniques, the availability in the participating centers was: intracranial pressure (ICP) (80.5%), jugular vein oxygen saturation (46.3%), tissue pressure (53.7%), transcranial doppler/duplex (90.2%), near-infrared spectroscopy (26.8%), intermittent (92.7%) or continuous electroencephalography (17.1%) and cerebral microdialysis (4.9%). With regard to clinical management, 61% of the

[☆] Please cite this article as: Llopart-Pou JA, Barea-Mendoza JA, Pérez-Bárcena J, Sánchez-Casado M, Ballesteros-Sanz MA, Chico-Fernández M et al. Encuesta de atención al paciente neurocrítico en España. Parte 1: Traumatismos del sistema nervioso central. *Med Clin (Barc)*. 2021;45:250–252.