

Reply to "Pulmonary toxicity by oxygen and COVID-19"[☆]



Respuesta a «Toxicidad pulmonar por oxígeno y COVID-19»

Dear Editor,

We read with great interest the letter in which the authors talk about the deleterious effect of using a high fraction of inspired oxygen (FiO₂) in patients with acute respiratory distress syndrome (ARDS) due pneumonia caused by SARS-CoV-2 where they advocate for the use of continuous positive airway pressure (CPAP) to reduce the high FiO₂ used in high-flow nasal oxygen (HFNO) therapy.¹

We agree with the authors when they talk about planning CPAP or non-invasive ventilation (NIV) as an alternative to HFNO. However, clinical practice guidelines say otherwise and they recommend HFNO and choose invasive mechanical ventilation (IMV) in cases of failed HFNO.² Based on these recommendations,² a Spanish multicenter registry (n=876 patients) showed a wide use of HFNO (49%) vs NIV, and CPAP (<5% in both modalities). We should mention the high rate of failure in the HFNO group (60%) with mortality rates > 30% in patients who required intubation.³

The efficacy of NIV in SARS-CoV-2-induced pneumonia was demonstrated in an Italian multicenter registry (N=110 patients) that compared the NIV-helmet to HFNO. Although the registry primary endpoint—ventilation-free days (20 days vs 18 days, *P*=.26)—was not achieved the NIV group had lower rates of intubation on day 28 vs the HFNO group [28% vs 51%; OR, 0.37 (0.17–0.82), *P*=.02]. At the same time, the NIV-helmet had better oxygenation, and less dyspnea compared to the HFNO group.⁴ Consistent with one of the arguments that support the use of HFNO,² the patients' tolerance was higher with HFNO compared to NIV.⁴ Similarly, in our series of 27 hypoxemic patients due to SARS-CoV-2-induced pneumonia, NIV was used as the first-line therapy in 21 patients (80.8%) basically with NIV specific ventilators in CPAP mode. CPAP failed in 10 patients (48%) with an associated mortality rate of 50%. No health personnel became infected in relation to the NIV as opposed to what has been reported in the clinical practice guidelines.²

Therefore, based on these guidelines,² we believe that the HFNO plays a key role in the early ventilatory therapy of hypoxemic patients. Unfortunately, the rate of failure

in hypoxemic patients is high,³ which may have overestimated the true efficacy of HFNO in advanced stages of ARDS. Like the authors say, a high FiO₂ in HFNO added to the pulmonary damage caused by the virus worsens ARDS, and eventually leads to IMV.¹ On the contrary, results from the latest clinical trials,³ and observational studies⁴ pave the way for the safe use of NIV in its different modes (CPAP or NIV) by applying positive end-expiratory pressure (PEEP) that recruits the damaged lung, which allows reaching a non-detrimental FiO₂¹ while avoiding IMV and its deleterious effect.^{3,5}

References

1. León-Jiménez A, Vázquez-Gandullo E, Montoro-Ballesteros F. Toxicidad pulmonar por oxígeno y COVID-19. *Med Intensiva*. 2020. <http://dx.doi.org/10.1016/j.medin.2021.04.008>.
2. Alhazzani W, Møller MH, Arabi YM, Loeb M, Gong MN, Fan E, et al. Surviving Sepsis Campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). *Intensive Care Med*. 2020;46:854–87.
3. Fernández R, González de Molina FJ, Batlle M, Fernández MM, Hernández S, Villagra A, Grupo de Semicríticos Covid. Soporte ventilatorio no invasivo en pacientes con neumonía por COVID-19: un registro multicéntrico español. *Med Intensiva*. 2020. <http://dx.doi.org/10.1016/j.medin.2021.02.002>.
4. Grieco DL, Menga LS, Cesarano M, Rosà T, Spadaro S, Bitondo MM, et al. Effect of helmet noninvasive ventilation vs high-flow nasal oxygen on days free of respiratory support in patients with COVID-19 and moderate to severe hypoxemic respiratory failure. *JAMA*. 2021;325:1731–43.
5. Belenguer-Muncharaz A, Hernández-Garcés H, López-Chicote C, Ribes-García S, Ochagavía-Barbarín J, Zaragoza-Crespo R. Eficacia de la ventilación no invasiva en pacientes ingresados por neumonía por SARS-CoV-2 en una unidad de cuidados intensivos. *Med Intensiva*. 2020. <http://dx.doi.org/10.1016/j.medin.2020.08.011>.

A. Belenguer-Muncharaz^{a,b,*}, H. Hernández-Garcés^a

^a Servicio de Medicina Intensiva, Hospital Universitario Dr. Peset, Valencia, Spain

^b Unidad Predepartamental Medicina, Facultad de Ciencias de la Salud, Universitat Jaume I (UJI), Castelló de la Plana, Spain

* Corresponding author.

E-mail address: belengualberto8@gmail.com (A. Belenguer-Muncharaz).

<https://doi.org/10.1016/j.medine.2022.04.005>
2173-5727/ © 2022 Published by Elsevier España, S.L.U.

[☆] Please cite this article as: Belenguer-Muncharaz A, Hernández-Garcés H. Respuesta a «Toxicidad pulmonar por oxígeno y COVID-19». *Med Intensiva*. 2022;46:354.