



IMAGES IN INTENSIVE MEDICINE

Reverse triggering Trigger reverso



V. Yordanov Zlatkov^{a,*}, C. Mayordomo García^b, A. Núñez Reiz^a

^a Servicio de Medicina Intensiva, Hospital Clínico San Carlos, Madrid, Spain

^b Servicio de Medicina Intensiva, Hospital Clínico Lozano Blesa, Zaragoza, Spain

Available online 13 August 2022



Figure 1

DOI of original article: <https://doi.org/10.1016/j.medin.2021.11.004>

* Corresponding author.

E-mail address: viktoryz1111@hotmail.com (V. Yordanov Zlatkov).

<https://doi.org/10.1016/j.medic.2022.07.004>

2173-5727/© 2021 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.

The image (Fig. 1) shows a registry of the electrical activity of the diaphragm using a neurally adjusted ventilatory assist (NAVA) esophageal catheter in a patient being ventilated in pressure control mode. The figure at the bottom superimposes two registries: pressure-time (yellow tracing) and electrical activity-time (grey tracing).

All inspirations are mandatory (yellow arrows). However, during each mandatory inspiration, the patient experiences spontaneous inspiration reflected by an increase in electrical activity of the diaphragm (grey arrows). This event coincides in time with a series of negative inflections in the pressure registry (white asterisks).

The registry corresponds to a patient with acute respiratory distress syndrome who suffers reverse triggering.

Conflicts of interest

None.