



ORIGINAL ARTICLE

Correlation and validity of imputed PaO₂/FiO₂ and SpO₂/FiO₂ in patients with invasive mechanical ventilation at 2600 m above sea level



G. Ortiz^a, A. Bastidas^{b,*}, M. Garay-Fernández^c, A. Lara^c, M. Benavides^c, E. Rocha^d, A. Buitrago^c, G. Díaz^c, J. Ordóñez^c, L.F. Reyes^b

^a Pulmonary Medicine, Universidad El Bosque, Intensive Care Unit, Hospital Santa Clara Bogotá, Colombia

^b School of Medicine, Universidad de la Sabana, Clínica Universidad de La Sabana, Chía, Colombia

^c Pulmonary Medicine Universidad El Bosque, Intensive Care Unit, Hospital Santa Clara Bogotá, Colombia

^d Intensive Care Unit, Hospital Santa Clara Bogotá, Colombia

Received 6 February 2021; accepted 4 May 2021

KEYWORDS

Mechanical ventilation;
SaO₂/FiO₂;
PaO₂/FiO₂;
Respiratory failure;
High altitude

Abstract

Objective: To establish the correlation and validity between PaO₂/FiO₂ obtained on arterial gases versus noninvasive methods (linear, nonlinear, logarithmic imputation of PaO₂/FiO₂ and SpO₂/FiO₂) in patients under mechanical ventilation living at high altitude.

Design: Ambispective descriptive multicenter cohort study.

Setting: Two intensive care units (ICU) from Colombia at 2600 m a.s.l.

Patients or participants: Consecutive critically ill patients older than 18 years with at least 24 h of mechanical ventilation were included from June 2016 to June 2019.

Interventions: None.

Variables: Variables analyzed were demographic, physiological measures, laboratory findings, oxygenation index and clinical condition. Nonlinear, linear and logarithmic imputation formulas were used to calculate PaO₂ from SpO₂, and at the same time the SpO₂/FiO₂ by severe hypoxemia diagnosis. The intraclass correlation coefficient, area under the ROC curve, sensitivity, specificity, positive predictive value, negative predictive value, positive and negative likelihood ratio were calculated.

Results: The correlation between PaO₂/FiO₂ obtained from arterial gases, PaO₂/FiO₂ derived from one of the proposed methods (linear, non-linear, and logarithmic formula), and SpO₂/FiO₂

Abbreviations: ARI, acute respiratory insufficiency; ARDS, acute respiratory distress syndrome; AUC-ROC, area under the curve receiver operating characteristic; CI, confidence interval; ICC, intraclass correlation coefficient; FiO₂, the fraction of inspired oxygen; MSNM, meters above sea level; PaCO₂, partial pressure of arterial carbon dioxide; PaO₂, partial pressure of arterial oxygen; PEEP, end-expiratory pressures; SOFA, sequential multiple organ damage assessment.

* Corresponding author.

E-mail address: alirio.bastidas@unisabana.edu.co (A. Bastidas).

measured by the intraclass correlation coefficient was high (greater than 0.77, $p < 0.001$). The different imputation methods and SpO₂/FiO₂ have a similar diagnostic performance in patients with severe hypoxemia (PaO₂/FiO₂ <150). PaO₂/FiO₂ linear imputation AUC ROC 0,84 (IC 0.81–0.87, $p < 0.001$), PaO₂/FiO₂ logarithmic imputation AUC ROC 0.84 (IC 0.80–0.87, $p < 0.001$), PaO₂/FiO₂ non-linear imputation AUC ROC 0.82 (IC 0.79–0.85, $p < 0.001$), SpO₂/FiO₂ oximetry AUC ROC 0.84 (IC 0.81–0.87, $p < 0.001$).

Conclusions: At high altitude, the SaO₂/FiO₂ ratio and the imputed PaO₂/FiO₂ ratio have similar diagnostic performance in patients with severe hypoxemia ventilated by various pathological conditions.

© 2021 Elsevier España, S.L.U. y SEMICYUC. All rights reserved.

PALABRAS CLAVE

Ventilación mecánica;
SaO₂/FiO₂;
PaO₂/FiO₂;
Insuficiencia
respiratoria;
Altitud elevada

Correlación y validez de la PaO₂/FiO₂ y SpO₂/FiO₂ imputadas en pacientes con ventilación mecánica invasiva a 2.600 metros sobre el nivel del mar

Resumen

Objetivo: Establecer la correlación y validez entre PaO₂/FiO₂ obtenida en gases arteriales versus métodos no invasivos (imputación lineal, no lineal, logarítmica de PaO₂/FiO₂ y SpO₂/FiO₂) en pacientes bajo ventilación mecánica que viven en altitudes elevadas.

Diseño: Estudio de cohorte multicéntrico descriptivo ambispectivo

Ámbito: Dos unidades de cuidados intensivos de Colombia a 2.600 m s.n.m.

Pacientes o participantes: Se incluyeron pacientes consecutivos en estado crítico mayores de 18 años con al menos 24 h de ventilación mecánica desde junio de 2016 a junio de 2019.

Intervenciones: Ninguna.

Variables: Las variables analizadas fueron demográficas, fisiológicas, hallazgos de laboratorio, índice de oxigenación y estado clínico. Se utilizaron fórmulas de imputación no lineales, lineales y logarítmicas para calcular la PaO₂ a partir de la SpO₂, y al mismo tiempo la SpO₂/FiO₂ mediante el diagnóstico de hipoxemia severa. Se calculó el coeficiente de correlación intraclass, el área bajo la curva ROC, la sensibilidad, la especificidad, el valor predictivo positivo, el valor predictivo negativo, la razón de verosimilitud positiva y negativa.

Resultados: La correlación entre PaO₂/FiO₂ obtenida a partir de gases arteriales, PaO₂/FiO₂ derivada de uno de los métodos propuestos (fórmula lineal, no lineal y logarítmica) y SpO₂/FiO₂ medida por el coeficiente de correlación intraclass fue alta (mayor a 0,77, $p < 0,001$). Los diferentes métodos de imputación y SpO₂/FiO₂ tienen un rendimiento diagnóstico similar en pacientes con hipoxemia severa (PaO₂/FiO₂ < 150). PaO₂/FiO₂ imputación lineal AUC ROC 0,84 (IC 0,81-0,87; $p < 0,001$), PaO₂/FiO₂ imputación logarítmica AUC ROC 0,84 (IC 0,80-0,87; $p < 0,001$), PaO₂/Imputación no lineal de FiO₂ AUC ROC 0,82 (IC 0,79-0,85; $p < 0,001$), oximetría de SpO₂/FiO₂ AUC ROC 0,84 (IC 0,81-0,87; $p < 0,001$).

Conclusiones: A gran altitud, el cociente SaO₂/FiO₂ y el cociente PaO₂/FiO₂ imputado tienen un rendimiento diagnóstico similar en pacientes con hipoxemia severa bajo ventilación mecánica invasiva por diversas patologías.

© 2021 Elsevier España, S.L.U. y SEMICYUC. Todos los derechos reservados.

Introduction

Acute respiratory insufficiency (ARI) is diagnosed in 11.3% of hospitalizations in the United States. Its incidence calculated for 2001 was 502 cases per 100,000 inhabitants, increasing to 704 cases per 100,000 inhabitants in 2009.¹ The relationship between the pressure of oxygen dissolved in the blood (PaO₂) and the fraction of inspired oxygen (FiO₂) (PaO₂/FiO₂), an index that classifies the severity of the disease in patients with acute respiratory failure, was incorporated in 2011 in the Berlin Consensus definition of adult acute respiratory distress syndrome (ARDS)² and has

been included in severity scores, such as in the sequential multiple organ damage assessment (SOFA) scale.^{3,4}

The measurement of arterial blood gases is required to determine PaO₂ and calculate the PaO₂/FiO₂ ratio.^{5,6} However, this technique can present limitations in its use, either because of difficulty in taking the sample (either because it requires trained personnel, complicated arterial accesses, constant puncture requirements, pain at the time of sample collection and skin laceration) or because it represents an increase in hospital costs.^{7,8}

A non-invasive substitute for the PaO₂/FiO₂ ratio is the ratio of the percentage of hemoglobin saturation measured

by pulse oximetry (SpO₂) and the inspired fraction of oxygen (FiO₂) (SpO₂/FiO₂).⁹ That allows the assessment and classification of the severity of respiratory failure¹⁰ without the need for arterial blood sampling.¹¹ Other methods have been proposed to estimate PaO₂ without direct measurement of oxygen in arterial blood. These models ensure their equivalence by imputing PaO₂ from SpO₂. These imputation techniques can be classified into those that simulate the hemoglobin dissociation curve (non-linear)^{12,13} and those that take into account other types of relationships (linear and logarithmic).^{14,15} The studies that have compared these formulas show an adequate relationship between measured and imputed PaO₂ values, with high correlation coefficients ranging from 0.75 to 0.9.^{12,13,15}

It must be taken into account that about 140 million people live at high altitudes (over 2500 m above sea level), which is why it is essential to understand how the oxygenation indices (PaO₂/FiO₂ and SpO₂/FiO₂) behave in these regions.¹⁶ However, we do not know of previous studies conducted in populations living at altitudes above 1500 m a.s.l., which evaluate the performance of different imputation methods and SpO₂/FiO₂ for the diagnosis of severe hypoxemia. Therefore, we aimed to establish the correlation and validity between PaO₂/FiO₂ obtained in arterial gases vs. non-invasive methods (linear, non-linear, logarithmic imputation of PaO₂/FiO₂ and SpO₂/FiO₂) in patients living at 2600 a.s.l. and under invasive mechanical ventilation.

Methodology

Ambispective cohort study was conducted in patients hospitalized in intensive care units with invasive ventilatory support in two third-level care hospitals in Colombia, the Santa Clara Hospital in the city of Bogota (Altitude: 2640 m above sea level) and the Clínica Universidad de La Sabana in the city of Chia (Altitude: 2562 m above sea level). The data was initially collected retrospectively, from June 2016 to April 2019, and prospectively from April to June 2019; information was obtained from the electronic medical records from the period of hospitalization.

Patients

Subjects over 18 years old, with at least 24h of invasive mechanical ventilation from any cause, with simultaneous arterial gases and SpO₂ measurements, were included, regardless of the saturation value, severity of the disease, type of ventilatory mode, vasopressor support, or radiological involvement. Patients without concomitant measurements of arterial gases, SpO₂ and ventilatory parameters were excluded. Patients with terminal disease, extracorporeal membrane oxygenation and those in whom FiO₂, PaO₂, and SpO₂ data had not been reported in the clinical records or had bad quality in the pulse wave were also excluded.

Variables

Information was collected on age, sex, skin pigmentation, type of pathology of entry, weight, height, vital signs,

values of hemoglobin, hematocrit, bilirubin, and creatinine, findings on chest radiography, with verification of the presence of infiltrates and the number of quadrants involved, SOFA severity score, vasopressor support, ventilatory support parameters (PEEP level >10 and <10 cmH₂O,^{17,18} peak pressure, plateau pressure, tidal volume), total arterial gas values and oxygenation rates, days of mechanical ventilation, intensive care unit (ICU) and hospital stays. Severe hypoxemia was the value of the PaO₂/FiO₂ ratio obtained from arterial gases less than 150.^{19,20}

Severinghaus-Ellis (no linear),^{21,22} Rice (linear)¹⁴ and Pandharipande (logarithmic linear)¹⁵ formulas were used to calculate imputed PaO₂ from SpO₂. The values of imputed PaO₂ were used for the calculation of imputed PaO₂/FiO₂; besides, SpO₂/FiO₂ was calculated simultaneously. The blood sample for arterial gases analysis was obtained through an arterial line; the oximeters used measure the saturation through an infrared sensor with a wavelength of 905 nm 2.0 mW with an error range of 1%; verification and calibration were performed daily during the study.

All the subjects admitted to the ICU of both institutions during the study period were included. In order to reduce errors in data collection and transcription, the data were recorded by double typing and reviewed by two members of the research team. This protocol was approved by the ethics committee of the Clínica Universidad de La Sabana.

Statistical analysis

An initial descriptive analysis was carried out, summarizing the qualitative variables in frequencies and percentages, the quantitative variables with normal distribution in means and standard deviation and if they did not meet the criteria of normality in medians and interquartile ranges. The qualitative variables were compared through the chi-square test and the quantitative ones with Student's t-test or Mann-Whitney *U* test according to their distribution.

We applied the intraclass correlation coefficient between the values obtained from imputed PaO₂, PaO₂/FiO₂ derived from the imputation methods, and the SpO₂/FiO₂ with the PaO₂/FiO₂ of the arterial gases. We considered the correlations thus: 0–0.3 null; 0.31–0.50 low; 0.51–0.70 moderate; 0.76–1.00 strong. The Bland-Altman method was used to evaluate the concordance. For the validity, the area under the ROC curve of the quantitative values obtained through the three imputation methods and the SpO₂/FiO₂ value was calculated, comparing it with the value of the arterial gases of severe hypoxemia (PaO₂/FiO₂ ≤ 150), to obtain the best cut-off point using the Youden index and calculating the values of sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), positive and negative likelihood ratio. The ROC area was also used to establish the equivalence ranges between the different SpO₂/FiO₂ measurements. A *p*-value of less than 0.05 was considered significant. The statistical analysis was performed with the SPSS 22.0 licensed program (Statistical Package for the Social Sciences, Chicago, IL).

Table 1 General characteristics of the cohort.

Characteristics	General population <i>n</i> = 664		PEEP < 10 cmH ₂ O <i>n</i> = 438		PEEP > 10 cmH ₂ O <i>n</i> = 226		<i>p</i> -Value
Age (years) (SD)	56.0	(19.9)	55.8	(20.1)	56.6	(19.5)	0.603
Sex (male), <i>n</i> (%)	444	(0.6)	291	(0.6)	153	(0.6)	0.744
<i>Condition on admission n (%)</i>							
Elective surgery	54	(0.0)	41	(0.1)	13	(0.0)	0.272
Emergency surgery	75	(0.1)	49	(0.1)	26	(0.1)	
Non-surgical	535	(0.8)	348	(0.7)	187	(0.8)	
Height cm (SD)	163.0	(9.0)	162.9	(9.1)	163.0	(8.8)	0.876
Weight kg (SD)	66.8	(157.2)	66.6	(12.4)	67.3	(12.8)	0.493
Vital signs (DE)	87.4	(17.7)	87.3	(18.0)	87.5	(17.0)	0.862
SBP mmHg	119.0	(24.0)	119.1	(24.4)	119.1	(23.2)	0.856
DBP mmHg	71.6	(17.0)	71.4	(17.4)	71.9	(16.3)	0.493
MAP mmHg	87.4	(17.7)	87.3	(18.0)	87.5	(17.0)	0.862
HR beats × min	86.8	(21.3)	84.9	(20.6)	90.6	(22.2)	0.001
RR resp × min	18.8	(4.7)	18.6	(4.7)	19.2	(4.7)	0.175
Temperature	36.9	(0.7)	36.8	(0.6)	36.9	(0.8)	0.037
<i>Laboratory workup (SD)</i>							
Leukocytes cell/ml	11,979.7	(7224.7)	12,145.1	(6975.9)	11,659.0	(7689.8)	0.426
Hb (g/dL)	11.90	(3.0)	11.79	(3.0)	12.11	(3.0)	0.200
Hematocrit (%)	36.4	(9.0)	36.2	(9.0)	37.0	(9.0)	0.280
Platelet count cell/ml	231,737.6	(102,667.8)	233,050.3	(98,410.4)	229,199.9	(110,618.9)	0.659
Total bilirubin (mg/dL)	1.27	(1.9)	1.22	(1.7)	1.34	(2.2)	0.483
Creatinine (mg/dL)	1.40	(1.4)	1.40	(1.5)	1.39	(1.3)	0.920
Sodium (mEq)	143.5	(8.5)	143.3	(9.1)	143.9	(7.1)	0.327
Potassium (mEq)	3.97	(0.8)	3.97	(0.7)	3.97	(0.8)	0.917
<i>Infiltrates on chest X-ray Quadrants involved n(%)</i>							
1	111	(0.3)	41	(0.3)	70	(0.4)	
2	127	(0.4)	43	(0.3)	84	(0.4)	
3	21	(0.0)	14	(0.1)	7	(0.0)	
4	31	(0.1)	22	(0.1)	9	(0.0)	
SOFA (IQR)	15	(2)	15	(2)	16	(2)	0.019
Vasopressor support	375	(0.5)	242	(0.5)	133	(0.5)	0.376

Notes: SD: standard deviation, SBP: systolic blood pressure, DBP: diastolic blood pressure, MAP: mean arterial pressure, HR: heart rate, RR: respiratory rate, Hb: hemoglobin, SOFA: Sepsis-related Organ Failure Assessment, IQR: interquartile range.

Results

We included a total of 664 patients during the study period, the general characteristics of the patients were described according to positive end-expiratory pressures (PEEP) greater or less than 10 cm H₂O and are showed in Table 1. Among the general characteristics, it was found that the patients had an mean age of 56 ± 20 years, with a male predominance 444/664 (67%), most of the pathological conditions on admission were non-surgical (81%), the mean SOFA was 15 ± 2 points, and 56% of the patients required vasoactive support.

Arterial gases and mechanical ventilation

The mean PaO₂ was 83.7 ± 27.5 mmHg, SpO₂ 92.4 ± 2.3%, PaCO₂ 33 ± 2.9 mmHg, and bicarbonate 21.7 ± 1.6 mEq/L, with no significant differences between subgroups. The pH was 7.35 ± 0.11 and the base excess -1.3 ± 1.4 mEq/L. The ventilatory parameters were a tidal volume of 481 ± 58.3 ml,

FIO₂ of 50 ± 19%, plateau pressure 19.3 ± 4.3 cmH₂O, and driving pressure of 10.8 cmH₂O. The mean time of mechanical ventilation was 7.1 days, ICU length of stay was 10.4 days and overall mortality was 18%. ([Supplementary Material](#))

Correlation and validity results

The correlation between PaO₂/FiO₂ obtained from arterial gases, PaO₂/FiO₂ derived from one of the proposed methods (linear, non-linear, and logarithmic formula), and SpO₂/FiO₂ measured by the intraclass correlation coefficient was high (greater than 0.77, *p* < 0.001). The different imputation methods and SpO₂/FiO₂ have a similar diagnostic performance in patients with severe hypoxemia (PaO₂/FiO₂ < 150), with an area under the ROC curve greater than 0.8 with a similar sensitivity and specificity, PaO₂/FiO₂ linear imputation AUC ROC 0.84 (IC 0.81–0.87, *p* < 0.001), PaO₂/FiO₂ logarithmic imputation AUC ROC 0.84 (IC 0.80–0.87, *p* < 0.001), PaO₂/FiO₂ non-linear imputation

Table 2 Results of validity between the formulas of imputation of PaO₂, PaO₂/FiO₂, SpO₂/FiO₂ and PaO₂, PaO₂/FiO₂ of arterial blood gases and severe Hypoxemia PaO₂/FiO₂ ≤ 150.

Method	AUC-ROC	CI 95%	Cut-off point	SENS	SPEC	PPV	NPV	LR+	LR-	p-Value
PaO ₂ linear imputation	0.83	(0.80, 0.86)	73.1	0.65	0.83	0.19	0.81	3.88	0.20	<0.001
PaO ₂ logarithmic imputation	0.84	(0.81, 0.87)	50.2	0.67	0.82	0.20	0.82	3.71	0.22	<0.001
PaO ₂ non-linear imputation	0.60	(0.55, 0.64)	70.8	0.40	0.79	0.13	0.71	1.86	0.27	<0.001
PaO ₂ /FiO ₂ linear imputation	0.84	(0.81, 0.87)	147.6	0.67	0.82	0.20	0.82	3.71	0.22	<0.001
PaO ₂ /FiO ₂ logarithmic imputation	0.84	(0.80, 0.87)	119.1	0.62	0.87	0.19	0.81	4.64	0.15	<0.001
PaO ₂ /FiO ₂ non-linear imputation	0.82	(0.79, 0.85)	141.3	0.65	0.83	0.19	0.81	3.87	0.20	<0.001
SpO ₂ /FiO ₂ oximetry	0.84	(0.81, 0.87)	204.4	0.61	0.87	0.18	0.81	4.62	0.15	<0.001

Notes: SENS: sensitivity; SPEC: specificity; AUC-ROC: area under the ROC curve; PPV: positive predictive values; NPV: negative predictive values; LR: likelihood ratio.




Figure 1 ROC curve with yield of PaO₂/FiO₂ imputed by the three methods of imputation and SpO₂/FiO₂.

Notes: AUC-ROC: area under the ROC-curve, PaO₂/FiO₂ linear imputation 0.84 (95%CI: 0.81–0.87); PaO₂/FiO₂ logarithmic imputation 0.84 (95%CI: 0.80–0.87); PaO₂/FiO₂ non-linear 0.82 (95%CI: 0.79–0.85); SpO₂/FiO₂ pulse oximetry 0.84 (95%CI: 0.81–0.87).

AUC ROC 0.82 (IC 0.79–0.85, *p* < 0.001), SpO₂/FiO₂ oximetry AUC ROC 0.84 (IC 0.81–0.87, *p* < 0.001). Table 2 and Fig. 1.

We found that the correlation of the different methods applied (imputation and SpO₂/FiO₂) decreased with hemoglobin levels equal to or less than 7g/dl (0.58 *p* < 0.001), as well as with total bilirubin values equal to or greater than 3mg/dL (0.68 *p* < 0.001), the results above concordance between imputed PaO₂/FiO₂ and SpO₂/FiO₂ with Bland–Altman graph, and the validity for

physiologic variable, mechanical ventilation and radiological findings between SpO₂/FiO₂ and severe Hypoxemia PaO₂/FiO₂ ≤ 150 showed in Supplementary Material.

In a practical way and to establish the usefulness of SpO₂/FiO₂ measurement, we present the ranges of values between SpO₂/FiO₂ and PaO₂/FiO₂ imputed by the three methods and PaO₂/FiO₂ measured by arterial gases in Table 3.

Discussion

This study found an adequate correlation between PaO₂/FiO₂ obtained by the three proposed imputation methods, SpO₂/FiO₂, and PaO₂/FiO₂ values obtained by arterial blood gases, considering the usefulness of these tools as non-invasive methods for the assessment of the oxygenation status in patients with invasive ventilatory support. We also found that the three imputation methods evaluated and the SpO₂/FiO₂ have an adequate diagnostic performance for severe hypoxemia (PaO₂/FiO₂ ≤ 150 mmHg by arterial gases) in patients living at high altitude.

It was previously determined that the concordance between the different non-invasive oxygenation indexes and the PaO₂/FiO₂ obtained by arterial gases is high^{7,16,23,24} Cineci and Gomez found an overall correlation index of 0.745 between SpO₂/FiO₂ and PaO₂/FiO₂ in patients with ARF¹¹; Rice et al. using imputation formulas, report a correspondence range between 0.73 and 0.88^{12,14,15} that are similar to those observed in our study (greater than 0.77). This accuracy can be influenced by the use of PEEP, hemoglobin levels, total bilirubin values, by the determinants of the affinity of oxygen for hemoglobin (pH, temperature, CO₂, 2.3-diphosphoglycerate, and fetal hemoglobin)^{12,15} and extreme values of arterial saturation,^{6,25} which can vary in critically-ill ventilated patients. Our results show an important decrease in the

Table 3 Values of correlation of SaO₂/FiO₂ by pulse oximetry and PaO₂/FiO₂ by three imputation techniques and arterial blood gases in patients under mechanical ventilation by ROC curve.

SpO ₂ /FiO ₂	PaO ₂ /FiO ₂ linear imputation		PaO ₂ /FiO ₂ logarithmic imputation		PaO ₂ /FiO ₂ non-linear imputation		PaO ₂ /FiO ₂ arterial blood gases		
>300	297	328	247	284	203	323	246	324	
250	299	250	297	149	247	149	203	225	246
200	249	148	250	100	149	129	149	182	225
150	199	92	148	64	149	98	129	137	182
100	149	37	92	30	64	68	98	100	137
<100	26	37	27	30	51	68	62	100	

correlation in subjects with hyperbilirubinemia (total bilirubin ≥ 3 mg/dL) and hemoglobin less than ≤ 7 mg/dL, without it being significantly affected by temperature, vasoactive support, and severity of multiorgan involvement.

Theoretically, imputation techniques that simulate the hemoglobin dissociation curve have a better diagnostic yield with the patient's oxygenation status. Brown's^{12,13} and Gadrey's²⁴ studies found that the diagnosis of moderate-to-severe hypoxemia is better when non-linear formulas are used. Despite this, our data show similar performance among the different imputation methods, without important changes in their diagnostic capacity when patients with arterial oxygen saturation greater than 97% are included. This difference is probably explained by the type of population studied and the barometric pressure to which our patients are exposed. It is important to emphasize that our results are aimed at the recognition of severe hypoxemia and not hyperoxemia, where the subrogated SpO₂ values may be more limited.²⁵

Although mechanical ventilators represent a closed circuit, there are not pressurized; therefore, patients ventilated at high altitude may have lower PaO₂ and SaO₂ values than those obtained at sea level.^{6,26,27} Previous observations show that the corresponding values of SaO₂/FiO₂ from 214 to 235 are equivalent to a PaO₂/FiO₂ (for arterial gases) of 200 in inhabitants of low-altitude areas.^{11,15} However, there are no studies in patients under invasive mechanical ventilation living at more than 1500 m a.s.l. that establish a single SpO₂/FiO₂ value corresponding to a given PaO₂/FiO₂. This limitation also affects the values obtained by imputation methods, since it is not possible to establish values that correspond to these measurements without any margin of error.¹³ Considering these difficulties, we propose a table of equivalence between SpO₂/FiO₂ ranges and PaO₂/FiO₂ values imputed by the three methods and that obtained by arterial gases, to establish values for the diagnosis and follow-up of patients with different degrees of hypoxemia that may be useful in clinical practice. We must not forget to evaluate these considering medical criteria and the condition of the patient.

We acknowledge that our study has certain limitations. First, a single arterial gas sampling was performed, which can affect the reliability of the measured values of the quantitative variables for the diagnosis of hypoxemia and the appearance of random errors. However, the samples were taken by trained personnel and keeping all the recommendations for adequate processing in the laboratory.

Second, some data were collected retrospectively, which can generate an information bias; however, we verified that the values obtained from the clinical histories corresponded to the data reported directly by the laboratory. Third, despite having a large population living at high altitude, our sample does not include all possible altitudes; but we consider that it is unlikely that there are significant differences in the reliability of the formulas. Fourth, SpO₂ values can be affected by methemoglobin and carboxyhemoglobin levels, which were not measured in the study. Despite this, none of the patients were suspected to have these disorders.

In conclusion, the present study found that at high altitude SaO₂/FiO₂ and imputed PaO₂/FiO₂ have a similar diagnostic yield in patients with severe hypoxemia, ventilated for various pathological conditions. Equivalence ranges between SaO₂/FiO₂ and PaO₂/FiO₂ values can be a non-invasive option in the follow-up of these subjects. It is necessary to develop additional prospective studies, where therapeutic goals of SpO₂/FiO₂ and imputed PaO₂/FiO₂ are explored to corroborate the safety ranges of these measurements.

Funding

The author received no financial support for the research, authorship, and/or publication of this article

Author contributions

GO, AB, MB, MGF and AL had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. GO, AB, MB, MGF, AL, GD, JO contributed substantially to the study design, data analysis and interpretation, and the writing of the manuscript. GO, AB contributed substantially to data statistical analysis.

Conflict of interests

The authors of this manuscript have no competing interests directly related to the manuscript's content.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.medin.2021.05.001](https://doi.org/10.1016/j.medin.2021.05.001).

References

1. Stefan MS, Shieh MS, Pekow PS, Rothberg MB, Steingrub JS, Lagu T, et al. Epidemiology and outcomes of acute respiratory failure in the United States, 2001 to 2009: a national survey. *J Hosp Med.* 2013;8:76–82.
2. Ferguson ND, Fan E, Camporota L, Antonelli M, Anzueto A, Beale R, et al. The Berlin definition of ARDS: an expanded rationale, justification, and supplementary material. *Intensive Care Med.* 2012;38:1573–82.
3. Moreno R, Vincent JL, Matos R, Mendonça A, Cantraine F, Thijs L, et al. The use of maximum SOFA score to quantify organ dysfunction/failure in intensive care Results of a prospective, multicentre study. Working Group on Sepsis related Problems of the ESICM. *Intensive Care Med.* 1999;25:686–96.
4. Seymour CW, Liu VX, Iwashyna TJ, Brunkhorst FM, Rea TD, Scherag A, et al. Assessment of clinical criteria for sepsis: for the third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA.* 2016;315:762–74.
5. Räsänen J, Downs JB, Malec DJ, DeHaven B, Garner WL. Real-time continuous estimation of gas exchange by dual oximetry. *Intensive Care Med.* 1988;14:118–22.
6. Karbing DS, Kjaergaard S, Smith BW, Espersen K, Allerød C, Andreassen S, et al. Variation in the PaO₂/FiO₂ ratio with FiO₂: mathematical and experimental description, and clinical relevance. *Crit Care.* 2007;11:R118.
7. Bashar FR, Vahedian-Azimi A, Farzanegan B, Goharani R, Shojaei S, Hatamian S, et al. Comparison of non-invasive to invasive oxygenation ratios for diagnosing acute respiratory distress syndrome following coronary artery bypass graft surgery: a prospective derivation-validation cohort study. *J Cardiothorac Surg.* 2018;13:123.
8. Chen W, Janz DR, Shaver CM, Bernard GR, Bastarache JA, Ware LB. Clinical characteristics and outcomes are similar in ARDS diagnosed by oxygen saturation/FiO₂ ratio compared with PaO₂/FiO₂ ratio. *Chest.* 2015;148:1477–83.
9. Adams JY, Rogers AJ, Schuler A, Marelich GP, Fresco JM, Taylor SL, et al. Association between peripheral blood oxygen saturation (SpO₂)/fraction of inspired oxygen (FiO₂) ratio time at risk and hospital mortality in mechanically ventilated patients. *Perm J.* 2020;24.
10. Pisani L, Roozeman JP, Simonis FD, Giangregorio A, van der Hoeven SM, Schouten LR, et al. Risk stratification using SpO₂(2)/FiO₂ and PEEP at initial ARDS diagnosis and after 24h in patients with moderate or severe ARDS. *Ann Intensive Care.* 2017;7:108.
11. Cinesi-Gómez C, García-García P, López-Pelayo I, Giménez JI, González-Torres LM, Bernal-Morell E. Correlation between oxyhaemoglobin saturation by pulse oximetry and partial pressure of oxygen in patients with acute respiratory failure. *Rev Clin Esp.* 2017;217:522–5.
12. Brown SM, Duggal A, Hou PC, Tidswell M, Khan A, Exline M, et al. Nonlinear imputation of PaO₂/FIO₂ From SpO₂/FIO₂ among mechanically ventilated patients in the ICU: a prospective, observational study. *Crit Care Med.* 2017;45:1317–24.
13. Brown SM, Grissom CK, Moss M, Rice TW, Schoenfeld D, Hou PC, et al. Nonlinear imputation of PaO₂/FiO₂ from SpO₂/FiO₂ among patients with acute respiratory distress syndrome. *Chest.* 2016;150:307–13.
14. Rice TW, Wheeler AP, Bernard GR, Hayden DL, Schoenfeld DA, Ware LB. Comparison of the SpO₂/FIO₂ ratio and the PaO₂/FIO₂ ratio in patients with acute lung injury or ARDS. *Chest.* 2007;132:410–7.
15. Pandharipande PP, Shintani AK, Hagerman HE, St Jacques PJ, Rice TW, Sanders NW, et al. Derivation and validation of SpO₂/FiO₂ ratio to impute for PaO₂/FiO₂ ratio in the respiratory component of the Sequential Organ Failure Assessment score. *Crit Care Med.* 2009;37:1317–21.
16. Batchinsky AI, Wendorff D, Jones J, Beely B, Roberts T, Choi JH, et al. Noninvasive SpO₂/FiO₂ ratio as surrogate for PaO₂/FiO₂ ratio during simulated prolonged field care and ground and high-altitude evacuation. *J Trauma Acute Care Surg.* 2020;89 2S Suppl 2:S126–31.
17. Alhazzani W, Møller MH, Arabi YM, Loeb M, Gong MN, Fan E, et al. Surviving sepsis campaign: guidelines on the management of critically ill adults with coronavirus disease 2019 (COVID-19). *Intensive Care Med.* 2020;46:854–87.
18. Briel M, Meade M, Mercat A, Brower RG, Talmor D, Walter SD, et al. Higher vs lower positive end-expiratory pressure in patients with acute lung injury and acute respiratory distress syndrome: systematic review and meta-analysis. *JAMA.* 2010;303:865–73.
19. Guérin C, Reigner J, Richard J-C, Beuret P, Gacouin A, Boulain T, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med.* 2013;368:2159–68.
20. Papazian L, Forel JM, Gacouin A, Penot-Ragon C, Perrin G, Loundou A, et al. Neuromuscular blockers in early acute respiratory distress syndrome. *N Engl J Med.* 2010;363:1107–16.
21. Severinghaus JW. Simple, accurate equations for human blood O₂ dissociation computations. *J Appl Physiol Respir Environ Exerc Physiol.* 1979;46:599–602.
22. Ellis RK. Determination of PO₂ from saturation. *J Appl Physiol (1985).* 1989;67:902.
23. Esteve F, Lopez-Delgado JC, Javierre C, Skaltsa K, Carrio ML, Rodríguez-Castro D, et al. Evaluation of the PaO₂/FiO₂ ratio after cardiac surgery as a predictor of outcome during hospital stay. *BMC Anesthesiol.* 2014;14:83.
24. Gadrey SM, Lau CE, Clay R, Rhodes GT, Lake DE, Moore CC, et al. Imputation of partial pressures of arterial oxygen using oximetry and its impact on sepsis diagnosis. *Physiol Meas.* 2019;40:115008.
25. Durlinger EMJ, Spoelstra-de Man AME, Smit B, de Grooth HJ, Girbes ARJ, Oudemans-van Straaten HM, et al. Hyperoxia: at what level of SpO₂ is a patient safe? A study in mechanically ventilated ICU patients. *J Crit Care.* 2017;39:199–204.
26. Gonzalez-Garcia M, Maldonado D, Barrero M, Casas A, Perez-Padilla R, Torres-Duque CA. Arterial blood gases and ventilation at rest by age and sex in an adult Andean population resident at high altitude. *Eur J Appl Physiol.* 2020;120:2729–36.
27. Jibaja M, Ortiz-Ruiz G, García F, Garay-Fernández M, de Jesús Montelongo F, Martínez J, et al. Hospital mortality and effect of adjusting PaO₂(2)/FiO₂ according to altitude above the sea level in acclimatized patients undergoing invasive mechanical ventilation. a multicenter study. *Arch Bronconeumol.* 2020;56:218–24.